

A NOTE ON CONSTRUCTING DIGRAPHS WITH PRESCRIBED PROPERTIES

W. Ananchuen*

School of Liberal Arts
Sukhothai Thammathirat Open University
Pakkred, Nonthaburi 11120
THAILAND

ABSTRACT

Let n be non-negative integer and k a positive integer. A digraph D is said to have property $Q(n,k)$ if for every subset of n vertices of D is dominated by at least k other vertices. For $q \equiv 5 \pmod{8}$ be a prime power. Define a quadruple Paley digraph $D_q^{(4)}$ as follows. The vertices of $D_q^{(4)}$ are the elements of the finite field F_q . Vertex a joins to vertex b by an arc if and only if $a - b = y^4$ for some $y \in F_q$. In this paper, we show for sufficiently large q , $D_q^{(4)}$ has property $Q(n,k)$.

1. INTRODUCTION

In this paper, our graphs are directed. For our purpose, all digraphs are finite and strict. If (x, y) is an arc in a digraph D , then we say vertex x dominates vertex y . A set of vertices A dominates a set of vertices B if every vertex of A dominates every vertex of B . A digraph D is said to have property $Q(n,k)$ if every subset of n vertices of D is dominated by at least k other vertices. Further, a digraph D is said to have property $Q(m,n,k)$ if for any set of $m + n$ distinct vertices of D there exist at least k other vertices each of which dominates the first m vertices and is dominated by the latter n vertices.

A special digraph arises in round robin tournaments. More precisely, consider a tournament T_q with q players $1, 2, \dots, q$ in which there are no draws. This gives rise to a digraphs in which either (a, b) or (b, a) is an arc for each pair a, b . Tournaments with property $Q(n, k)$ have been studied by Ananchuen and Caccetta [2] Bollobás [3] and Graham and Spencer [4].

* This research has been supported by The Thailand Research Fund grant BRG/07/2541.