

บทคัดย่อ

ชื่อโครงการวิจัย

การจำแนกพีชคณิตจำกัดด้วยฟังก์ชันที่ไม่สมนัยหนึ่งต่อหนึ่ง

ผู้วิจัย

รองศาสตราจารย์ ดร. ธีรวัชร รัตนประเสริฐ

คณะวิทยาศาสตร์ มหาวิทยาลัยศิลปากร e-mail address : ratach@su.ac.th

แหล่งทุนอุดหนุน

สำนักงานกองทุนสนับสนุนการวิจัย

ระยะเวลา

28 กรกฎาคม 2548 - 28 กรกฎาคม 2551

คำหลัก

unary operation, pre-period, partial operation, n-ary operation, invariant equivalence relation, clone, semigroup, band

สำหรับแต่ละ unary operation f บนเซตจำกัด A ซึ่ง $|A| = k$ จะมีโซ่ลดลง $A \supseteq \text{Im } f \supseteq \text{Im } f^2 \supseteq \dots \supseteq \text{Im } f^n = \text{Im } f^{n+1}$ โดยเรียกจำนวนเต็มบวกตัวน้อยสุด $\lambda(f)$ ที่ทำให้ $\text{Im } f^{\lambda(f)} = \text{Im } f^{\lambda(f)+1}$ ว่า pre-period ของ f และ pre-period ของ f มีค่าจาก 0 ถึง $k - 1$ ถ้า $\lambda(f) = k - 1$ เมื่อ $k \geq 1$ เราเรียก f ว่า long-tailed (LT) -operation และถ้า $\lambda(f) = k - 2$ เมื่อ $k \geq 2$ เราเรียก f ว่า (LT_1) -operation ได้มีการจำแนก Unary (LT) - และ (LT_1) -operations ตลอดจนบรรยาย equivalence relations ทั้งหมดซึ่งไม่แปรเปลี่ยนภายใต้ f ในโครงการวิจัยนี้ เราศึกษา ให้นิยามและจำแนก n -ary operations สำหรับทุก $n > 1$ ซึ่งเป็นทั้ง (LT) - และ (LT_1) -operations รวมถึงบรรยาย invariant equivalence relations ทั้งหมดซึ่งไม่แปรเปลี่ยนภายใต้ f ผลการศึกษาเหล่านี้สามารถประยุกต์กับสาขาวิชาอื่นๆ ที่ดำเนินการศึกษาด้วยวิธี iteration และ recursion นอกจากนี้ยังจำแนก unary operations บนเซตจำกัดทั้งหมดซึ่ง $\lambda(f) = k - t$ สำหรับทุก $1 \leq t \leq n - 1$ และให้รายละเอียดเกี่ยวกับ equivalence relations บนเซตจำกัดซึ่งไม่แปรเปลี่ยนภายใต้ unary operations เหล่านี้

เราใช้สัญลักษณ์ $f : A \rightarrow A$ แทน partial unary operation ซึ่งนิยามบน A ถ้า f เป็น partial unary operation เราแทนโดเมนของ f ด้วย $\text{dom } f$ และให้ $\text{Im } f = \{ f(a) \mid a \in \text{Dom } f \}$ แทนเซตส่วนฉายของ f และถ้า f เป็น proper partial operation บน A ที่ไม่สมนัยหนึ่งต่อหนึ่งบน $\text{dom } f$ แล้ว $|A| > |\text{dom } f| \geq |\text{Im } f|$ และมีจำนวนเต็มบวกตัวน้อยสุด $\lambda(f)$ ที่ทำให้ $\text{Im } f^{\lambda(f)} = \text{Im } f^{\lambda(f)+1}$ ในโครงการวิจัยนี้เราศึกษาเพื่อจำแนก partial unary operations ที่มี $\lambda(f) = k$ และ $\lambda(f) = k - 1$ และในทั้งสองกรณีเราบรรยาย equivalence relations บน A ทั้งหมดซึ่งไม่แปรเปลี่ยนภายใต้ f ผลงานเหล่านี้เป็นการวางนัยทั่วไปของกรณี total unary operations

เราอาจศึกษา algebraic structure ของ Clone ได้หลายแบบ และเพราะเราอาจนิยาม $(n+1)$ -ary operation บน $O^n(A)$ โดย $S^n(f, g_1, \dots, g_n)(a_1, \dots, a_n) := f(g_1(a_1, \dots, a_n), \dots, g_n(a_1, \dots, a_n))$ สำหรับทุก n -ary operations f, g_1, \dots, g_n บน A และ $a_1, \dots, a_n \in A$ จึงได้นิยาม binary operation $+$ บน $O^n(A)$ โดย $f + g := S^n(f, g_1, \dots, g_n)$ ที่ทำให้ $(O^n(A); +)$ เป็น semigroup สำหรับโครงการวิจัยนี้ แทนที่จะศึกษา clone บนเซตจำกัดใดๆ ซึ่งเป็นโครงสร้างที่ซับซ้อนและยุ่งยาก เราศึกษา semigroups ของ n -ary operations ซึ่งเป็น subsemigroups ของ $(O^n(A); +)$ และศึกษาสมบัติของ semigroup เหล่านี้แทน ตลอดจนศึกษา Green's relations และจำแนก constant subsemigroups, rectangular bands และ normal bands ใน $(O^n(A); +)$

ABSTRACT

Research Title	Classification of finite algebras by unary non-bijective functions
Researcher	Ass. Prof. Dr. Chawewan Ratanaprasert, Faculty of Science, Silpakorn University. E-mail address : ratach@su.ac.th
Research Grants	The Thailand Research Fund
Period	July 28, 2005 – July 28, 2008
Keywords	unary operation, pre-period, partial operation, n-ary operation, invariant equivalence relation, clone, semigroup, band

Iterating a unary operation f defined on a finite set A with $|A| = k$, one obtains the descending chain $A \supseteq \text{Im } f \supseteq \text{Im } f^2 \supseteq \dots \supseteq \text{Im } f^m = \text{Im } f^{m+1}$. The least integer $\lambda(f)$ with $\text{Im } f^{\lambda(f)} = \text{Im } f^{\lambda(f)+1}$ is called the pre-period of f . The pre-period of f is an integer between 0 and $k - 1$. If $\lambda(f) = k - 1$ and $k \geq 1$, then f is called a long-tailed (LT)-operation and if $\lambda(f) = k - 2$ for $k \geq 2$, f is said to be an (LT_1) -operation. Unary (LT) - and (LT_1) -operations and their invariant equivalence relations have been characterized. In the project, we consider the iteration of n -ary operations for $n > 1$, define and characterize (LT) - and (LT_1) -operations and their invariant equivalence relations. The results can be applied in all fields where iteration and recursion plays a role.

We denote a partial unary operation f defined on A by $f : A \rightarrow \circ A$ and denote the domain of f by $\text{dom } f$ and also let $\text{Im } f = \{ f(a) \mid a \in \text{Dom } f \}$ be the image of f . If $f : A \rightarrow \circ A$ is a proper partial operation on A which is not bijective on its domain, then $|A| > |\text{dom } f| \geq |\text{Im } f|$ and there is a least integer $\lambda(f)$ with $\text{Im } f^{\lambda(f)} = \text{Im } f^{\lambda(f)+1}$. In the project, we characterize partial unary operations with $\lambda(f) = k$ and $\lambda(f) = k - 1$. In both cases, we describe the equivalence relations on A which are invariant with respect to such partial unary operations. This generalizes similar results for total unary operations.

There are several ways to regard a clone as an algebraic structure. If f, g_1, \dots, g_n are n -ary operations defined on A , by $S^n(f, g_1, \dots, g_n)(a_1, \dots, a_n) := f(g_1(a_1, \dots, a_n), \dots, g_n(a_1, \dots, a_n))$ for all $a_1, \dots, a_n \in A$ an $(n+1)$ -ary operation on the set $O^n(A)$ of all n -ary operations can be defined; and one can derive a binary operation $+$ defined by $f+g := S^2(f, g_1, \dots, g_n)$ and obtains a semigroup $(O^n(A); +)$. The collection of all clones of operations on a finite set forms a complete lattice. This lattice is well-described if $|A| = 2$. If $|A| > 2$, this lattice is uncountably infinite and very complex. In the project, instead of clones we study semigroups of n -ary operations, i.e. subsemigroups of the semigroup $(O^n(A); +)$ and their properties. We consider Green's relations for the semigroup $(O^n(A); +)$, characterize all constant subsemigroups of $(O^n(A); +)$, all semilattices, rectangular bands and normal bands contained in $(O^n(A); +)$.