



# รายงานวิจัยฉบับสมบูรณ์

โครงการ

“ฟังก์ชันวางนัยทั่วไป และ ทฤษฎีบทจุดตรึง

ในปริภูมิบานาค”

โดย ศาสตราจารย์ อำนวย ขนนั่นไทย และ คณะ

กรกฎาคม 2552

## รายงานวิจัยฉบับสมบูรณ์

โครงการ “ฟังก์ชันวางนัยทั่วไป และ ทฤษฎีบทจุดตรึงในปริภูมิบานาค”

คณะผู้วิจัย

1. ศาสตราจารย์ อำนวย จนนไทย

2. ศาสตราจารย์ ดร. สุเทพ สวนใต้

ภาควิชาคณิตศาสตร์ คณะวิทยาศาสตร์

มหาวิทยาลัยเชียงใหม่

สนับสนุนโดยสำนักงานกองทุนสนับสนุนการวิจัย

(ความเห็นในรายงานนี้เป็นของผู้วิจัย

สกว. ไม่จำเป็นต้องเห็นด้วยเสมอไป )

## ABSTRACT

The first main purpose of this project is to study the operators concerning the heat equations and the wave equation such operator are the Laplace, the ultra hyperbolic the Diamond operators and the mixed operator. In doing research for such operators, we are succeeded in obtaining the interesting solution that all solutions cover all old area of solution before and all such solution are beautiful uniqueness.

The second purpose of this project is to construct new iterative methods for approximating a fixed point and common fixed points of nonlinear mappings. In this part, we introduce a new three-step iteration with errors for nonexpansive nonself-mappings in a uniformly convex Banach space. Weak and strong convergence theorems of the new three- step iteration under certain control conditions are established. We also modify Noor iterations for non-Lipshitzian mappings in Banach spaces and prove weak and strong convergence theorems of the modified Noor iterations under some control conditions. For finding a common fixed point of a finite family of nonexpansive mappings, we introduce a new iterative method for them and prove weak and strong convergence theorems under some suitable control conditions. Moreover, we introduce new methods for finding a common element of a fixed point set of nonlinear mappings and the set of solutions of equilibrium problems. Our results improve and extend many results in this area.

## EXECUTIVE SUMMARY

**Title :** Generalized Functions and Fixed Point Theory in Banach Spaces

ฟังก์ชันวงนัยทั่วไป และ ทฤษฎีบทจุดตรึงในปริภูมิบานาค

**Researchers :** 1. Prof. Amnuay Kananthai, Head of the Project

2. Prof. Dr. Suthep Suantai

Department of Mathematics, Faculty of Science, Chiang Mai University

**Budget:** 2,000,000 Bath

**Research Duration :** 31 July 2006 - 30 July 2009

### Principles Theory, Rationale and / or Hypotheses

Generalized functions and fixed point theory play an important role in mathematical analysis that have the applications widely in the other fields related to science and technology. Basically, Generalized functions cover all Classical functions (Ordinary functions). It is well known that the generalized functions can be applied to solve the problems of the wave equations, particularly the wave functions which are not continuous such as shock wave. That kind of the wave functions is so difficult to interpret in the term of ordinary function. At the beginning in the year 1950, the Russian mathematician, S. L. Sobolev studied partial differential equation which related to the generalized function and he was the first in setting the background of such generalized functions in the year 1960, L. Schwartz studied from S. L. Sobolev by extending and developing some new concepts and obtain many properties and theorems in such area.

In the area of generalized functions, he also discovered some operators that concern the partial differential equations, for examples, the elliptic operator, the hyperbolic operator and the parabolic operator. The famous elliptic operator is the Laplacian  $\Delta$  that defined by

$$\Delta = \frac{\partial^2}{\partial x_1^2} + \frac{\partial^2}{\partial x_2^2} + \dots + \frac{\partial^2}{\partial x_n^2}$$

And the hyperbolic operator is the wave operator  $\square = \frac{\partial^2}{\partial t^2} - \Delta$  and also the parabolic operator is the heat operator defined by  $L = \frac{\partial}{\partial t} - \Delta$

In the year 1987, S. E. Trione studied the ultra-hyperbolic operator which is an extension of the wave operator. The ultra-hyperbolic operator iterated  $k$ -times is defined by

$$\square^k = \left[ \sum_{i=1}^p \frac{\partial^2}{\partial x_i^2} - \sum_{j=p+1}^{p+q} \frac{\partial^2}{\partial x_j^2} \right]^k$$

$p + q = n$  where  $n$  is the dimension of the Euclidean space  $\mathbb{R}^n$  and  $k$  is a nonnegative integer.

In the year 1994, M. A. Tellez has shown that the operator  $\square^k$  exists only for  $n$  is an odd with  $p$  is odd and  $q$  is even.

In the year 1997, A. Kananthai established the new operator that is called the Diamond operator

$$\diamond \text{ iterated } k\text{-times defined by } \diamond^k = \left[ \left( \sum_{i=1}^p \frac{\partial^2}{\partial x_i^2} \right)^2 - \left( \sum_{j=p+1}^{p+q} \frac{\partial^2}{\partial x_j^2} \right)^2 \right]^k, \quad p + q = n.$$

The diamond operator covers all the Laplacian and ultra operators. He also obtained the elementary solution for such Diamond operator.

In the year 2001, A. Kananthai and S. Suantai extend the Diamond operator to be the operator

$$\oplus^k = \left[ \left( \sum_{i=1}^p \frac{\partial^2}{\partial x_i^2} \right)^4 - \left( \sum_{j=p+1}^{p+q} \frac{\partial^2}{\partial x_j^2} \right)^4 \right]^k \quad \text{and obtained the interesting elementary solution.}$$

All operators that have been mentioned are based on the area of generalized functions.

In our research, we will study the operator in the form of nonlinear equations which are the new frontier research.

In studying the problems in science and technology, usually those problems are formulated in term of equations or inequalities. So the question arise from this point that how we know the existence of a solution of such equation and inequalities, and once we know the existence of the solution, the second question will be asked, how can we find that solution. So there are two problems which are concerned in solving the solution of linear and nonlinear equation :

1. The existence of the solutions of such equations and
2. The method of solving the solutions of such equations.

So we are interested in studying those two problems in a general Banach space setting.

#### Research Objectives

1. Study various properties of the Laplacian, Ultra hyperbolic, Diamond and the compound operators.
2. Study the elementary solutions of those operators mentioned in 1.
3. Study the solutions of the partial differential operators related to non-linear wave and heat equations.
4. Construct and study new fixed point iteration methods for approximating fixed points of nonlinear mappings in a Banach space.
5. Study the existence and uniqueness of the fixed points of generalized contraction mappings

6. Study the geometric properties related to fixed point theory.
7. To build the young researchers in the area of generalized functions and fixed point theory.

#### **Usefulness of the research**

1. Obtain various properties of the Laplacian, Ultra hyperbolic, Diamond and the compound operators
2. Obtain the elementary solutions of those operators mentioned in 1.
3. Obtain the solutions of the partial differential operators related to non-linear wave and heat equations.
4. Obtain new fixed point iteration methods for approximating fixed points of nonlinear mappings in a Banach space.
5. Obtain the theorems of the existence and uniqueness of the fixed points of generalized contraction mappings
6. Obtain some geometric properties related to fixed point theory

## RESEARCH CONTENTS

### Chapter I

Introduction

### Chapter II

The Solution of the Partial Differential Operators  
Related to Generalized Functions

### Chapter III

Fixed Point Theory in Banach Spaces

# Chapter I

## Introduction

Generalized functions and fixed point theory play an important role in mathematical analysis that have the applications widely in the other fields related to science and technology. Basically, Generalized functions cover all Classical functions (Ordinary functions). It is well known that the generalized functions can be applied to solve the problems of the wave equations, particularly the wave functions which are not continuous such as shock wave. That kind of the wave functions is so difficult to interpret in the term of ordinary function. At the beginning in the year 1950, the Russian mathematician, S. L. Sobolev studied partial differential equation which related to the generalized function and he was the first in setting the background of such generalized functions in the year 1960, L. Schwartz studied from S. L. Sobolev by extending and developing some new concepts and obtain many properties and theorems in such area.

In the area of generalized functions, he also discovered some operators that concern the partial differential equations, for examples, the elliptic operator, the hyperbolic operator and the parabolic operator. The famous elliptic operator is the Laplacian  $\Delta$  that defined by

$$\Delta = \frac{\partial^2}{\partial x_1^2} + \frac{\partial^2}{\partial x_2^2} + \dots + \frac{\partial^2}{\partial x_n^2}$$

And the hyperbolic operator is the wave operator  $\square = \frac{\partial^2}{\partial t^2} - \Delta$  and also the parabolic operator is

the heat operator defined by  $L = \frac{\partial}{\partial t} - \Delta$

In the year 1987, S. E. Trione studied the ultra-hyperbolic operator which is an extension of the wave operator. The ultra-hyperbolic operator iterated k-times is defined by

$$\square^k = \left[ \sum_{i=1}^p \frac{\partial^2}{\partial x_i^2} - \sum_{j=p+1}^{p+q} \frac{\partial^2}{\partial x_j^2} \right]^k$$

$p + q = n$  where  $n$  is the dimension of the Euclidean space  $R^n$  and  $k$  is a nonnegative integer.

In the year 1994, M. A. Tellez has shown that the operator  $\square^k$  exists only for  $n$  is an odd with  $p$  is odd and  $q$  is even.

In the year 1997, A. Kananthai established the new operator that is called the Diamond operator

$$\diamond \text{ iterated } k\text{-times defined by } \diamond^k = \left[ \left( \sum_{i=1}^p \frac{\partial^2}{\partial x_i^2} \right)^2 - \left( \sum_{j=p+1}^{p+q} \frac{\partial^2}{\partial x_j^2} \right)^2 \right]^k, \quad p + q = n.$$

The diamond operator covers all the Laplacian and ultra operators. He also obtained the elementary solution for such Diamond operator.

In the year 2001, A. Kananthai and S. Suantai extend the Diamond operator to be the operator

$$\oplus^k = \left[ \left( \sum_{i=1}^p \frac{\partial^2}{\partial x_i^2} \right)^4 - \left( \sum_{j=p+1}^{p+q} \frac{\partial^2}{\partial x_j^2} \right)^4 \right]^k \quad \text{and obtained the interesting elementary solution.}$$

All operators that have been mentioned are based on the area of generalized functions.

In our research, we will study the operator in the form of nonlinear equations which are the new frontier research.

In studying the problems in science and technology, usually those problems are formulated in term of equations or inequalities. So the question arise from this point that how we know the existence of a solution of such equation and inequalities, and once we know the existence of the solution, the second question will be asked, how can we find that solution. So there are two problems which are concerned in solving the solution of linear and nonlinear equation :

1. The existence of the solutions of such equations and
2. The method of solving the solutions of such equations.

So we are interested in studying those two problems in a general Banach space setting.

## Chapter 2

### Some operators related to the heat and the wave equations

One part of doing research is the title "Some operator related to the heat and the wave equations". For the past 3 years, we have succeeded in doing research in such operators. We obtained many papers that can be classified in the following groups.

**The first group** are the reprints papers consisting of the following paper

1. A. Kananthai and K. Nonlaopon, On the generalized nonlinear ultra-hyperbolic heat equation related to the spectrum, Computational and Applied Mathematics, Volume 28 N. 2, pp. 1-10, 2009.
2. W. Satsanit and A. Kananthai, On the ultra-hyperbolic wave operator, International Journal of Pure and Applied Mathematics, Volume 52 N. 1, pp. 117-126, 2009.
3. C. Bunpog and A. Kananthai, On the Green Function of the Operator Related to the Bessel Helmholtz Operator and the Bessel Klein-Gordon Operator, Journal of Applied Functional Analysis, Volume 4 pp 10-19, 2009.

**The second group**, the accepted papers.

1. W. Satsanit and A. Kananthai, Diamond operator related to Bihmonic equation, Far East Journal of Applied Mathematics.
2. W. Satsanit and A. Kananthai, The operator and its spectrum related to heat equation, International Journal of Pure and Applied Mathematics.

**The third group**, submissions paper.

1. Amnuay Kananthai, On the Diamond-Wave Operator, submitted to Journal of Applied Mathematics and Computation.

2. Amnuay Kananthai, On the Nonlinear heat equation related to the operator, submitted to

*Nonlinear Analysis and Application.*

## On the generalized nonlinear ultra-hyperbolic heat equation related to the spectrum

A. KANANTHAI and K. NONLAOPON\*

Department of Mathematics, Chiang Mai University, Chiang Mai, 50200 Thailand

E-mail: malamnka@science.cmu.ac.th

**Abstract.** In this paper, we study the nonlinear equation of the form

$$\frac{\partial}{\partial t} u(x, t) - c^2 \square^k u(x, t) = f(x, t, u(x, t))$$

where  $\square^k$  is the ultra-hyperbolic operator iterated  $k$ -times, defined by

$$\square^k = \left( \frac{\partial^2}{\partial x_1^2} + \frac{\partial^2}{\partial x_2^2} + \cdots + \frac{\partial^2}{\partial x_p^2} - \frac{\partial^2}{\partial x_{p+1}^2} - \frac{\partial^2}{\partial x_{p+2}^2} - \cdots - \frac{\partial^2}{\partial x_{p+q}^2} \right)^k,$$

$p + q = n$  is the dimension of the Euclidean space  $\mathbb{R}^n$ ,  $(x, t) = (x_1, x_2, \dots, x_n, t) \in \mathbb{R}^n \times (0, \infty)$ ,  $k$  is a positive integer and  $c$  is a positive constant.

On the suitable conditions for  $f$ ,  $u$  and for the spectrum of the heat kernel, we can find the unique solution in the compact subset of  $\mathbb{R}^n \times (0, \infty)$ . Moreover, if we put  $k = 1$  and  $q = 0$  we obtain the solution of nonlinear equation related to the heat equation.

**Mathematical subject classification:** author, please, provide the AMS classif.

**Key words:** author, please, provide the keywords.

### 1 Introduction

It is well known that for the heat equation

$$\frac{\partial}{\partial t} u(x, t) = c^2 \Delta u(x, t) \quad (1.1)$$

#752/08. Received: 07/III/08. Accepted: 08/III/09.

\*Supported by The Royal Golden Jubilee Project grant no. PHD/0221/2543.

with the initial condition

$$u(x, 0) = f(x)$$

where  $\Delta = \sum_{i=1}^n \frac{\partial^2}{\partial x_i^2}$  is the Laplace operator and  $(x, t) = (x_1, x_2, \dots, x_n, t) \in \mathbb{R}^n \times (0, \infty)$ , and  $f$  is a continuous function, we obtain the solution

$$u(x, t) = \frac{1}{(4c^2\pi t)^{n/2}} \int_{\mathbb{R}^n} \exp\left[-\frac{|x-y|^2}{4c^2t}\right] f(y) dy \quad (1.2)$$

as the solution of (1.1).

Now, (1.2) can be written as  $u(x, t) = E(x, t) * f(x)$  where

$$E(x, t) = \frac{1}{(4c^2\pi t)^{n/2}} \exp\left[-\frac{|x|^2}{4c^2t}\right]. \quad (1.3)$$

$E(x, t)$  is called the *heat kernel*, where  $|x|^2 = x_1^2 + x_2^2 + \dots + x_n^2$  and  $t > 0$ , see [1, p. 208–209].

Moreover, we obtain  $E(x, t) \rightarrow \delta$  as  $t \rightarrow 0$ , where  $\delta$  is the Dirac-delta distribution. We also have extended (1.1) to be the equation

$$\frac{\partial}{\partial t} u(x, t) = c^2 \square u(x, t) \quad (1.4)$$

where  $\square$  is the *ultra-hyperbolic operator*, defined by

$$\square = \left( \frac{\partial^2}{\partial x_1^2} + \frac{\partial^2}{\partial x_2^2} + \dots + \frac{\partial^2}{\partial x_p^2} - \frac{\partial^2}{\partial x_{p+1}^2} - \frac{\partial^2}{\partial x_{p+2}^2} - \dots - \frac{\partial^2}{\partial x_{p+q}^2} \right).$$

We obtain the *ultra-hyperbolic heat kernel*

$$E(x, t) = \frac{(i)^q}{(4c^2\pi t)^{n/2}} \exp\left[\frac{\sum_{i=1}^p x_i^2 - \sum_{j=p+1}^{p+q} x_j^2}{4c^2t}\right]$$

where  $p + q = n$  is the dimension of the Euclidean space  $\mathbb{R}^n$  and  $i = \sqrt{-1}$ . For finding the kernel  $E(x, t)$  see [4].

In this paper, we extend (1.4) to be the general of the nonlinear form

$$\frac{\partial}{\partial t} u(x, t) - c^2 \square^k u(x, t) = f(x, t, u(x, t)) \quad (1.5)$$

for  $(x, t) \in \mathbb{R}^n \times (0, \infty)$  and with the following conditions on  $u$  and  $f$  as follows,

(1)  $u(x, t) \in C^{(2k)}(\mathbb{R}^n)$  for any  $t > 0$  where  $C^{(2k)}(\mathbb{R}^n)$  is the space of continuous function with  $2k$ -derivatives.

(2)  $f$  satisfies the Lipchitz condition, that is

$$|f(x, t, u) - f(x, t, w)| \leq A|u - w|$$

where  $A$  is constant and  $0 < A < 1$ .

(3)

$$\int_0^\infty \int_{\mathbb{R}^n} |f(x, t, u(x, t))| dx dt < \infty$$

for  $x = (x_1, x_2, \dots, x_n) \in \mathbb{R}^n$ ,  $t \in (0, \infty)$  and  $u(x, t)$  is continuous function on  $\mathbb{R}^n \times (0, \infty)$ .

Under such conditions of  $f, u$  and for the spectrum of  $E(x, t)$ , we obtain the convolution

$$u(x, t) = E(x, t) * f(x, t, u(x, t))$$

as a unique solution in the compact subset of  $\mathbb{R}^n \times (0, \infty)$  and  $E(x, t)$  is an elementary solution defined by (2.5).

## 2 Preliminaries

**Definition 2.1.** Let  $f(x) \in \mathbb{L}_1(\mathbb{R}^n)$ -the space of integrable function in  $\mathbb{R}^n$ . The Fourier transform of  $f(x)$  is defined by

$$\widehat{f}(\xi) = \frac{1}{(2\pi)^{n/2}} \int_{\mathbb{R}^n} e^{-i(\xi, x)} f(x) dx \quad (2.1)$$

where  $\xi = (\xi_1, \xi_2, \dots, \xi_n)$ ,  $x = (x_1, x_2, \dots, x_n) \in \mathbb{R}^n$ ,  $(\xi, x) = \xi_1 x_1 + \xi_2 x_2 + \dots + \xi_n x_n$  is the usual inner product in  $\mathbb{R}^n$  and  $dx = dx_1 dx_2 \dots dx_n$ .

Also, the inverse of Fourier transform is defined by

$$f(x) = \frac{1}{(2\pi)^{n/2}} \int_{\mathbb{R}^n} e^{i(\xi, x)} \widehat{f}(\xi) d\xi. \quad (2.2)$$

**Definition 2.2.** The spectrum of the kernel  $E(x, t)$  defined by (2.5) is the bounded support of the Fourier transform  $\widehat{E}(\xi, t)$  for any fixed  $t > 0$ .

**Definition 2.3.** Let  $\xi = (\xi_1, \xi_2, \dots, \xi_n)$  be a point in  $\mathbb{R}^n$  and we write

$$u = \xi_1^2 + \xi_2^2 + \dots + \xi_p^2 - \xi_{p+1}^2 - \xi_{p+2}^2 - \dots - \xi_{p+q}^2, \quad p + q = n.$$

Denote by

$$\Gamma_+ = \{\xi \in \mathbb{R}^n : \xi_1 > 0 \text{ and } u > 0\}$$

the set of an interior of the forward cone, and  $\bar{\Gamma}_+$  denotes the closure of  $\Gamma_+$ .

Let  $\Omega$  be spectrum of  $E(x, t)$  defined by Definition 2.2 for any fixed  $t > 0$  and  $\Omega \subset \bar{\Gamma}_+$ . Let  $\widehat{E}(\xi, t)$  be the Fourier transform of  $E(x, t)$  and define

$$\widehat{E}(\xi, t) = \begin{cases} \frac{1}{(2\pi)^{n/2}} \exp \left[ c^2 t \left( \sum_{j=p+1}^{p+q} \xi_j^2 - \sum_{i=1}^p \xi_i^2 \right)^k \right] & \text{for } \xi \in \Gamma_+, \\ 0 & \text{for } \xi \notin \Gamma_+. \end{cases} \quad (2.3)$$

**Lemma 2.1.** Let  $L$  be the operator defined by

$$L = \frac{\partial}{\partial t} - c^2 \square^k \quad (2.4)$$

where  $\square^k$  is the ultra-hyperbolic operator iterated  $k$ -times defined by

$$\square^k = \left( \frac{\partial^2}{\partial x_1^2} + \frac{\partial^2}{\partial x_2^2} + \dots + \frac{\partial^2}{\partial x_p^2} - \frac{\partial^2}{\partial x_{p+1}^2} - \frac{\partial^2}{\partial x_{p+2}^2} - \dots - \frac{\partial^2}{\partial x_{p+q}^2} \right)^k,$$

$p + q = n$  is the dimension of  $\mathbb{R}^n$ ,  $(x_1, x_2, \dots, x_n) \in \mathbb{R}^n$ ,  $t \in (0, \infty)$ ,  $k$  is a positive integer and  $c$  is a positive constant. Then we obtain

$$E(x, t) = \frac{1}{(2\pi)^n} \int_{\Omega} \exp \left[ c^2 t \left( \sum_{j=p+1}^{p+q} \xi_j^2 - \sum_{i=1}^p \xi_i^2 \right)^k + i(\xi, x) \right] d\xi \quad (2.5)$$

as a elementary solution of (2.4) in the spectrum  $\Omega \subset \mathbb{R}^n$  for  $t > 0$ .

**Proof.** Let  $LE(x, t) = \delta(x, t)$  where  $E(x, t)$  is the kernel or the elementary solution of operator  $L$  and  $\delta$  is the Dirac-delta distribution. Thus

$$\frac{\partial}{\partial t} E(x, t) - c^2 \square^k E(x, t) = \delta(x) \delta(t).$$

Take the Fourier transform defined by (2.1) to both sides of the equation, we obtain

$$\frac{\partial}{\partial t} \widehat{E(\xi, t)} - c^2 \left( \sum_{j=p+1}^{p+q} \xi_j^2 - \sum_{i=1}^p \xi_i^2 \right)^k \widehat{E(\xi, t)} = \frac{1}{(2\pi)^{n/2}} \delta(t).$$

Thus

$$\widehat{E(\xi, t)} = \frac{H(t)}{(2\pi)^{n/2}} \exp \left[ c^2 t \left( \sum_{j=p+1}^{p+q} \xi_j^2 - \sum_{i=1}^p \xi_i^2 \right)^k \right]$$

where  $H(t)$  is the Heaviside function. Since  $H(t) = 1$  for  $t > 0$ . Therefore,

$$\widehat{E(\xi, t)} = \frac{1}{(2\pi)^{n/2}} \exp \left[ c^2 t \left( \sum_{j=p+1}^{p+q} \xi_j^2 - \sum_{i=1}^p \xi_i^2 \right)^k \right]$$

which has been already defined by (2.3). Thus

$$E(x, t) = \frac{1}{(2\pi)^{n/2}} \int_{\mathbb{R}^n} e^{i(\xi, x)} \widehat{E(\xi, t)} d\xi = \frac{1}{(2\pi)^{n/2}} \int_{\Omega} e^{i(\xi, x)} \widehat{E(\xi, t)} d\xi$$

where  $\Omega$  is the spectrum of  $E(x, t)$ . Thus from (2.3)

$$E(x, t) = \frac{1}{(2\pi)^n} \int_{\Omega} \exp \left[ c^2 t \left( \sum_{j=p+1}^{p+q} \xi_j^2 - \sum_{i=1}^p \xi_i^2 \right)^k + i(\xi, x) \right] d\xi \quad \text{for } t > 0.$$

□

**Definition 2.4.** Let us extend  $E(x, t)$  to  $\mathbb{R}^n \times \mathbb{R}$  by setting

$$E(x, t) = \begin{cases} \frac{1}{(2\pi)^n} \int_{\Omega} \exp \left[ c^2 t \left( \sum_{j=p+1}^{p+q} \xi_j^2 - \sum_{i=1}^p \xi_i^2 \right)^k + i(\xi, x) \right] d\xi & \text{for } t > 0, \\ 0 & \text{for } t \leq 0, \end{cases}$$

### 3 Main Results

**Theorem 3.1.** The kernel  $E(x, t)$  defined by (2.5) have the following properties:

- (1)  $E(x, t) \in C^\infty$ -the space infinitely differentiable.

$$(2) \left( \frac{\partial}{\partial t} - c^2 \square^k \right) E(x, t) = 0 \text{ for } t > 0.$$

(3)

$$|E(x, t)| \leq \frac{2^{2-n}}{\pi^{n/2}} \frac{M(t)}{\Gamma\left(\frac{p}{2}\right) \Gamma\left(\frac{q}{2}\right)}, \text{ for } t > 0,$$

where  $M(t)$  is a function of  $t$  in the spectrum  $\Omega$  and  $\Gamma$  denote the Gamma function. Thus  $E(x, t)$  is bounded for any fixed  $t > 0$ .

$$(4) \lim_{t \rightarrow 0} E(x, t) = \delta.$$

**Proof.**

(1) From (2.5), since

$$\frac{\partial^n}{\partial x^n} E(x, t) = \frac{1}{(2\pi)^n} \int_{\Omega} \frac{\partial^n}{\partial x^n} \exp \left[ c^2 t \left( \sum_{j=p+1}^{p+q} \xi_j^2 - \sum_{i=1}^p \xi_i^2 \right)^k + i(\xi, x) \right] d\xi.$$

Thus  $E(x, t) \in C^\infty$  for  $x \in \mathbb{R}^n, t > 0$ .

(2) By computing directly, we obtain

$$\left( \frac{\partial}{\partial t} - c^2 \square^k \right) E(x, t) = 0.$$

(3) We have

$$E(x, t) = \frac{1}{(2\pi)^n} \int_{\Omega} \exp \left[ c^2 t \left( \sum_{j=p+1}^{p+q} \xi_j^2 - \sum_{i=1}^p \xi_i^2 \right)^k + i(\xi, x) \right] d\xi.$$

$$|E(x, t)| \leq \frac{1}{(2\pi)^n} \int_{\Omega} \exp \left[ c^2 t \left( \sum_{j=p+1}^{p+q} \xi_j^2 - \sum_{i=1}^p \xi_i^2 \right)^k \right] d\xi.$$

By changing to bipolar coordinates

$$\xi_1 = r\omega_1, \xi_2 = r\omega_2, \dots, \xi_p = r\omega_p \quad \text{and}$$

$$\xi_{p+1} = s\omega_{p+1}, \xi_{p+2} = s\omega_{p+2}, \dots, \xi_{p+q} = s\omega_{p+q}$$

where  $\sum_{i=1}^p \omega_i^2 = 1$  and  $\sum_{j=p+1}^{p+q} \omega_j^2 = 1$ . Thus

$$|E(x, t)| \leq \frac{1}{(2\pi)^n} \int_{\Omega} \exp \left[ c^2 t (s^2 - r^2)^k \right] r^{p-1} s^{q-1} dr ds d\Omega_p d\Omega_q$$

where  $d\xi = r^{p-1} s^{q-1} dr ds d\Omega_p d\Omega_q$ ,  $d\Omega_p$  and  $\Omega_q$  are the elements of surface area of the unit sphere in  $\mathbb{R}^p$  and  $\mathbb{R}^q$  respectively. Since  $\Omega \subset \mathbb{R}^n$  is the spectrum of  $E(x, t)$  and we suppose  $0 \leq r \leq R$  and  $0 \leq s \leq L$  where  $R$  and  $L$  are constants. Thus we obtain

$$\begin{aligned} |E(x, t)| &\leq \frac{\Omega_p \Omega_q}{(2\pi)^n} \int_0^R \int_0^L \exp \left[ c^2 t (s^2 - r^2)^k \right] r^{p-1} s^{q-1} ds dr \\ &= \frac{\Omega_p \Omega_q}{(2\pi)^n} M(t) \quad \text{for any fixed } t > 0 \text{ in the spectrum } \Omega \\ &= \frac{2^{2-n}}{\pi^{n/2}} \frac{M(t)}{\Gamma(\frac{p}{2})\Gamma(\frac{q}{2})} \end{aligned} \quad (3.1)$$

where

$$M(t) = \int_0^R \int_0^L \exp \left[ c^2 t (s^2 - r^2)^k \right] r^{p-1} s^{q-1} ds dr \quad (3.2)$$

is a function of

$$t > 0, \quad \Omega_p = \frac{2\pi^{p/2}}{\Gamma(\frac{p}{2})} \quad \text{and} \quad \Omega_q = \frac{2\pi^{q/2}}{\Gamma(\frac{q}{2})}.$$

Thus, for any fixed  $t > 0$ ,  $E(x, t)$  is bounded.

(4) By (2.5), we have

$$E(x, t) = \frac{1}{(2\pi)^n} \int_{\Omega} \exp \left[ c^2 t \left( \sum_{j=p+1}^{p+q} \xi_j^2 - \sum_{i=1}^p \xi_i^2 \right)^k + i(\xi, x) \right] d\xi.$$

Since  $E(x, t)$  exists, then

$$\begin{aligned} \lim_{t \rightarrow 0} E(x, t) &= \frac{1}{(2\pi)^n} \int_{\Omega} e^{i(\xi, x)} d\xi \\ &= \frac{1}{(2\pi)^n} \int_{\mathbb{R}^n} e^{i(\xi, x)} d\xi \\ &= \delta(x), \quad \text{for } x \in \mathbb{R}^n. \end{aligned}$$

See [3, p. 396, Eq. (10.2.19b)]. □

**Theorem 3.2.** *Given the nonlinear equation*

$$\frac{\partial}{\partial t} u(x, t) - c^2 \square^k u(x, t) = f(x, t, u(x, t)) \quad (3.3)$$

*for  $(x, t) \in \mathbb{R}^n \times (0, \infty)$ ,  $k$  is positive number and with the following conditions on  $u$  and  $f$  as follows,*

(1)  $u(x, t) \in C^{(2k)}(\mathbb{R}^n)$  for any  $t > 0$  where  $C^{(2k)}(\mathbb{R}^n)$  is the space of continuous function with  $2k$ -derivatives.

(2)  $f$  satisfies the Lipchitz condition, that is

$$|f(x, t, u) - f(x, t, w)| \leq A|u - w|$$

where  $A$  is constant and  $0 < A < 1$ .

(3)

$$\int_0^\infty \int_{\mathbb{R}^n} |f(x, t, u(x, t))| dx dt < \infty$$

for  $x = (x_1, x_2, \dots, x_n) \in \mathbb{R}^n$ ,  $t \in (0, \infty)$  and  $u(x, t)$  is continuous function on  $\mathbb{R}^n \times (0, \infty)$ .

Then, for the spectrum of  $E(x, t)$  we obtain the convolution

$$u(x, t) = E(x, t) * f(x, t, u(x, t)) \quad (3.4)$$

as a unique solution of (3.3) for  $x \in \Omega_0$  where  $\Omega_0$  is an compact subset of  $\mathbb{R}^n$ ,  $0 \leq t \leq T$  with  $T$  is constant and  $E(x, t)$  is an elementary solution defined by (2.5) and also  $u(x, t)$  is bounded.

In particular, if we put  $k = 1$  and  $q = 0$  in (3.3) then (3.3) reduces to the nonlinear heat equation.

**Proof.** Convoluting both sides of (3.3) with  $E(x, t)$  and then we obtain the solution

$$u(x, t) = E(x, t) * f(x, t, u(x, t))$$

or

$$u(x, t) = \int_{-\infty}^{\infty} \int_{\mathbb{R}^n} E(r, s) f(x - r, t - s, u(x - r, t - s)) dr ds$$

where  $E(r, s)$  is given by Definition 2.4.

We next show that  $u(x, t)$  is bounded on  $\mathbb{R}^n \times (0, \infty)$ . We have

$$\begin{aligned} |u(x, t)| &\leq \int_{-\infty}^{\infty} \int_{\mathbb{R}^n} |E(r, s)| |f(x - r, t - s, u(x - r, t - s))| dr ds \\ &\leq \frac{2^{2-n}}{\pi^{n/2}} \frac{N \cdot M(t)}{\Gamma(\frac{n}{2}) \Gamma(\frac{q}{2})} \end{aligned}$$

by the condition (3) and (3.1) where

$$N = \int_0^\infty \int_{\mathbb{R}^n} |f(x, t, u(x, t))| dx dt.$$

Thus  $u(x, t)$  is bounded on  $\mathbb{R}^n \times (0, \infty)$ .

To show that  $u(x, t)$  is unique, suppose there is another solution  $w(x, t)$  of equation (3.3). Let the operator

$$L = \frac{\partial}{\partial t} - c^2 \Delta^k$$

then (3.3) can be written in the form

$$L u(x, t) = f(x, t, u(x, t)).$$

Thus

$$L u(x, t) - L w(x, t) = f(x, t, u(x, t)) - f(x, t, w(x, t)).$$

By the condition (2) of the Theorem,

$$|L u(x, t) - L w(x, t)| \leq A |u(x, t) - w(x, t)|. \quad (3.5)$$

Let  $\Omega_0 \times (0, T]$  be compact subset of  $\mathbb{R}^n \times (0, \infty)$  and  $L: C^{(2k)}(\Omega_0) \rightarrow C^{(2k)}(\Omega_0)$  for  $0 \leq t \leq T$ .

Now  $(C^{(2k)}(\Omega_0), \|\cdot\|)$  is a Banach space where  $u(x, t) \in C^{(2k)}(\Omega_0)$  for  $0 \leq t \leq T$ ,  $\|\cdot\|$  given by

$$\|u(x, t)\| = \sup_{x \in \Omega_0} |u(x, t)|.$$

Then, from (3.5) with  $0 < A < 1$ , the operator  $L$  is a contraction mapping on  $C^{(2k)}(\Omega_0)$ . Since  $(C^{(2k)}(\Omega_0), \|\cdot\|)$  is a Banach space and  $L: C^{(2k)}(\Omega_0) \rightarrow$

$C^{(2k)}(\Omega_0)$  is a contraction mapping on  $C^{(2k)}(\Omega_0)$ , by Contraction Theorem, see [3, p. 300], we obtain the operator  $L$  has a fixed point and has uniqueness property. Thus  $u(x, t) = w(x, t)$ . It follows that the solution  $u(x, t)$  of (3.3) is unique for  $(x, t) \in \Omega_0 \times (0, T]$  where  $u(x, t)$  is defined by (3.4).

In particular, if we put  $k = 1$  and  $q = 0$  in (3.3) then (3.3) reduces to the nonlinear heat equation

$$\frac{\partial}{\partial t} u(x, t) - c^2 \Delta u(x, t) = f(x, t, u(x, t))$$

which has solution

$$u(x, t) = E(x, t) * f(x, t, u(x, t))$$

where  $E(x, t)$  is defined by (2.5) with  $k = 1$  and  $q = 0$ . That is complete of proof.  $\square$

**Acknowledgement.** The authors would like to thank The Thailand Research Fund for financial support.

#### REFERENCES

- [1] F. John, "Partial Differential Equations", 4<sup>th</sup> Edition, Springer-Verlag, New York, (1982).
- [2] R. Haberman, "Elementary Applied Partial Differential Equations", 2<sup>nd</sup> Edition, Prentice-Hall International, Inc. (1983).
- [3] E. Kreyszig, "Introductory Functional Analysis with Applications", John Wiley & Sons Inc., (1978).
- [4] K. Nonlaopon and A. Kananthai, *On the Ultrahyperbolic Heat Kernel*, International Journal of Applied Mathematics, **13** (2) (2003), 215–225.

Volume 4, Number 1

January 2009

ISSN:1559-1948 (PRINT), 1559-1956 (ONLINE)

EUDOXUS PRESS,LLC



# JOURNAL OF APPLIED FUNCTIONAL ANALYSIS

# On the Green Function of the $(\diamond_B + m^4)^k$ Operator Related to the Bessel-Helmholtz Operator and the Bessel Klein-Gordon Operator

Chalermpon Bunpog and Amnuay Kananthai

Department of Mathematics, Chiang Mai University, Chiang Mai, 50200 Thailand  
malamka@science.cmu.ac.th

## Abstract

In this paper, we study the Green function of the operator  $(\diamond_B + m^4)^k$  which is iterated  $k$ -times and is defined by

$$(\diamond_B + m^4)^k = \left[ \left( \sum_{i=1}^p B_{x_i} \right)^2 - \left( \sum_{j=p+1}^{p+q} B_{x_j} \right)^2 + m^4 \right]^k, \quad (0.1)$$

where  $m$  is a positive real number and  $p+q=n$  is the dimension of  $\mathbb{R}_n^+$  and  $k$  is a nonnegative integer and  $B_{x_i} = \frac{\partial^2}{\partial x_i^2} + \frac{2v_i}{x_i} \frac{\partial}{\partial x_i}$ ,  $2v_i = 2\alpha_i + 1$ ,  $\alpha_i > -\frac{1}{2}$ ,  $x_i > 0$ . At first we study the Green function of the operator  $(\diamond_B + m^4)^k$ , we have that such a Green function related to the elementary solutions of the Bessel-Helmholtz operator  $(\Delta_B + m^2)^k$  iterated  $k$ -times and the Bessel Klein-Gordon operator  $(\square_B + m^2)^k$  iterated  $k$ -times. We also apply such a Green function to solve the solution of the equation  $(\diamond_B + m^4)^k u(x) = f(x)$  where  $f$  is a generalized function and  $u(x)$  is an unknown function for  $x \in \mathbb{R}_n^+$ .

**Keywords:** Green function, Bessel diamond operator, Helmholtz operator, Klein-Gordon operator

## 1 Introduction

A. Kananthai [1] first introduced the diamond operator  $\diamond^k$  iterated  $k$ -times, defined by

$$\diamond^k = \left[ \left( \sum_{i=1}^p \frac{\partial^2}{\partial x_i^2} \right)^2 - \left( \sum_{j=p+1}^{p+q} \frac{\partial^2}{\partial x_j^2} \right)^2 \right]^k,$$

the equation  $\diamond^k u(x) = f(x)$ , see [2], has been already studied and the convolution  $u(x) = (-1)^k R_{2k}^H(x) * R_{2k}^e * f(x)$  has been obtained as a solution of such an equation.

Later the equation  $(\diamond + m^4)^k u(x) = f(x)$ , see [3], has been studied and the convolution  $u(x) = (W_{2k}^H(u, m) * W_{2k}^e(v, m)) * (s^{*k})^{*-1}(x) * f(x)$  has been obtained a solution of such an equation.

Furthermore, Hüseyin Yildirim, Mzeki Sarikaya and Sermin Öztürk [4] first introduced the Bessel diamond operator  $\diamond_B^k$  iterated  $k$ -times, defined by

$$\diamond_B^k = \left[ \left( \sum_{i=1}^p B_{x_i} \right)^2 - \left( \sum_{j=p+1}^{p+q} B_{x_j} \right)^2 \right]^k \quad (1.1)$$

where  $B_{x_i} = \frac{\partial^2}{\partial x_i^2} + \frac{2v_i}{x_i} \frac{\partial}{\partial x_i}$ ,  $2v_i = 2\alpha_i + 1$ ,  $\alpha_i > -\frac{1}{2}$ ,  $x_i > 0$ . The operator  $\diamond_B^k$  can be expressed by  $\diamond_B^k = \Delta_B^k \square_B^k = \square_B^k \Delta_B^k$ , where

$$\Delta_B^k = \left( \sum_{i=1}^p B_{x_i} \right)^k \quad (1.2)$$

and

$$\square_B^k = \left[ \sum_{i=1}^p B_{x_i} - \sum_{j=p+1}^{p+q} B_{x_j} \right]^k \quad (1.3)$$

The equation  $\diamond_B^k u(x) = \delta(x)$ , see ([4], p.382), has been already studied and the convolution  $u(x) = (-1)^k S_{2k} * R_{2k}$  has been obtained as a solution of such an equation where the function  $S_{2k}$  and  $R_{2k}$  are defined by (2.1) and (2.2), respectively, with  $\alpha = \beta = 2k$ . In this work, we study the equation of the form

$$(\diamond_B + m^4)^k G(x) = \delta(x).$$

We obtain the elementary solution  $G(x) = (T_{2k}(x) * W_{2k}(x)) * (C^{*k})^{*-1}(x)$ , where the symbol  $*k$  denotes the convolution of itself  $k$ -times and the symbol  $*-1$  is an inverse of the convolution algebra,  $T_{2k}(x)$  is the elementary solution of the Bessel-Helmholtz operator  $(\Delta_B + m^2)^k$  iterated  $k$ -times, that is  $T_{2k}(x)$  satisfy the equation

$$(\Delta_B + m^2)^k u(x) = \delta(x)$$

and  $W_{2k}(x)$  is the elementary solution of the Bessel Klein-Gordon operator  $(\square_B + m^2)^k$  iterated  $k$ -times, that is  $W_{2k}(x)$  satisfy the equation

$$(\square_B + m^2)^k u(x) = \delta(x)$$

and  $C(x)$  is defined by

$$C(x) = \delta(x) - m^2(T_2(x) + W_2(V)) + 2m^4(T_2(x) * W_2(V)).$$

Moreover, we apply such a Green function to obtain the solution of the equation

$$(\diamond_B + m^4)^k u(x) = f(x).$$

where  $f$  is a generalized function.

## 2 Preliminaries

**Definition 2.1** Let  $x = (x_1, x_2, \dots, x_n), \nu = (\nu_1, \nu_2, \dots, \nu_n) \in \mathbb{R}_n^+$ . For any complex number  $\alpha$ , we define the function  $S_\alpha(x)$  by

$$S_\alpha(x) = \frac{2^{n+2|\nu|-2\alpha} \Gamma\left(\frac{n+2|\nu|-\alpha}{2}\right) |x|^{\alpha-n-2|\nu|}}{\prod_{i=1}^n 2^{\nu_i-\frac{1}{2}} \Gamma(\nu_i + \frac{1}{2})} \quad (2.1)$$

**Definition 2.2** Let  $x = (x_1, x_2, \dots, x_n), \nu = (\nu_1, \nu_2, \dots, \nu_n) \in \mathbb{R}_n^+$ , and denote by  $V = x_1^2 + x_2^2 + \dots + x_p^2 - x_{p+1}^2 - x_{p+2}^2 - \dots - x_{p+q}^2$  the nondegenerated quadratic form. Denote the interior of the forward cone by  $\Gamma_+ = \{x \in \mathbb{R}_n^+ : x_1 > 0, x_2 > 0, \dots, x_n > 0, V > 0\}$ . The function  $R_\beta(x)$  is defined by

$$R_\beta(x) = \frac{V^{\frac{\beta-n-2|\nu|}{2}}}{K_n(\beta)}, \quad (2.2)$$

where

$$K_n(\beta) = \frac{\pi^{\frac{n+2|\nu|-1}{2}} \Gamma\left(\frac{2+\beta-n-2|\nu|}{2}\right) \Gamma\left(\frac{1-\beta}{2}\right) \Gamma(\beta)}{\Gamma\left(\frac{2+\beta-p-2|\nu|}{2}\right) \Gamma\left(\frac{p-\beta}{2}\right)},$$

where  $\beta$  is a complex number.

**Definition 2.3** Let  $x = (x_1, x_2, \dots, x_n) \in \mathbb{R}_n^+$ , For any complex number  $\alpha$ , we define the function

$$T_\alpha(x) = \sum_{r=0}^{\infty} \frac{(-1)^r \Gamma\left(\frac{\eta}{2} + r\right)}{r! \Gamma\left(\frac{\eta}{2}\right)} (m^2)^r (-1)^{\frac{\alpha}{2}+r} S_{\alpha+2r}(x), \quad (2.3)$$

where  $\eta$  is a complex number and  $S_{\alpha+2r}(x)$  is defined in definition 2.1.

**Definition 2.4** Let  $x = (x_1, x_2, \dots, x_n)$ , For any complex number  $\beta$ , we define the function

$$W_\beta(x) = \sum_{r=0}^{\infty} \frac{(-1)^r \Gamma\left(\frac{\eta}{2} + r\right)}{r! \Gamma\left(\frac{\eta}{2}\right)} (m^2)^r R_{\beta+2r}(x), \quad (2.4)$$

where  $\eta$  is a complex number and  $R_{\beta+2r}(x)$  is defined in definition 2.2.

**Lemma 2.1** Given the equation  $\Delta_B^k u(x) = \delta(x)$  for  $x \in \mathbb{R}_n^+$ , where  $\Delta_B^k$  is defined by (1.2). Then

$$u(x) = (-1)^k S_{2k}(x)$$

where  $S_{2k}(x)$  is defined by (2.1), with  $\alpha = 2k$ .

**Proof.** See ([4], p.379). □

**Lemma 2.2** Given the equation  $\square_B^k u(x) = \delta(x)$  for  $x \in \mathbb{R}_n^+$ , where  $\square_B^k$  is defined by (1.3). Then

$$u(x) = R_{2k}(x)$$

where  $R_{2k}(x)$  is defined by (2.2), with  $\beta = 2k$

**Proof.** See ([4], p.379). □

**Lemma 2.3** (The elementary solution of the Bessel-Helmholtz operator).

Given the equation  $(\Delta_B + m^2)^k u(x) = \delta(x)$  for  $x \in \mathbb{R}_n^+$ , where  $\Delta_B$  is defined by (1.2) with  $k = 1$ . Then

$$u(x) = T_{2k}(x)$$

where  $T_{2k}(x)$  is defined by (2.3), with  $\alpha = 2k$ .

**Proof.** At first, the following formula is valid ([5], p.3),

$$\Gamma\left(\frac{\eta}{2} + r\right) = \frac{\eta}{2} \left(\frac{\eta}{2} + 1\right) \cdots \left(\frac{\eta}{2} + r - 1\right) \Gamma\left(\frac{\eta}{2}\right).$$

Equivalently,

$$\begin{aligned} (-1)^r \frac{1}{r!} \Gamma\left(\frac{\eta}{2} + r\right) &= \frac{(-1)^r \frac{\eta}{2} \left(\frac{\eta}{2} + 1\right) \cdots \left(\frac{\eta}{2} + r - 1\right) \Gamma\left(\frac{\eta}{2}\right)}{r!} \\ &= \frac{\left(-\frac{\eta}{2}\right) \left(-\frac{\eta}{2} - 1\right) \cdots \left[-\left(\frac{\eta}{2} + r - 1\right)\right]}{r!} \Gamma\left(\frac{\eta}{2}\right). \end{aligned}$$

We have,

$$(-1)^r \frac{1}{r!} \Gamma\left(\frac{\eta}{2} + r\right) = \binom{-\frac{\eta}{2}}{r} \Gamma\left(\frac{\eta}{2}\right).$$

Then, we obtain the function  $T_\alpha(x)$  is defined by Definition 2.3 become

$$T_\alpha(x) = \sum_{r=0}^{\infty} \binom{-\frac{\eta}{2}}{r} (m^2)^r (-1)^{\frac{\alpha}{2}+r} S_{\alpha+2r}(x). \quad (2.5)$$

Putting  $\alpha = \eta = 2k$  in (2.5), we have

$$T_{2k}(x) = \sum_{r=0}^{\infty} \binom{-k}{r} (m^2)^r (-1)^{k+r} S_{2k+2r}(x).$$

Since the operator  $\Delta_B$  is linearly continuous and has 1-1 mapping, then it has inverse, by Lemma 2.1 we obtain

$$\begin{aligned} T_{2k}(x) &= \sum_{r=0}^{\infty} \binom{-k}{r} (m^2)^r \delta(x) * \Delta_B^{-k-r} \\ &= (\Delta_B + m^2)^{-k} \delta(x), \end{aligned} \quad (2.6)$$

where  $(\Delta_B + m^2)^{-k}$  is the inverse operator of the operator  $(\Delta_B + m^2)^k$ . By applying the operator  $(\Delta_B + m^2)^k$  to both sides of (2.6), we obtain

$$(\Delta_B + m^2)^k T_{2k}(x) = (\Delta_B + m^2)^k (\Delta_B + m^2)^{-k} \delta(x).$$

Thus

$$(\Delta_B + m^2)^k T_{2k}(x) = \delta(x).$$

□

**Lemma 2.4** (The elementary solution of the Bessel Klein-Gordon operator).

Given the equation  $(\square_B + m^2)^k u(x) = \delta(x)$  for  $x \in \mathbb{R}_n^+$ , where  $\square_B$  is defined by (1.3) with  $k = 1$ . Then

$$u(x) = W_{2k}(x)$$

where  $W_{2k}(x)$  is defined by (2.4), with  $\alpha = 2k$ .

**Proof.** The proof of lemma 2.4 is similar to the proof of Lemma 2.3. □

**Lemma 2.5** Let  $T_{2k}(x)$  and  $W_{2k}(x)$  be defined by (2.3) and (2.4) respectively, where  $\alpha = \beta = 2k$ . Then the convolution  $T_{2k}(x) * W_{2k}(x)$  exist and it is lie in  $S'$ , where  $S'$  is a space of tempered distribution.

**Proof.** From (2.3) and (2.4) with  $\alpha = \beta = 2k$ , we have

$$\begin{aligned} T_{2k}(x) * W_{2k}(x) &= \left( \sum_{r=0}^{\infty} \frac{(-1)^r \Gamma(k+r)}{r! \Gamma(k)} (m^2)^r (-1)^{k+r} S_{2k+2r}(x) \right) \\ &\quad * \left( \sum_{r=0}^{\infty} \frac{(-1)^r \Gamma(k+r)}{r! \Gamma(k)} (m^2)^r R_{2k+2r}(x) \right) \\ &= \sum_{r=0}^{\infty} \sum_{s=0}^{\infty} \frac{(-1)^s \Gamma(k+s)}{s! \Gamma(k)} (m^2)^s \cdot \frac{(-1)^r \Gamma(k+r)}{r! \Gamma(k)} (m^2)^r \\ &\quad (-1)^{k+r} S_{2k+2r}(x) * R_{2k+2r}(x). \end{aligned}$$

Hüseyin Yıldırım, Mzeki Sarikaya and Sermin Öztürk ([4], p.380) has shown that  $S_{2k+2r}(x) * R_{2k+2r}(x)$  exists and is a tempered distribution. It follows that  $T_{2k}(x) * W_{2k}(x)$  exists and also is a tempered distribution. □

**Lemma 2.6** *Let  $T_2(x)$  and  $W_2(x)$  be defined by (2.3) and (2.4) respectively, where  $\alpha = \beta = 2$ . Then*

$$[(\Delta_B + m^2)(\square_B + m^2) - m^2(\Delta_B + \square_B)] (T_2(x) * W_2(x)) = C(x), \quad (2.7)$$

where  $C(x) = \delta(x) - m^2(T_2(x) + W_2(x)) + 2m^4(T_2(x) * W_2(x))$

**Proof.** We have

$$\begin{aligned} & [(\Delta_B + m^2)(\square_B + m^2) - m^2(\Delta_B + \square_B)] (T_2(x) * W_2(x)) = \\ & [(\Delta_B + m^2)(\square_B + m^2) (T_2(x) * W_2(x)) - m^2(\Delta_B + \square_B) (T_2(x) * W_2(x))] = \\ & [(\Delta_B + m^2)T_2(x) * (\square_B + m^2)W_2(x) - m^2(\Delta_B T_2(x) * W_2(x) + T_2(x) * \square_B W_2(x))] . \end{aligned} \quad (2.8)$$

From Lemma 2.3 and Lemma 2.4, for  $k = 1$  we have

$$(\Delta_B + m^2)T_2(x) = \delta(x) \quad \text{and} \quad (\square_B + m^2)W_2(x) = \delta(x),$$

respectively. Moreover,

$$\Delta_B T_2(x) = \delta(x) - m^2 T_2(x)$$

and

$$\square_B W_2(x) = \delta(x) - m^2 W_2(x),$$

thus(2.8) become

$$\begin{aligned} & [(\Delta_B + m^2)(\square_B + m^2) - m^2(\Delta_B + \square_B)] (T_2(x) * W_2(x)) = \\ & \delta(x) * \delta(x) - m^2 [(\delta(x) - m^2 T_2(x)) * W_2(x) + T_2(x) * (\delta(x) - m^2 W_2(x))] = \\ & \delta(x) - m^2 [W_2(x) - m^2 T_2(x) * W_2(x) + T_2(x) - m^2 T_2(x) * W_2(x)] = \\ & \delta(x) - m^2 (T_2(x) + W_2(x)) - 2m^4 (T_2(x) * W_2(x)) = C(x). \end{aligned}$$

□

**Lemma 2.7** *Let  $S_\alpha(x)$  be the function, defined by (2.1). Then*

$$S_\alpha(x) * S_\beta(x) = S_{\alpha+\beta}(x),$$

where  $\alpha$  and  $\beta$  are a positive **even** numbers.

**Proof.** See([4],p.380)

□

**Lemma 2.8** *Let  $R_\beta(x)$  be the function, defined by (2.2). Then*

$$R_\beta(x) * R_\alpha(x) = R_{\beta+\alpha}(x),$$

where  $\alpha$  and  $\beta$  are a positive **even** numbers.

**Proof.** Since  $R_\beta(x)$  and  $R_\alpha(x)$  are tempered distributions (see [4], p.380). Let  $\text{Supp} R_\beta(x) = K \subset \bar{\Gamma}_+$ , where  $K$  is a compact set and  $\bar{\Gamma}_+$  is a closure of  $\Gamma_+$  appears in Definition 2.2, then  $R_\beta(x) * R_\alpha(x)$  exists and is well defined. To show that  $R_\beta(x) * R_\alpha(x) = R_{\beta+\alpha}(x)$ , by Lemma 2.2  $\square_B^k u(x) = \delta(x)$  Then  $u(x) = R_{2k}(x)$ . Now,  $\square_B^k u(x) = \square_B^r \square_B^{k-r} u(x) = \delta(x)$  for  $r < k$ , then by Lemma 2.2  $\square_B^{k-r} u(x) = R_{2r}(x)$ . Convolving both sides by  $R_{2(k-r)}(x)$  we obtain

$$R_{2(k-r)}(x) * \square_B^{k-r} u(x) = R_{2(k-r)}(x) * R_{2r}(x)$$

or,

$$\square_B^{k-r} R_{2(k-r)}(x) * u(x) = R_{2(k-r)}(x) * R_{2r}(x)$$

by Lemma 2.2 again, we have

$$\delta(x) * u(x) = R_{2(k-r)}(x) * R_{2r}(x).$$

It follow that

$$u(x) = R_{2(k-r)}(x) * R_{2r}(x).$$

Since  $u(x) = R_{2k}(x)$ , thus

$$R_{2(k-r)}(x) * R_{2r}(x) = R_{2k}(x).$$

Let  $\beta = 2(k-r)$  and  $\alpha = 2r$ , actually  $\beta$  and  $\alpha$  are positive even numbers. It follows that  $R_\beta(x) * R_\alpha(x) = R_{\beta+\alpha}(x)$  as required.  $\square$

### 3 Main Results

**Theorem 3.1** *Given the equation*

$$(\diamond_B + m^4)^k G(x) = \delta(x) \quad (3.1)$$

where  $(\diamond_B + m^4)^k$  is the operator iterated  $k$ -times defined by (0.1),  $\delta$  is the Dirac-delta distribution,  $x = (x_1, x_2, \dots, x_n) \in \mathbb{R}_n^+$  and  $k$  is a nonnegative integer. Then we obtain  $G(x) = T_{2k}(x) * W_{2k}(x) * (C^{*k}(x))^{*-1}$  is a Green function for the operator  $(\diamond_B + m^4)^k$  iterated  $k$ -time where  $\diamond_B$  is defined by (1.1) with  $k = 1$ ,  $m$  is a nonnegative real number and

$$C(x) = \delta(x) - m^2(T_2(x) + W_2(x)) + 2m^4(T_2(x) * W_2(x)) \quad (3.2)$$

$C^{*k}(x)$  denote the convolution of  $C$  it self  $k$ -time,  $(C^{*k}(x))^{*-1}$  denote the inverse of  $C^{*k}(x)$  in the convolution algebra. Moreover  $C(x)$  is a tempered distribution.

**Proof.** Since  $(\diamond_B + m^4)^k = ((\Delta_B + m^2)(\square_B + m^2) - m^2(\Delta_B + \square_B))^k$ .

$$\begin{aligned} & [(\Delta_B + m^2)(\square_B + m^2) - m^2(\Delta_B + \square_B)] \cdot \\ & [(\Delta_B + m^2)(\square_B + m^2) - m^2(\Delta_B + \square_B)]^{k-1} G(x) = \delta(x) \end{aligned} \quad (3.3)$$

From Lemma 2.5 we have  $T_2(x) * W_2(x)$  exists and is a tempered distribution. Convolving both sides of the above equation by  $T_2(x) * W_2(x)$ , we obtain

$$\begin{aligned} & [(\Delta_B + m^2)(\square_B + m^2) - m^2(\Delta_B + \square_B)] (T_2(x) * W_2(x)) * \\ & [(\Delta_B + m^2)(\square_B + m^2) - m^2(\Delta_B + \square_B)]^{k-1} G(x) = (T_2(x) * W_2(x)) * \delta(x) \end{aligned}$$

by Lemma 2.6, we have

$$C(x) * [(\Delta_B + m^2)(\square_B + m^2) - m^2(\Delta_B + \square_B)]^{k-1} G(x) = (T_2(x) * W_2(x)) * \delta(x).$$

Keeping on convolving both sides of the above equation by  $T_2(x) * W_2(x)$  up to  $k-1$  times, we have

$$C^{*k}(x) * G(x) = (T_2(x) * W_2(x))^{*k},$$

where  $*k$  denotes the convolution of itself  $k$ -times.

By Lemma 2.7, Lemma 2.8 and definitions of  $T_\alpha(x)$  and  $W_\beta(x)$ , we have

$$(T_2(x) * W_2(x))^{*k} = T_{2k}(x) * W_{2k}(x),$$

then

$$C^{*k}(x) * G(x) = T_{2k}(x) * W_{2k}(x).$$

Now, consider the function  $C^{*k}(x)$ , since  $\delta(x)$ ,  $T_2(x)$ ,  $W_2(x)$  and  $T_2(x) * W_2(x)$  are lies in  $\mathcal{S}'$  where  $\mathcal{S}'$  is a space of tempered distribution, then  $C(x) \in \mathcal{S}'$ , moreover by ([6], p.152) we obtain  $C^{*k}(x) \in \mathcal{S}'$ . Since  $T_{2k}(x) * W_{2k}(x) \in \mathcal{S}'$ , choose  $\mathcal{S}' \subset \mathcal{D}'_{\mathcal{R}}$  where  $\mathcal{D}'_{\mathcal{R}}$  is the right-side distribution which is a subspace of  $\mathcal{D}'$  of distribution. Thus  $T_{2k}(x) * W_{2k}(x) \in \mathcal{D}'_{\mathcal{R}}$ , it follow that  $T_{2k}(x) * W_{2k}(x)$  is an element of convolution algebra, thus by ([7], p.150-151), we have that the equation (2.8) has a unique solution

$$G(x) = T_{2k}(x) * W_{2k}(x) * (C^{*k}(x))^{*-1}$$

where  $(C^{*k}(x))^{*-1}$  is an inverse of  $C^{*k}$  in the convolution algebra,  $G(x)$  is called the Green function of the operator  $(\diamond_B + m^4)^k$ . Since  $T_{2k}(x) * W_{2k}(x)$  and  $(C^{*k}(x))^{*-1}$  are lies in  $\mathcal{S}'$ , then by ([6], p.152) again, we have  $T_{2k}(x) * W_{2k}(x) * (C^{*k}(x))^{*-1} \in \mathcal{S}'$ . Hence  $G(x)$  is a tempered distribution.  $\square$

**Theorem 3.2** *Given the equation*

$$(\diamond_B + m^4)^k u(x) = f(x) \quad (3.4)$$

*where  $f$  is a given generalized function and  $u(x)$  is an unknown function, we obtain*

$$u(x) = G(x) * f(x)$$

*is a unique solution of the equation (3.4) where  $G(x)$  is a Green function for  $(\diamond_B + m^4)^k$ .*

**Proof.** Convolving both sides of (3.4) by  $G(x)$  where  $G(x)$  is a Green function for  $(\diamond_B + m^4)^k$  in theorem 3.1, we obtain

$$G(x) * (\diamond_B + m^4)^k u(x) = G(x) * f(x)$$

or,

$$(\diamond_B + m^4)^k G(x) * u(x) = G(x) * f(x)$$

applying the Theorem 3.1, we have

$$\delta(x) * u(x) = G(x) * f(x).$$

Therefor,

$$u(x) = G(x) * f(x).$$

Since  $G(x)$  is unique. Hence  $u(x)$  is a unique solution of the equation (3.4).  $\square$

#### Acknowledgement.

The authors would like to thank The Commission on Higher Education Scholarship and Graduate School, Chiang Mai University, Thailand for financial support.

## References

- [1] A. Kananthai, *On the solution of the n-dimensional Diamond operator*, Applied Mathematics and computation, vol. 88, Elsevier Science Inc., New York, 1997, p. 27-37.
- [2] A. Kananthai, *On the Diamond operator related to the wave equation*, Nonlinear Analysis, 2001, 47 (2), p. 1373-1382.
- [3] A. Kananthai, *On the Green function of the Diamond operator related to the Klein-Gordon operator*, Bull. Cal. Math. Soc., 2001, 93 (5), p.353-360.

- [4] Hüseyin Yildirim, Mzeki Sarikaya and Sermin Öztürk, *The solution of the  $n$ -dimensional Bessel diamond operator and the Fourier-Bessel transform of their convolution*, Proc. Indian Acad. Sci. (Math. Sci.) Vol. 114, No.4, November 2004, 375–387.
- [5] Bateman, Manuscript Project, *Higher Trascendental Functions*, Vol.I, Mc-Graw Hill, New York, 1953.
- [6] Donoghue, W. F., *Distributions and Fourier transform*, Academic Press, (1969).
- [7] Zemanian, A. H., *Distribution Theory and Transform Analysis*, Mc-Graw Hill, New York, (1964).

# ON THE ULTRA-HYPERBOLIC WAVE OPERATOR

Wanchak Satsanit<sup>1</sup>, Amnuay Kananthai<sup>2</sup> §

<sup>1,2</sup>Department of Mathematics

Faculty of Science

Chiang Mai University

Chiang Mai, 50200, THAILAND

<sup>2</sup>e-mail: malamnka@science.cmu.ac.th

**Abstract:** In this paper, we study the generalized wave equation of the form

$$\frac{\partial^2}{\partial t^2} u(x, t) + c^2 (\square)^k u(x, t) = 0$$

with the initial conditions

$$u(x, 0) = f(x), \quad \frac{\partial}{\partial t} u(x, 0) = g(x),$$

where  $u(x, t) \in \mathbb{R}^n \times [0, \infty)$ ,  $\mathbb{R}^n$  is the  $n$ -dimensional Euclidean space,  $\square^k$  is the ultra-hyperbolic operator iterated  $k$ -times defined by

$$\square^k = \left( \frac{\partial^2}{\partial x_1^2} + \frac{\partial^2}{\partial x_2^2} + \cdots + \frac{\partial^2}{\partial x_p^2} - \frac{\partial^2}{\partial x_{p+1}^2} - \frac{\partial^2}{\partial x_{p+2}^2} - \cdots - \frac{\partial^2}{\partial x_{p+q}^2} \right)^k,$$

$p+q=n$ ,  $c$  is a positive constant,  $k$  is a nonnegative integer,  $f$  and  $g$  are continuous and absolutely integrable functions. We obtain  $u(x, t)$  as a solution for such equation. Moreover, by  $\epsilon$ -approximation we also obtain the asymptotic solution  $u(x, t) = O(\epsilon^{-n/k})$ . In particular, if we put  $n = 1$ ,  $k = 2$  and  $q = 0$ , the  $u(x, t)$  reduces to the solution of the beam equation

$$\frac{\partial^2}{\partial t^2} u(x, t) + c^2 \frac{\partial^4}{\partial x^4} u(x, t) = 0.$$

**AMS Subject Classification:** 35L05

**Key Words:** generalized wave equation, beam equation, tempered distribution

Received: March 12, 2009

© 2009 Academic Publications

Correspondence author

## 1. Introduction

It is well known that for the 1-dimensional wave equation

$$\frac{\partial^2}{\partial t^2}u(x, t) = c^2 \frac{\partial^2}{\partial x^2}u(x, t), \quad (1)$$

we obtain  $u(x, t) = f(x + ct) + g(x - ct)$  as a solution of the equation, where  $f$  and  $g$  are continuous. Also for the  $n$ -dimensional wave equation

$$\frac{\partial^2}{\partial t^2}u(x, t) + c^2 \Delta u(x, t) = 0, \quad (2)$$

with the initial condition

$$u(x, 0) = f(x) \quad \text{and} \quad \frac{\partial}{\partial t}u(x, 0) = g(x),$$

where  $f$  and  $g$  are given continuous functions. By solving the Cauchy problem for such equation, the Fourier transform has been applied and the solution is given by

$$\hat{u}(\xi, t) = \hat{f}(\xi) \cos(2\pi|\xi|t) + \hat{g}(\xi) \frac{\sin(2\pi|\xi|t)}{2\pi|\xi|}$$

where  $|\xi|^2 = \xi_1^2 + \xi_2^2 + \dots + \xi_n^2$  (see [2, p. 177]). By using the inverse Fourier transform, we obtain  $u(x, t)$  in the convolution form, that is

$$u(x, t) = f(x) * \Psi_t(x) + g(x) * \Phi_t(x) \quad (3)$$

where  $\Phi_t$  is an inverse Fourier transform of  $\hat{\Phi}_t(\xi) = \frac{\sin(2\pi|\xi|t)}{2\pi|\xi|}$  and  $\Psi_t$  is an inverse Fourier transform of  $\hat{\Psi}_t(\xi) = \cos(2\pi|\xi|t) = \frac{\partial}{\partial t}\hat{\Phi}_t(\xi)$ .

In this paper, we study the equation

$$\frac{\partial^2}{\partial t^2}u(x, t) + c^2 (\square)^k u(x, t) = 0 \quad (4)$$

with  $u(x, 0) = f(x)$  and  $\frac{\partial}{\partial t}u(x, 0) = g(x)$ , where  $c$  is a positive constant,  $k$  is a nonnegative integer,  $f$  and  $g$  are continuous functions and absolutely integrable. The equation (4) is motivated by the heat equation of the form

$$\frac{\partial}{\partial t}u(x, t) = -c^2 (\square)^k u(x, t)$$

(see [3], more general: [1]-[4]). We obtain

$$u(x, t) = f(x) * \Psi_t(x) + g(x) * \Phi_t(x) \quad (5)$$

as a solution of (4) where  $\Phi_t$  is an inverse Fourier transform of

$$\hat{\Phi}_t(\xi) = \frac{\sin c \left( \sqrt{s^2 - r^2} \right)^k t}{c \left( \sqrt{s^2 - r^2} \right)^k}$$

## ON THE ULTRA-HYPERBOLIC WAVE OPERATOR

and  $\Psi_t$  is an inverse Fourier transform of  $\widehat{\Psi}_t(\xi) = \cos c \left( \sqrt{s^2 - r^2} \right)^k t = \frac{\partial}{\partial t} \widehat{\Phi}_t(\xi)$  where  $r^2 = \xi_1^2 + \xi_2^2 + \dots + \xi_p^2$  and  $s^2 = \xi_{p+1}^2 + \xi_{p+2}^2 + \dots + \xi_{p+q}^2$ . Moreover, if we put  $k = 1$  and  $q = 0$  in (4) then (5) reduces to the solution of the  $n$ -dimensional wave equation and also if  $k = 2, n = 1$  and  $q = 0$  in (4) then (5) reduces to the solution of beam equation.

We also study the asymptotic form of  $u(x, t)$  in (5) by using  $\epsilon$  approximation and obtain  $u(x, t) = O(\epsilon^{-n/k})$ .

### 2. Preliminaries

We shall need the following definitions.

**Definition 1.** Let  $f \in L_1(\mathbb{R}^n)$ -the space of integrable function in  $\mathbb{R}^n$ . The Fourier transform of  $f(x)$  is defined by

$$\widehat{f}(\xi) = \frac{1}{(2\pi)^{n/2}} \int_{\mathbb{R}^n} e^{-i(\xi, x)} f(x) dx, \quad (6)$$

where  $\xi = (\xi_1, \xi_2, \dots, \xi_n)$ ,  $x = (x_1, x_2, \dots, x_n) \in \mathbb{R}^n$ ,  $(\xi, x) = \xi_1 x_1 + \xi_2 x_2 + \dots + \xi_n x_n$  is the inner product in  $\mathbb{R}^n$  and  $dx = dx_1 dx_2 \dots dx_n$ .

Also, the inverse of Fourier transform is defined by

$$f(\xi) = \frac{1}{(2\pi)^{n/2}} \int_{\mathbb{R}^n} e^{i(\xi, x)} \widehat{f}(x) dx. \quad (7)$$

**Lemma 2.** Given the function

$$f(x) = \exp \left[ -\sqrt{-\sum_{i=1}^p x_i^2 + \sum_{j=p+1}^{p+q} x_j^2} \right],$$

where  $(x_1, x_2, \dots, x_n) \in \mathbb{R}^n$ ,  $p + q = n$ ,  $\sum_{i=1}^p x_i^2 < \sum_{j=p+1}^{p+q} x_j^2$ . Then

$$\left| \int_{\mathbb{R}^n} f(x) dx \right| \leq \frac{\Omega_p \Omega_q}{2} \cdot \frac{\Gamma(n) \Gamma(\frac{p}{2}) \Gamma(\frac{2-n}{2})}{\Gamma(\frac{2-q}{2})},$$

where  $\Gamma$  denotes the Gamma function. That is  $\int_{\mathbb{R}^n} f(x) dx$  is bounded.

*Proof.*

$$\int_{\mathbb{R}^n} f(x) dx = \int_{\mathbb{R}^n} \exp \left[ -\sqrt{-\sum_{i=1}^p x_i^2 + \sum_{j=p+1}^{p+q} x_j^2} \right] dx.$$

Let us transform to bipolar coordinates defined by

$$x_1 = r\omega_1, \quad x_2 = r\omega_2, \dots, \quad x_p = r\omega_p,$$

$$dx_1 = r d\omega_1, \quad dx_2 = r d\omega_2, \dots, \quad dx_p = r d\omega_p,$$

and

$$x_{p+1} = s\omega_{p+1}, \quad x_{p+2} = s\omega_{p+2}, \dots, \quad x_{p+q} = s\omega_{p+q},$$

$$dx_{p+1} = s d\omega_{p+1}, \quad dx_{p+2} = s d\omega_{p+2}, \dots, \quad dx_{p+q} = s d\omega_{p+q},$$

where  $\omega_1^2 + \omega_2^2 + \dots + \omega_p^2 = 1$  and  $\omega_{p+1}^2 + \omega_{p+2}^2 + \dots + \omega_{p+q}^2 = 1$ . Thus

$$\int_{\mathbb{R}^n} f(x) dx = \int_{\mathbb{R}^n} \exp \left[ -\sqrt{s^2 - r^2} \right] r^{p-1} s^{q-1} dr ds d\Omega_p d\Omega_q,$$

where  $dx = r^{p-1} s^{q-1} dr ds d\Omega_p d\Omega_q$ ,  $d\Omega_p$  and  $d\Omega_q$  are the elements of surface area on the unit sphere in  $\mathbb{R}^p$  and  $\mathbb{R}^q$  respectively,

$$\left| \int_{\mathbb{R}^n} f(x) dx \right| \leq \int_{\mathbb{R}^n} \exp \left[ -\sqrt{s^2 - r^2} \right] r^{p-1} s^{q-1} dr ds d\Omega_p d\Omega_q.$$

By computing directly, we obtain

$$\int_{\mathbb{R}^n} f(x) dx = \Omega_p \Omega_q \int_0^\infty \int_0^s \exp \left[ -\sqrt{s^2 - r^2} \right] r^{p-1} s^{q-1} dr ds,$$

where  $\Omega_p = \frac{2\pi^{p/2}}{\Gamma(p/2)}$  and  $\Omega_q = \frac{2\pi^{q/2}}{\Gamma(q/2)}$ . Thus

$$\left| \int_{\mathbb{R}^n} f(x) dx \right| \leq \Omega_p \Omega_q \int_0^\infty \int_0^s \exp \left[ -\sqrt{s^2 - r^2} \right] r^{p-1} s^{q-1} dr ds.$$

Put  $r = s \sin \theta$ ,  $dr = s \cos \theta d\theta$  and  $0 \leq \theta \leq \frac{\pi}{2}$ ,

$$\begin{aligned} \left| \int_{\mathbb{R}^n} f(x) dx \right| &\leq \Omega_p \Omega_q \int_0^\infty \int_0^s e^{-\sqrt{s^2 - s^2 \sin^2 \theta}} (s \sin \theta)^{p-1} s^{q-1} s \cos \theta d\theta ds \\ &= \Omega_p \Omega_q \int_0^\infty \int_0^s e^{-s \cos \theta} s^{p+q-1} (\sin \theta)^{p-1} \cos \theta d\theta ds. \end{aligned}$$

Put  $y = s \cos \theta$ ,  $ds = \frac{dy}{\cos \theta}$ ,

$$\begin{aligned} \left| \int_{\mathbb{R}^n} f(x) dx \right| &\leq \Omega_p \Omega_q \int_0^{\pi/2} \int_0^\infty e^{-y} \left( \frac{y}{\cos \theta} \right)^{n-1} (\sin \theta)^{p-1} \cos \theta d\theta \frac{dy}{\cos \theta} \\ &= \Omega_p \Omega_q \int_0^{\pi/2} \int_0^\infty e^{-y} y^{n-1} (\cos \theta)^{1-n} (\sin \theta)^{p-1} dy d\theta \\ &= \Omega_p \Omega_q \Gamma(n) \int_0^{\pi/2} (\cos \theta)^{1-n} (\sin \theta)^{p-1} d\theta \\ &= \frac{\Omega_p \Omega_q}{2} \Gamma(n) \beta \left( \frac{p}{2}, \frac{2-n}{2} \right), \end{aligned}$$

$$\left| \int_{\mathbb{R}^n} f(x) dx \right| \leq \frac{\Omega_p \Omega_q}{2} \frac{\Gamma(n) \Gamma(\frac{p}{2}) \Gamma(\frac{2-n}{2})}{\Gamma(\frac{2-n}{2})}.$$

That is  $\int_{\mathbb{R}^n} f(x) dx$  is bounded. □

## ON THE ULTRA-HYPERBOLIC WAVE OPERATOR

### 3. Main Results

**Theorem 3.** *Given the equation*

$$\frac{\partial^2}{\partial t^2} u(x, t) + c^2 (\square)^k u(x, t) = 0 \quad (8)$$

*with initial conditions*

$$u(x, 0) = f(x) \quad \text{and} \quad \frac{\partial}{\partial t} u(x, 0) = g(x), \quad (9)$$

*where  $u(x, t) \in \mathbb{R}^n \times [0, \infty)$ ,  $\square^k$  is the ultra-hyperbolic operator iterated  $k$ -times,  $c$  is a positive constant,  $k$  is a nonnegative integer,  $f$  and  $g$  are continuous functions and absolutely integrable for  $x \in \mathbb{R}^n$ . Then (8) has a unique solution*

$$u(x, t) = f(x) * \Psi_t(x) + g(x) * \Phi_t(x) \quad (10)$$

*and satisfy the condition (9), where  $\Phi_t$  is an inverse Fourier transform of*

$$\widehat{\Phi}_t(\xi) = \frac{\sin c \left( \sqrt{s^2 - r^2} \right)^k t}{c \left( \sqrt{s^2 - r^2} \right)^k}$$

*and  $\Psi_t$  is an inverse Fourier transform of*

$$\widehat{\Psi}_t(\xi) = \cos c \left( \sqrt{s^2 - r^2} \right)^k t = \frac{\partial}{\partial t} \widehat{\Phi}_t(\xi),$$

*where  $r^2 = \xi_1^2 + \xi_2^2 + \dots + \xi_p^2$  and  $s^2 = \xi_{p+1}^2 + \xi_{p+2}^2 + \dots + \xi_{p+q}^2$ .*

*Proof.* By applying the Fourier transform defined by (6) to (8) and obtain

$$\frac{\partial^2}{\partial t^2} \widehat{u}(\xi, t) + c^2 \left( -\xi_1^2 - \xi_2^2 - \dots - \xi_p^2 + \xi_{p+1}^2 + \xi_{p+2}^2 + \dots + \xi_{p+q}^2 \right)^k \widehat{u}(\xi, t) = 0,$$

$$\frac{\partial^2}{\partial t^2} \widehat{u}(\xi, t) + c^2 \left( -\sum_{i=1}^p \xi_i^2 + \sum_{j=p+1}^{p+q} \xi_j^2 \right)^k \widehat{u}(\xi, t) = 0$$

*and let  $s > r$ . Thus we have*

$$\frac{\partial^2}{\partial t^2} \widehat{u}(\xi, t) + c^2 (s^2 - r^2)^k \widehat{u}(\xi, t) = 0$$

$$\widehat{u}(\xi, t) = A(\xi) \cos c \left( \sqrt{s^2 - r^2} \right)^k t + B(\xi) \sin c \left( \sqrt{s^2 - r^2} \right)^k t.$$

*By (9),  $\widehat{u}(\xi, 0) = A(\xi) = \widehat{f}(\xi)$*

$$\begin{aligned} \frac{\partial \widehat{u}(\xi, t)}{\partial t} &= -c \left( \sqrt{s^2 - r^2} \right)^k A(\xi) \sin c \left( \sqrt{s^2 - r^2} \right)^k t \\ &\quad + c \left( \sqrt{s^2 - r^2} \right)^k B(\xi) \cos c \left( \sqrt{s^2 - r^2} \right)^k t, \end{aligned}$$

$$\begin{aligned}
\frac{\partial \widehat{u}(\xi, 0)}{\partial t} &= 0 + c \left( \sqrt{s^2 - r^2} \right)^k B(\xi) = \widehat{g}(\xi), \\
B(\xi) &= \frac{\widehat{g}(\xi)}{c \left( \sqrt{s^2 - r^2} \right)^k}, \\
\widehat{u}(\xi, t) &= \widehat{f}(\xi) \cos c \left( \sqrt{s^2 - r^2} \right)^k t + \frac{\widehat{g}(\xi)}{c \left( \sqrt{s^2 - r^2} \right)^k} \sin c \left( \sqrt{s^2 - r^2} \right)^k t. \quad (11)
\end{aligned}$$

By applying the inverse Fourier transform (11), we obtain the solution  $u(x, t)$  in the convolution form of (8). Now we need to show the existence of  $\Phi_t(x)$  and  $\Psi_t(x)$ .

Let us consider the Fourier transform

$$\widehat{\Phi}_t(x) = \frac{\sin c \left( \sqrt{s^2 - r^2} \right)^k t}{c \left( \sqrt{s^2 - r^2} \right)^k} \quad \text{and} \quad \Psi_t(x) = \cos c \left( \sqrt{s^2 - r^2} \right)^k t.$$

They are all tempered distributions but they are not  $L_1(\mathbb{R}^n)$  the space of integrable function. So we cannot compute the inverse Fourier transform  $\Phi_t(x)$  and  $\Psi_t(x)$  directly. Thus we compute the inverse  $\Phi_t(x)$  and  $\Psi_t(x)$  by using the method of  $\epsilon$ -approximation.

Let us define

$$\begin{aligned}
\widehat{\phi}_t^\epsilon(\xi) &= e^{-\epsilon c \left( \sqrt{s^2 - r^2} \right)^k} \widehat{\phi}_t(\xi) = e^{-\epsilon c \left( \sqrt{s^2 - r^2} \right)^k} \frac{\sin c \left( \sqrt{s^2 - r^2} \right)^k t}{c \left( \sqrt{s^2 - r^2} \right)^k} \\
&\quad \text{for } \epsilon > 0. \quad (12)
\end{aligned}$$

We see that  $\phi_t^\epsilon(x) \in L_1(\mathbb{R}^n)$  and  $\widehat{\phi}_t^\epsilon(x) \rightarrow \widehat{\phi}_t(x)$  uniformly as  $\epsilon \rightarrow 0$ . So that  $\phi_t(x)$  will be limit in the topology of tempered distribution of  $\phi_t^\epsilon(x)$ . Now

$$\begin{aligned}
\Phi_t^\epsilon(x) &= \frac{1}{(2\pi)^{n/2}} \int_{\mathbb{R}^n} e^{i(\xi, x)} \widehat{\Phi}_t^\epsilon(\xi) d\xi \\
&= \frac{1}{(2\pi)^{n/2}} \int_{\mathbb{R}^n} e^{i(\xi, x)} e^{-\epsilon c \left( \sqrt{s^2 - r^2} \right)^k} \frac{\sin c \left( \sqrt{s^2 - r^2} \right)^k t}{c \left( \sqrt{s^2 - r^2} \right)^k} d\xi \\
|\Phi_t^\epsilon(x)| &\leq \frac{1}{(2\pi)^{n/2}} \int_{\mathbb{R}^n} \frac{e^{-\epsilon c \left( \sqrt{s^2 - r^2} \right)^k}}{c \left( \sqrt{s^2 - r^2} \right)^k} d\xi. \quad (13)
\end{aligned}$$

# ON THE ULTRA-HYPERBOLIC WAVE OPERATOR

By changing to bipolar coordinates. Now, put

$$\xi_1 = rw_1, \xi_2 = rw_2, \dots, \xi_p = rw_p$$

and

$$\xi_{p+1} = sw_{p+1}, \xi_{p+2} = sw_{p+2}, \dots, \xi_p = sw_{p+q}, \quad p+q=n,$$

where  $w_1^2 + w_2^2 + \dots + w_p^2 = 1$  and  $w_{p+1}^2 + w_{p+2}^2 + \dots + w_{p+q}^2 = 1$ .

$$|\Phi_i^\epsilon(x)| \leq \frac{1}{(2\pi)^{n/2}} \int_{\mathbb{R}^n} \frac{e^{-\epsilon c(\sqrt{s^2-r^2})^k}}{c(\sqrt{s^2-r^2})^k} r^{p-1} s^{q-1} dr ds d\Omega_p d\Omega_q,$$

where  $d\xi = r^{p-1} s^{q-1} dr ds d\Omega_p d\Omega_q$ ,  $d\Omega_p$  and  $d\Omega_q$  are the elements of surface area of the unit sphere in  $\mathbb{R}^p$  and  $\mathbb{R}^q$  respectively, where  $\Omega_p = \frac{2\pi^{p/2}}{\Gamma(p/2)}$ ,  $\Omega_q = \frac{2\pi^{q/2}}{\Gamma(q/2)}$ ,

$$|\Phi_i^\epsilon(x)| \leq \frac{\Omega_p \Omega_q}{(2\pi)^{n/2}} \int_0^\infty \int_0^s \frac{e^{-\epsilon c(\sqrt{s^2-r^2})^k}}{c(\sqrt{s^2-r^2})^k} r^{p-1} s^{q-1} dr ds.$$

Put  $r = s \sin \theta$ ,  $dr = s \cos \theta d\theta$  and  $0 \leq \theta \leq \frac{\pi}{2}$ .

$$\begin{aligned} |\Phi_i^\epsilon(x)| &\leq \frac{\Omega_p \Omega_q}{(2\pi)^{n/2}} \int_0^\infty \int_0^{\pi/2} \frac{e^{-\epsilon c(\sqrt{s^2-s^2 \sin^2 \theta})^k}}{c(\sqrt{s^2-s^2 \sin^2 \theta})^k} (s \sin \theta)^{p-1} s^{q-1} s \cos \theta d\theta ds, \\ &= \frac{\Omega_p \Omega_q}{c(2\pi)^{n/2}} \int_0^\infty \int_0^{\pi/2} \frac{e^{-\epsilon c(s \cos \theta)^k}}{(s \cos \theta)^k} (s)^{p-1} s^{q-1} s (\sin \theta)^{p-1} \cos \theta d\theta ds. \end{aligned}$$

Put  $y = \epsilon c(s \cos \theta)^k = \epsilon c s^k \cos^k \theta$ ,  $s^k = \frac{y}{\epsilon c \cos^k \theta}$ ,  $ds = \frac{dy}{\epsilon c k s^{k-1} \cos^k \theta} = \frac{s dy}{k y}$ , thus

$$\begin{aligned} |\Phi_i^\epsilon(x)| &\leq \frac{\Omega_p \Omega_q}{c(2\pi)^{n/2}} \int_0^{\pi/2} \int_0^\infty \frac{e^{-y} s^{n-1}}{y/(\epsilon c)} (\sin \theta)^{p-1} \cos \theta \frac{s}{k y} dy d\theta \\ &= \frac{\Omega_p \Omega_q}{(2\pi)^{n/2}} \int_0^{\pi/2} \int_0^\infty \frac{e^{-y} \epsilon}{k y^2} \left( \frac{y}{\epsilon c \cos^k \theta} \right)^{n/k} (\sin \theta)^{p-1} \cos \theta dy d\theta \\ &= \frac{\Omega_p \Omega_q}{(2\pi)^{n/2}} \int_0^{\pi/2} \int_0^\infty \frac{e^{-y} y^{n/k-2}}{c^{n/k} k \epsilon^{n/k-1}} (\sin \theta)^{p-1} (\cos \theta)^{1-n} dy d\theta \\ &= \frac{\Omega_p \Omega_q}{(2\pi)^{n/2}} \frac{\Gamma(\frac{n}{k}-1)}{k \epsilon^{\frac{n}{k}-1} c^{n/k}} \int_0^{\pi/2} (\sin \theta)^{p-1} (\cos \theta)^{1-n} d\theta \\ &= \frac{\Omega_p \Omega_q}{2 c^{n/k} (2\pi)^{n/2} k \epsilon^{n/k-1}} \Gamma\left(\frac{n}{k}-1\right) \beta\left(\frac{p}{2}, \frac{2-n}{2}\right), \\ |\Phi_i^\epsilon(x)| &\leq \frac{\Omega_p \Omega_q}{2 c^{n/k} (2\pi)^{n/2} k \epsilon^{n/k-1}} \frac{\Gamma\left(\frac{n}{k}-1\right) \Gamma\left(\frac{p}{2}\right) \Gamma\left(\frac{2-n}{2}\right)}{\Gamma\left(\frac{2-q}{2}\right)}. \end{aligned}$$

Similarly, we defined  $\widehat{\Psi}_t^\epsilon(\xi) = e^{-\epsilon c(\sqrt{s^2-r^2})^k} \cos c(\sqrt{s^2-r^2})^k t$  and

$$\begin{aligned}\Psi_t^\epsilon(x) &= \frac{1}{(2\pi)^{n/2}} \int_{\mathbb{R}^n} e^{i(\xi, x)} \widehat{\Psi}_t^\epsilon(\xi) d\xi \\ &= \frac{1}{(2\pi)^{n/2}} \int_{\mathbb{R}^n} e^{i(\xi, x)} e^{-\epsilon c(\sqrt{s^2-r^2})^k} \cos c(\sqrt{s^2-r^2})^k t d\xi, \\ |\Psi_t^\epsilon(x)| &\leq \frac{1}{(2\pi)^{n/2}} \int_{\mathbb{R}^n} e^{-\epsilon c(\sqrt{s^2-r^2})^k} d\xi \\ &= \frac{1}{(2\pi)^{n/2}} \int_0^\infty \int_0^s e^{-\epsilon c(\sqrt{s^2-r^2})^k} r^{p-1} s^{q-1} dr ds,\end{aligned}$$

Put  $r = s \sin \theta$ ,  $dr = s \cos \theta d\theta$  and  $0 \leq \theta \leq \frac{\pi}{2}$

$$\begin{aligned}|\Psi_t^\epsilon(x)| &\leq \frac{\Omega_p \Omega_q}{(2\pi)^{n/2}} \int_0^\infty \int_0^{\pi/2} e^{-\epsilon c(s \cos \theta)^k} (s \sin \theta)^{p-1} s^{q-1} s \cos \theta d\theta ds \\ &= \frac{\Omega_p \Omega_q}{(2\pi)^{n/2}} \int_0^\infty \int_0^{\pi/2} e^{-\epsilon c(s \cos \theta)^k} s^{p+q-1} (\sin \theta)^{p-1} \cos \theta d\theta ds.\end{aligned}$$

Put  $y = \epsilon c(s \cos \theta)^k$ ,  $ds = s \frac{dy}{ky}$ ,

$$\begin{aligned}|\Psi_t^\epsilon(x)| &\leq \frac{\Omega_p \Omega_q}{k(2\pi)^{n/2}} \int_0^{\pi/2} \int_0^\infty \frac{e^{-y}}{y} \left( \frac{y}{\epsilon c \cos^k \theta} \right)^{n/k} (\sin \theta)^{p-1} \cos \theta dy d\theta \\ &= \frac{\Omega_p \Omega_q}{k(2\pi)^{n/2}} \int_0^{\pi/2} \int_0^\infty \frac{e^{-y} y^{n/k-1}}{c^{n/k} \epsilon^{n/k}} (\sin \theta)^{p-1} (\cos \theta)^{1-n} dy d\theta \\ &= \frac{\Omega_p \Omega_q}{(2\pi)^{n/2} k c^{n/k} \epsilon^{n/k}} \Gamma\left(\frac{n}{k}\right) \int_0^{\pi/2} (\sin \theta)^{p-1} (\cos \theta)^{1-n} d\theta, \\ |\Psi_t^\epsilon(x)| &\leq \frac{\Omega_p \Omega_q}{2(2\pi)^{n/2} k c^{n/k} \epsilon^{n/k}} \frac{\Gamma\left(\frac{n}{k}\right) \Gamma\left(\frac{p}{2}\right) \Gamma\left(\frac{2-n}{2}\right)}{\Gamma\left(\frac{2-q}{2}\right)}.\end{aligned}$$

Set

$$u^\epsilon(x, t) = f(x) * \Psi_t^\epsilon(x) + g(x) * \Phi_t^\epsilon(x) \quad (14)$$

which is  $\epsilon$ -approximation of  $u(x, t)$  in (14) for  $\epsilon \rightarrow 0$ ,  $u^\epsilon(x, t) \rightarrow u(x, t)$  uniformly. Now

$$u^\epsilon(x, t) = \int_{\mathbb{R}^n} f(r) \Psi_t^\epsilon(x-r) dr + \int_{\mathbb{R}^n} g(r) \Phi_t^\epsilon(x-r) dr.$$

Thus

$$|u^\epsilon(x, t)| \leq |\Psi_t^\epsilon(x-r)| \int_{\mathbb{R}^n} |f(r)| dr + |\Phi_t^\epsilon(x-r)| \int_{\mathbb{R}^n} |g(r)| dr$$

## ON THE ULTRA-HYPERBOLIC WAVE OPERATOR

$$\begin{aligned}
 &\leq \frac{\Omega_p \Omega_q}{(2\pi)^{n/2} k c^{n/k} \epsilon^{n/k}} \frac{\Gamma\left(\frac{n}{k}\right) \Gamma\left(\frac{p}{2}\right) \Gamma\left(\frac{2-n}{2}\right)}{\Gamma\left(\frac{2-q}{2}\right)} M \\
 &\quad + \frac{\Omega_p \Omega_q}{(2\pi)^{n/2} k c^{n/k} \epsilon^{n/k-1}} \frac{\Gamma\left(\frac{n}{k} - 1\right) \Gamma\left(\frac{p}{2}\right) \Gamma\left(\frac{2-n}{2}\right)}{\Gamma\left(\frac{2-q}{2}\right)} N, \\
 \epsilon^{n/k} |u^\epsilon(x, t)| &\leq \frac{\Omega_p \Omega_q}{2(2\pi)^{n/2} k c^{n/k}} \frac{\Gamma\left(\frac{n}{k}\right) \Gamma\left(\frac{p}{2}\right) \Gamma\left(\frac{2-n}{2}\right)}{\Gamma\left(\frac{2-q}{2}\right)} M \\
 &\quad + \frac{\Omega_p \Omega_q \epsilon}{2(2\pi)^{n/2} k c^{n/k}} \frac{\Gamma\left(\frac{n}{k} - 1\right) \Gamma\left(\frac{p}{2}\right) \Gamma\left(\frac{2-n}{2}\right)}{\Gamma\left(\frac{2-q}{2}\right)} N,
 \end{aligned}$$

where  $M = \int_{\mathbb{R}^n} |f(r)| dr$  and  $N = \int_{\mathbb{R}^n} |g(r)| dr$ , since  $f$  and  $g$  are absolutely integrable.

$$\lim_{\epsilon \rightarrow 0} \epsilon^{n/k} |u^\epsilon(x, t)| \leq \frac{\Omega_p \Omega_q}{2(2\pi)^{n/2} k c^{n/k}} \frac{\Gamma\left(\frac{n}{k}\right) \Gamma\left(\frac{p}{2}\right) \Gamma\left(\frac{2-n}{2}\right)}{\Gamma\left(\frac{2-q}{2}\right)} = K.$$

It follows that  $u(x, t) = O(\epsilon^{-n/k})$  for  $n \neq k$  as  $\epsilon \rightarrow 0$ .

In particular, if we put  $k = 2, n = 1$  and  $q = 0$  then (8) reduces to the solution of the beam equation, see [1, p. 47]

$$\frac{\partial^2}{\partial t^2} u(x, t) + c^2 \frac{\partial^4}{\partial x^4} u(x, t) = 0,$$

with the initial conditions

$$u(x, 0) = f(x) \quad \text{and} \quad \frac{\partial}{\partial t} u(x, 0) = g(x),$$

where  $f$  and  $g$  are continuous and absolutely integrable for  $x \in \mathbb{R}^n$ . Thus we obtain  $u(x, t) = O(\epsilon^{-1/2})$  which is a solution of such beam equation.

## Acknowledgements

The authors would like to thank The Thailand Research Fund for financial support.

W. Satsanit, A. Kananthai

### References

- [1] J. David Logan, *An Introduction to Nonlinear Partial Differential Equations*, A Wiley-Interscience Publication, John Wiley and Sons (1997).
- [2] G.B. Folland, *Introduction to Partial Differential Equation*, Princeton University Press, Princeton, New Jersey (1995).
- [3] A. Kanathai, K. Nonlaopon, On the generalized heat kernel, *Computational Technologies*, **9**, No. 1 (2004), 3-10.



## DIAMOND OPERATOR RELATED TO BIHARMONIC EQUATIONS

WANCHAK SATSANIT and AMNUAY KANANTHAI

Department of Mathematics  
Chiangmai University  
Chiangmai, 50200, Thailand  
e-mail: [malamka@science.cmu.ac.th](mailto:malamka@science.cmu.ac.th)

### Abstract

In this paper, we study the generalized wave equation of the form

$$\frac{\partial^2}{\partial t^2} u(x, t) - c^2 (\diamond)^k u(x, t) = 0$$

with the initial conditions

$$u(x, 0) = f(x), \quad \frac{\partial}{\partial t} u(x, 0) = g(0),$$

where  $u(x, t) \in \mathbb{R}^n \times [0, \infty)$ ,  $\mathbb{R}^n$  is the  $n$ -dimensional Euclidean space,

$\diamond^k$  is the Diamond operator iterated  $k$ -times defined by

$$\diamond^k = \left[ \left( \sum_{i=1}^p \frac{\partial^2}{\partial x_i^2} \right)^2 - \left( \sum_{i=p+1}^{p+q} \frac{\partial^2}{\partial x_i^2} \right)^2 \right]^k,$$

$\diamond$  can be written as the product of the operators in the form  $\diamond = \Delta \square$

$$= \square \Delta, \quad \text{where } \Delta = \sum_{i=1}^n \frac{\partial^2}{\partial x_i^2} \text{ is the Laplacian and } \square = \sum_{i=1}^p \frac{\partial^2}{\partial x_i^2}$$

2000 Mathematics Subject Classification: Kindly provide.

Keywords and phrases: biharmonic wave equation, Diamond operator, tempered distribution.

Received March 27, 2009

$-\sum_{j=p+1}^{p+q} \frac{\partial^2}{\partial x_j^2}$  is the ultra-hyperbolic.  $p+q=n$ ,  $c$  is a positive constant,

$k$  is a nonnegative integer,  $f$  and  $g$  are continuous and absolutely integrable functions. We obtain  $u(x, t)$  as a solution for such equation. Moreover, by  $\varepsilon$ -approximation we also obtain the asymptotic solution  $u(x, t) = O(\varepsilon^{-n/2k})$ . In particular, if we put  $n=1$ ,  $k=2$  and  $p=0$ , the  $u(x, t)$  reduces to the solution of the biharmonic wave equation

$$\frac{\partial^2}{\partial t^2} u(x, t) + c^2 (\Delta)^4 u(x, t) = 0.$$

### 1. Introduction

It is well known that for the 1-dimensional wave equation

$$\frac{\partial^2}{\partial t^2} u(x, t) = c^2 \frac{\partial^2}{\partial x^2} u(x, t), \quad (1.1)$$

we obtain  $u(x, t) = f(x+ct) + g(x-ct)$  as a solution of the equation where  $f$  and  $g$  are continuous.

Also for the  $n$ -dimensional wave equation

$$\frac{\partial^2}{\partial t^2} u(x, t) + c^2 \Delta u(x, t) = 0, \quad (1.2)$$

with the initial condition

$$u(x, 0) = f(x) \quad \text{and} \quad \frac{\partial}{\partial t} u(x, 0) = g(x),$$

where  $f$  and  $g$  are given continuous functions. By solving the Cauchy problem for such equation, the Fourier transform has been applied and the solution is given by

$$\hat{u}(\xi, t) = \hat{f}(\xi) \cos(2\pi |\xi| t) + \hat{g}(\xi) \frac{\sin(2\pi |\xi| t)}{2\pi |\xi|},$$

where  $r^2 = \xi_1^2 + \xi_2^2 + \dots + \xi_p^2$ ,  $s^2 = \xi_{p+1}^2 + \xi_{p+2}^2 + \dots + \xi_{p+q}^2$  (see [1, p. 177]).

By using the inverse Fourier transform, we obtain  $u(x, t)$  in the convolution form,

# DIAMOND OPERATOR RELATED TO BIHARMONIC EQUATION

that is,

$$u(x, t) = f(x) * \Psi_t(x) + g(x) * \Phi_t(x), \quad (1.3)$$

where  $\Phi_t$  is an inverse Fourier transform of  $\hat{\Phi}_t(\xi) = \frac{\sin(2\pi|\xi|t)}{2\pi|\xi|}$  and  $\Psi_t$  is an inverse Fourier transform of  $\hat{\Psi}_t(\xi) = \cos(2\pi|\xi|t) = \frac{\partial}{\partial t} \hat{\Phi}_t(\xi)$ .

In 1996, Kananthai [2] introduced the *Diamond operator*  $\diamond$  defined by

$$\diamond = \left( \sum_{i=1}^p \frac{\partial^2}{\partial x_i^2} \right)^2 - \left( \sum_{j=p+1}^{p+q} \frac{\partial^2}{\partial x_j^2} \right)^2, \quad p + q = n$$

or  $\diamond$  can be written as the product of the operators in the form  $\diamond = \Delta \square = \square \Delta$

where  $\Delta = \sum_{i=1}^n \frac{\partial^2}{\partial x_i^2}$  is the Laplacian and  $\square = \sum_{i=1}^p \frac{\partial^2}{\partial x_i^2} - \sum_{j=p+1}^{p+q} \frac{\partial^2}{\partial x_j^2}$  is the ultra-

hyperbolic. The Fourier transform of the Diamond operator has also been studied and the elementary solution of such operator, see [3]. Next, G. Sritantana, A. Kananthai study the equation

$$\frac{\partial^2}{\partial t^2} u(x, t) + c^2 (-\Delta)^k u(x, t) = 0$$

see [7, pp. 23-29], where

$$\Delta^k = \left( \frac{\partial^2}{\partial x_1^2} + \frac{\partial^2}{\partial x_2^2} + \dots + \frac{\partial^2}{\partial x_p^2} + \frac{\partial^2}{\partial x_{p+1}^2} + \frac{\partial^2}{\partial x_{p+2}^2} + \dots + \frac{\partial^2}{\partial x_{p+q}^2} \right)^k.$$

Next, W. Satsanit, A. Kananthai study the equation

$$\frac{\partial^2}{\partial t^2} u(x, t) + c^2 (\square)^k u(x, t) = 0$$

see [6], where

$$\square^k = \left( \frac{\partial^2}{\partial x_1^2} + \frac{\partial^2}{\partial x_2^2} + \dots + \frac{\partial^2}{\partial x_p^2} - \frac{\partial^2}{\partial x_{p+1}^2} - \frac{\partial^2}{\partial x_{p+2}^2} - \dots - \frac{\partial^2}{\partial x_{p+q}^2} \right)^k,$$

we obtain the solution related to the beam equation.

In this paper, we study the equation

$$\frac{\partial^2}{\partial t^2} u(x, t) + c^2 (\diamond)^k u(x, t) = 0 \quad (1.4)$$

with  $u(x, 0) = f(x)$  and  $\partial/\partial t u(x, 0) = g(x)$ , where  $c$  is a positive constant,  $k$  is a nonnegative integer,  $f$  and  $g$  are continuous functions and absolutely integrable. The equation (1.4) is motivated by the heat equation of the form

$$\frac{\partial}{\partial t} u(x, t) = -c^2 (\diamond)^k u(x, t)$$

(see [4, 1-4]). We obtain

$$u(x, t) = f(x) * \Psi_t(x) + g(x) * \Phi_t(x) \quad (1.5)$$

as a solution of (1.4), where  $\Phi_t$  is an inverse Fourier transform of  $\hat{\Phi}_t(\xi)$

$$= \frac{\sin c(\sqrt{s^4 - r^4})^k t}{c(\sqrt{s^4 - r^4})^k} \quad \text{and} \quad \Psi_t \text{ is an inverse Fourier transform of } \hat{\Psi}_t(\xi)$$

$$= \cos c(\sqrt{s^4 - r^4})^k t = \frac{\partial}{\partial t} \hat{\Phi}_t(\xi), \quad \text{where } r^2 = \xi_1^2 + \xi_2^2 + \dots + \xi_p^2 \quad \text{and } s^2 = \xi_{p+1}^2$$

$$+ \xi_{p+2}^2 + \dots + \xi_{p+q}^2. \quad \text{Moreover, if we put } k = 2 \text{ and } p = 0 \text{ in (1.4), then (1.5)}$$

reduces to the solution of the  $n$ -dimensional biharmonic wave equation and also if  $k = 1$ ,  $n = 1$  and  $p = 0$  in (1.4), then (1.5) reduces to the solution of beam equation.

We also study the asymptotic form of  $u(x, t)$  in (1.5) by using  $\epsilon$ -approximation and obtain  $u(x, t) = O(\epsilon^{-n/2k})$ .

## 2. Preliminaries

We shall need the following definitions

**Definition 2.1.** Let  $f \in L_1(\mathbb{R}^n)$  the space of integrable function in  $\mathbb{R}^n$ . The Fourier transform of  $f(x)$  is defined by

$$\hat{f}(\xi) = \frac{1}{(2\pi)^{n/2}} \int_{\mathbb{R}^n} e^{i(\xi, x)} f(x) dx, \quad (2.1)$$

## DIAMOND OPERATOR RELATED TO BIHARMONIC EQUATION

where  $\xi = (\xi_1, \xi_2, \dots, \xi_n)$ ,  $x = (x_1, x_2, \dots, x_n) \in \mathbb{R}^n$ ,  $(\xi, x) = \xi_1 x_1 + \xi_2 x_2 + \dots + \xi_n x_n$  is the inner product in  $\mathbb{R}^n$  and  $dx = dx_1 dx_2 \dots dx_n$ .

Also, the inverse of Fourier transform is defined by

$$f(\xi) = \frac{1}{(2\pi)^{n/2}} \int_{\mathbb{R}^n} e^{i(\xi, x)} \hat{f}(x) dx. \quad (2.2)$$

**Lemma 2.1.** *Given the function*

$$f(x) = \exp \left[ - \sqrt{ - \left( \sum_{i=1}^p x_i^2 \right)^2 + \left( \sum_{j=p+1}^{p+q} x_j^2 \right)^2 } \right],$$

where  $(x_1, x_2, \dots, x_n) \in \mathbb{R}^n$ ,  $p + q = n$ ,  $\sum_{i=1}^p x_i^2 < \sum_{j=p+1}^{p+q} x_j^2$ . Then

$$\left| \int_{\mathbb{R}^n} f(x) dx \right| \leq \frac{\Omega_p \Omega_q}{8} \cdot \frac{\Gamma\left(\frac{n}{2}\right) \Gamma\left(\frac{p}{4}\right) \Gamma\left(\frac{4-n}{4}\right)}{\Gamma\left(\frac{4-q}{4}\right)},$$

where  $\Gamma$  denotes the Gamma function. That is,  $\int_{\mathbb{R}^n} f(x) dx$  is bounded.

**Proof.** First note that

$$\int_{\mathbb{R}^n} f(x) dx = \int_{\mathbb{R}^n} \exp \left[ - \sqrt{ - \left( \sum_{i=1}^p x_i^2 \right)^2 + \left( \sum_{j=p+1}^{p+q} x_j^2 \right)^2 } \right] dx.$$

Now, we transform to bipolar coordinates defined by

$$x_1 = r\omega_1, \quad x_2 = r\omega_2, \dots, \quad x_p = r\omega_p,$$

$$dx_1 = r d\omega_1, \quad dx_2 = r d\omega_2, \dots, \quad dx_p = r d\omega_p$$

and

$$x_{p+1} = s\omega_{p+1}, \quad x_{p+2} = s\omega_{p+2}, \dots, \quad x_{p+q} = s\omega_{p+q},$$

$$dx_{p+1} = s d\omega_{p+1}, \quad dx_{p+2} = s d\omega_{p+2}, \dots, \quad dx_{p+q} = s d\omega_{p+q},$$

# DIAMOND OPERATOR RELATED TO BIHARMONIC EQUATION

where  $\xi = (\xi_1, \xi_2, \dots, \xi_n)$ ,  $x = (x_1, x_2, \dots, x_n) \in \mathbb{R}^n$ ,  $(\xi, x) = \xi_1 x_1 + \xi_2 x_2 + \dots + \xi_n x_n$  is the inner product in  $\mathbb{R}^n$  and  $dx = dx_1 dx_2 \dots dx_n$ .

Also, the inverse of Fourier transform is defined by

$$f(\xi) = \frac{1}{(2\pi)^{n/2}} \int_{\mathbb{R}^n} e^{i(\xi, x)} \hat{f}(x) dx. \quad (2.2)$$

**Lemma 2.1.** *Given the function*

$$f(x) = \exp \left[ - \sqrt{ - \left( \sum_{i=1}^p x_i^2 \right)^2 + \left( \sum_{j=p+1}^{p+q} x_j^2 \right)^2 } \right],$$

where  $(x_1, x_2, \dots, x_n) \in \mathbb{R}^n$ ,  $p + q = n$ ,  $\sum_{i=1}^p x_i^2 < \sum_{j=p+1}^{p+q} x_j^2$ . Then

$$\left| \int_{\mathbb{R}^n} f(x) dx \right| \leq \frac{\Omega_p \Omega_q}{8} \cdot \frac{\Gamma\left(\frac{n}{2}\right) \Gamma\left(\frac{p}{4}\right) \Gamma\left(\frac{4-n}{4}\right)}{\Gamma\left(\frac{4-q}{4}\right)},$$

where  $\Gamma$  denotes the Gamma function. That is,  $\int_{\mathbb{R}^n} f(x) dx$  is bounded.

**Proof.** First note that

$$\int_{\mathbb{R}^n} f(x) dx = \int_{\mathbb{R}^n} \exp \left[ - \sqrt{ - \left( \sum_{i=1}^p x_i^2 \right)^2 + \left( \sum_{j=p+1}^{p+q} x_j^2 \right)^2 } \right] dx.$$

Now, we transform to bipolar coordinates defined by

$$x_1 = r\omega_1, \quad x_2 = r\omega_2, \quad \dots, \quad x_p = r\omega_p,$$

$$dx_1 = r d\omega_1, \quad dx_2 = r d\omega_2, \quad \dots, \quad dx_p = r d\omega_p$$

and

$$x_{p+1} = s\omega_{p+1}, \quad x_{p+2} = s\omega_{p+2}, \quad \dots, \quad x_{p+q} = s\omega_{p+q},$$

$$dx_{p+1} = s d\omega_{p+1}, \quad dx_{p+2} = s d\omega_{p+2}, \quad \dots, \quad dx_{p+q} = s d\omega_{p+q},$$

where  $\omega_1^2 + \omega_2^2 + \dots + \omega_p^2 = 1$  and  $\omega_{p+1}^2 + \omega_{p+2}^2 + \omega_{p+q}^2 = 1$ .

Thus

$$\int_{\mathbb{R}^n} f(x) dx = \int_{\mathbb{R}^n} \exp[-\sqrt{s^4 - r^4}] r^{p-1} s^{q-1} dr ds d\Omega_p d\Omega_q,$$

where  $dx = r^{p-1} s^{q-1} dr ds d\Omega_p d\Omega_q$ ,  $d\Omega_p$  and  $d\Omega_q$  are the elements of surface area on the unit sphere in  $\mathbb{R}^p$  and  $\mathbb{R}^q$ , respectively.

$$\left| \int_{\mathbb{R}^n} f(x) dx \right| \leq \int_{\mathbb{R}^n} \exp[-\sqrt{s^4 - r^4}] r^{p-1} s^{q-1} dr ds d\Omega_p d\Omega_q.$$

By a direct computation, we obtain

$$\int_{\mathbb{R}^n} f(x) dx = \Omega_p \Omega_q \int_0^\infty \int_0^s \exp[-\sqrt{s^4 - r^4}] r^{p-1} s^{q-1} dr ds,$$

where  $\Omega_p = \frac{2\pi^{p/2}}{\Gamma(p/2)}$  and  $\Omega_q = \frac{2\pi^{q/2}}{\Gamma(q/2)}$ . Thus

$$\left| \int_{\mathbb{R}^n} f(x) dx \right| \leq \Omega_p \Omega_q \int_0^\infty \int_0^s \exp[-\sqrt{s^4 - r^4}] r^{p-1} s^{q-1} dr ds.$$

Put  $r^2 = s^2 \sin \theta$ ,  $2r dr = s^2 \cos \theta d\theta$  and  $0 \leq \theta \leq \frac{\pi}{2}$ , to have

$$\begin{aligned} \left| \int_{\mathbb{R}^n} f(x) dx \right| &\leq \Omega_p \Omega_q \int_0^\infty \int_0^s e^{-\sqrt{s^4 - s^4 \sin^2 \theta}} s^{p-2} (\sin \theta)^{\frac{p-2}{2}} s^{q+1} \cos \theta d\theta ds \\ &= \frac{\Omega_p \Omega_q}{2} \int_0^\infty \int_0^s e^{-s^2 \cos \theta} s^{p+q-1} (\sin \theta)^{\frac{p-2}{2}} \cos \theta d\theta ds. \end{aligned}$$

Put  $y = s^2 \cos \theta$ ,  $ds = \frac{dy}{2s \cos \theta}$ , to have

$$\begin{aligned} \left| \int_{\mathbb{R}^n} f(x) dx \right| &\leq \frac{\Omega_p \Omega_q}{4} \int_0^{\pi/2} \int_0^\infty e^{-y} \left( \frac{y}{\cos \theta} \right)^{\frac{n-2}{2}} (\sin \theta)^{\frac{p-2}{2}} \cos \theta d\theta \frac{dy}{\cos \theta} \\ &= \frac{\Omega_p \Omega_q}{4} \int_0^{\pi/2} \int_0^\infty e^{-y} y^{\frac{n-2}{2}} (\cos \theta)^{\frac{2-n}{2}} (\sin \theta)^{\frac{p-2}{2}} dy d\theta \end{aligned}$$

# DIAMOND OPERATOR RELATED TO BIHARMONIC EQUATION

$$= \frac{\Omega_p \Omega_q}{4} \Gamma\left(\frac{n}{2}\right) \int_0^{\pi/2} (\cos \theta)^{\frac{2-n}{2}} (\sin \theta)^{\frac{p-2}{2}} d\theta$$

$$= \frac{\Omega_p \Omega_q}{8} \Gamma\left(\frac{n}{2}\right) \beta\left(\frac{p}{4}, \frac{4-n}{4}\right).$$

Therefore,

$$\left| \int_{\mathbb{R}^n} f(x) dx \right| \leq \frac{\Omega_p \Omega_q}{8} \frac{\Gamma\left(\frac{n}{2}\right) \Gamma\left(\frac{p}{4}\right) \Gamma\left(\frac{4-n}{4}\right)}{\Gamma\left(\frac{4-q}{4}\right)}.$$

Thus it follows that  $\int_{\mathbb{R}^n} f(x) dx$  is bounded.

## 3. Main Results

**Theorem 3.1.** *Given the equation*

$$\frac{\partial^2}{\partial t^2} u(x, t) + c^2 (\diamond)^k u(x, t) = 0 \quad (3.1)$$

*with initial conditions*

$$u(x, 0) = f(x) \quad \text{and} \quad \frac{\partial}{\partial t} u(x, 0) = g(x), \quad (3.2)$$

*where  $u(x, t) \in \mathbb{R}^n \times [0, \infty)$ ,  $\diamond^k$  is the Diamond operator iterated  $k$ -times,  $c$  is a positive constant,  $k$  is a nonnegative integer,  $f$  and  $g$  are continuous functions and absolutely integrable for  $x \in \mathbb{R}^n$ . Then (3.1) has a unique solution*

$$u(x, t) = f(x) * \Psi_t(x) + g(x) * \Phi_t(x) \quad (3.3)$$

*and satisfies the condition (3.2) where  $\Phi_t$  is the inverse Fourier transform of*

$$\hat{\Phi}_t(\xi) = \frac{\sin c(\sqrt{s^4 - r^4})^k t}{c(\sqrt{s^4 - r^4})^k}$$

*and  $\Psi_t$  is the inverse Fourier transform of*

$$\hat{\Psi}_t(\xi) = \cos c(\sqrt{s^4 - r^4})^k t = \frac{\partial}{\partial t} \hat{\Phi}(\xi),$$

*with  $r^2 = \xi_1^2 + \xi_2^2 + \dots + \xi_p^2$  and  $s^2 = \xi_{p+1}^2 + \xi_{p+2}^2 + \dots + \xi_{p+q}^2$ .*

**Proof.** By applying the Fourier transform defined by (2.1) to (3.1), we obtain

$$\frac{\partial^2}{\partial t^2} \hat{u}(\xi, t) + c^2 \left( - \left( \sum_{i=1}^p \xi_i^2 \right) + \left( \sum_{j=p+1}^{p+q} \xi_j^2 \right) \right)^k \hat{u}(\xi, t) = 0.$$

Let  $s > r$ . Thus

$$\frac{\partial^2}{\partial t^2} \hat{u}(\xi, t) + c^2 (s^4 - r^4)^k \hat{u}(\xi, t) = 0,$$

$$\hat{u}(\xi, t) = A(\xi) \cos c(\sqrt{s^4 - r^4})^k t + B(\xi) \sin c(\sqrt{s^4 - r^4})^k t.$$

By (3.2),  $\hat{u}(\xi, 0) = A(\xi) = \hat{f}(\xi)$ ,

$$\begin{aligned} \frac{\partial \hat{u}(\xi, t)}{\partial t} &= -c(\sqrt{s^4 - r^4})^k A(\xi) \sin c(\sqrt{s^4 - r^4})^k t \\ &\quad + c(\sqrt{s^4 - r^4})^k B(\xi) \cos c(\sqrt{s^4 - r^4})^k t. \end{aligned}$$

$$\frac{\partial \hat{u}(\xi, 0)}{\partial t} = 0 + c(\sqrt{s^4 - r^4})^k B(\xi) = \hat{g}(\xi),$$

$$B(\xi) = \frac{\hat{g}(\xi)}{c(\sqrt{s^4 - r^4})^k},$$

$$\hat{u}(\xi, t) = \hat{f}(\xi) \cos c(\sqrt{s^4 - r^4})^k t + \frac{\hat{g}(\xi)}{c(\sqrt{s^4 - r^4})^k} \sin c(\sqrt{s^4 - r^4})^k t. \quad (3.4)$$

By applying the inverse Fourier transform (3.4), we obtain the solution  $u(x, t)$  in the convolution form of (3.1). Now, we need to show the existence of  $\Phi_t(x)$  and  $\Psi_t(x)$ . Consider the Fourier transforms

$$\widehat{\Phi}_t(x) = \frac{\sin c(\sqrt{s^4 - r^4})^k t}{c(\sqrt{s^4 - r^4})^k} \quad \text{and} \quad \Psi_t(x) = \cos c(\sqrt{s^4 - r^4})^k t.$$

These are all tempered distributions not lying in the space  $L_1(\mathbb{R}^n)$  of integrable functions. So we cannot compute the inverse Fourier transforms  $\Phi_t(x)$  and  $\Psi_t(x)$

# DIAMOND OPERATOR RELATED TO BIHARMONIC EQUATION

directly. Thus we compute the inverse  $\Phi_t(x)$  and  $\Psi_t(x)$  by using the method of  $\varepsilon$ -approximation.

Define

$$\widehat{\phi}_t^\varepsilon(\xi) = e^{-\varepsilon c(\sqrt{s^4 - r^4})^k} \widehat{\phi}_t(\xi) = e^{-\varepsilon c(\sqrt{s^4 - r^4})^k} \frac{\sin c(\sqrt{s^4 - r^4})^k t}{c(\sqrt{s^4 - r^4})^k} \text{ for } \varepsilon > 0. \quad (3.5)$$

We see that  $\phi_t^\varepsilon(x) \in L_1(\mathbb{R}^n)$  and  $\widehat{\phi}_t^\varepsilon(x) \rightarrow \widehat{\phi}_t(x)$  uniformly as  $\varepsilon \rightarrow 0$ . So that  $\phi_t(x)$  will be limit in the topology of tempered distribution of  $\phi_t^\varepsilon(x)$ . Now

$$\begin{aligned} \Phi_t^\varepsilon(x) &= \frac{1}{(2\pi)^{n/2}} \int_{\mathbb{R}^n} e^{i(\xi, x)} \widehat{\phi}_t^\varepsilon(\xi) d\xi \\ &= \frac{1}{(2\pi)^{n/2}} \int_{\mathbb{R}^n} e^{i(\xi, x)} e^{-\varepsilon c(\sqrt{s^4 - r^4})^k} \frac{\sin c(\sqrt{s^4 - r^4})^k t}{c(\sqrt{s^4 - r^4})^k} d\xi, \\ |\Phi_t^\varepsilon(x)| &\leq \frac{1}{(2\pi)^{n/2}} \int_{\mathbb{R}^n} \frac{e^{-\varepsilon c(\sqrt{s^4 - r^4})^k}}{c(\sqrt{s^4 - r^4})^k} d\xi. \end{aligned} \quad (3.6)$$

By changing to bipolar coordinates and putting

$$\xi_1 = rw_1, \quad \xi_2 = rw_2, \dots, \quad \xi_p = rw_p,$$

and

$$\xi_{p+1} = sw_{p+1}, \quad \xi_{p+2} = sw_{p+2}, \dots, \quad \xi_p = sw_{p+q}, \quad p + q = n,$$

where  $w_1^2 + w_2^2 + \dots + w_p^2 = 1$  and  $w_{p+1}^2 + w_{p+2}^2 + \dots + w_{p+q}^2 = 1$ , we obtain

$$|\Phi_t^\varepsilon(x)| \leq \frac{1}{(2\pi)^{n/2}} \int_{\mathbb{R}^n} \frac{e^{-\varepsilon c(\sqrt{s^4 - r^4})^k}}{c(\sqrt{s^4 - r^4})^k} r^{p-1} s^{q-1} dr ds d\Omega_p d\Omega_q,$$

where  $d\xi = r^{p-1} s^{q-1} dr ds d\Omega_p d\Omega_q$ ,  $d\Omega_p$  and  $d\Omega_q$  are the elements of surface area

of the unit spheres in  $\mathbb{R}^p$  and  $\mathbb{R}^q$ , respectively, with  $\Omega_p = \frac{(2\pi)^{p/2}}{\Gamma(p/2)}$ ,  $\Omega_q =$

$\frac{(2\pi)^{q/2}}{\Gamma(q/2)}$ . Now,

$$|\Phi_t^\varepsilon(x)| \leq \frac{\Omega_p \Omega_q}{(2\pi)^{n/2}} \int_0^\infty \int_0^s \frac{e^{-\varepsilon c(\sqrt{s^4-r^4})^k}}{c(\sqrt{s^4-r^4})^k} r^{p-1} s^{q-1} dr ds.$$

Putting  $r^2 = s^2 \sin \theta$ ,  $2rdr = s^2 \cos \theta d\theta$  and  $0 \leq \theta \leq \frac{\pi}{2}$ , we get

$$\begin{aligned} |\Phi_t^\varepsilon(x)| &\leq \frac{\Omega_p \Omega_q}{2(2\pi)^{n/2}} \int_0^\infty \int_0^{\pi/2} \frac{e^{-\varepsilon c(\sqrt{s^4-s^4 \sin^2 \theta})^k}}{c(\sqrt{s^4-s^4 \sin^2 \theta})^k} (\sin \theta)^{\frac{p-2}{2}} s^{p+q-1} \cos \theta d\theta ds \\ &= \frac{\Omega_p \Omega_q}{2c(2\pi)^{n/2}} \int_0^\infty \int_0^{\pi/2} \frac{e^{-\varepsilon c(s^2 \cos \theta)^k}}{c(s^2 \cos \theta)^k} s^{p+q-1} (\sin \theta)^{\frac{p-2}{2}} \cos \theta d\theta ds. \end{aligned}$$

Putting  $y = \varepsilon c(s^2 \cos \theta)^k = \varepsilon c s^{2k} \cos^k \theta$ ,  $s^{2k} = \frac{y}{c\varepsilon \cos^k \theta}$ ,  $ds = \frac{s dy}{2ky}$ , it follows

that

$$\begin{aligned} |\Phi_t^\varepsilon(x)| &\leq \frac{\Omega_p \Omega_q}{4c(2\pi)^{n/2}} \int_0^{\pi/2} \int_0^\infty \frac{e^{-y} s^{n-1}}{y/(\varepsilon c)} (\sin \theta)^{\frac{p-2}{2}} \cos \theta \frac{s}{ky} dy d\theta \\ &= \frac{\Omega_p \Omega_q}{4(2\pi)^{n/2}} \int_0^{\pi/2} \int_0^\infty \frac{e^{-y} \varepsilon}{ky^2} \left( \frac{y}{c\varepsilon \cos^k \theta} \right)^{n/2k} (\sin \theta)^{p-2/2} \cos \theta dy d\theta \\ &= \frac{\Omega_p \Omega_q}{4(2\pi)^{n/2}} \int_0^{\pi/2} \int_0^\infty \frac{e^{-y} y^{n/2k-2} \varepsilon}{c^{n/2k} k \varepsilon^{n/2k-1}} (\sin \theta)^{\frac{p-2}{2}} (\cos \theta)^{\frac{2-n}{2}} dy d\theta \\ &= \frac{\Omega_p \Omega_q}{4(2\pi)^{n/2}} \frac{\Gamma\left(\frac{n}{2k}-1\right)}{k \varepsilon^{\frac{n}{2k}-1} c^{n/2k}} \int_0^{\pi/2} (\sin \theta)^{\frac{p-2}{2}} (\cos \theta)^{\frac{2-n}{2}} d\theta \\ &= \frac{\Omega_p \Omega_q}{8c^{n/2k} (2\pi)^{n/2} k \varepsilon^{n/2k-1}} \Gamma\left(\frac{n}{2k}-1\right) \beta\left(\frac{p}{4}, \frac{4-n}{4}\right), \end{aligned}$$

and

$$|\Phi_t^\varepsilon(x)| \leq \frac{\Omega_p \Omega_q}{8c^{n/2k} (2\pi)^{n/2} k \varepsilon^{n/2k-1}} \frac{\Gamma\left(\frac{n}{2k}-1\right) \Gamma\left(\frac{p}{4}\right) \Gamma\left(\frac{4-n}{4}\right)}{\Gamma\left(\frac{4-q}{4}\right)}.$$

Similarly, we define  $\widehat{\Psi}_t^\varepsilon(\xi) = e^{-\varepsilon c(\sqrt{s^4 - r^4})^k} \cos c(\sqrt{s^4 - r^4})^k t$  and

$$\begin{aligned} \Psi_t^\varepsilon(x) &= \frac{1}{(2\pi)^{n/2}} \int_{\mathbb{R}^n} e^{i(\xi, x)} \widehat{\Psi}_t^\varepsilon(\xi) d\xi \\ &= \frac{1}{(2\pi)^{n/2}} \int_{\mathbb{R}^n} e^{i(\xi, x)} e^{-\varepsilon c(\sqrt{s^4 - r^4})^k} \cos c(\sqrt{s^4 - r^4})^k t d\xi, \\ |\Psi_t^\varepsilon(x)| &\leq \frac{1}{(2\pi)^{n/2}} \int_{\mathbb{R}^n} e^{-\varepsilon c(\sqrt{s^4 - r^4})^k} d\xi \\ &= \frac{1}{(2\pi)^{n/2}} \int_0^\infty \int_0^s e^{-\varepsilon c(\sqrt{s^4 - r^4})^k} r^{p-1} s^{q-1} dr ds. \end{aligned}$$

Putting  $r^2 = s^2 \sin \theta$ ,  $2r dr = s^2 \cos \theta d\theta$  and  $0 \leq \theta \leq \frac{\pi}{2}$ , we obtain

$$\begin{aligned} |\Psi_t^\varepsilon(x)| &\leq \frac{\Omega_p \Omega_q}{2(2\pi)^{n/2}} \int_0^\infty \int_0^{\pi/2} e^{-\varepsilon c(s^2 \cos \theta)^k} (\sin \theta)^{\frac{p-2}{2}} s^{p+q-1} \cos \theta d\theta ds \\ &= \frac{\Omega_p \Omega_q}{2(2\pi)^{n/2}} \int_0^\infty \int_0^{\pi/2} e^{-\varepsilon c(s^2 \cos \theta)^k} s^{p+q-1} (\sin \theta)^{p-2/2} \cos \theta d\theta ds. \end{aligned}$$

Next, putting  $y = \varepsilon c(s^2 \cos \theta)^k$ ,  $ds = s \frac{dy}{2ky}$ , we have

$$\begin{aligned} |\Psi_t^\varepsilon(x)| &\leq \frac{\Omega_p \Omega_q}{4k(2\pi)^{n/2}} \int_0^{\pi/2} \int_0^\infty \frac{e^{-y}}{y} \left( \frac{y}{c\varepsilon \cos^k \theta} \right)^{n/2k} (\sin \theta)^{\frac{p-2}{2}} \cos \theta dy d\theta \\ &= \frac{\Omega_p \Omega_q}{4k(2\pi)^{n/2}} \int_0^{\pi/2} \int_0^\infty \frac{e^{-y} y^{n/2k-1}}{c^{n/2k} \varepsilon^{n/2k}} (\sin \theta)^{\frac{p-2}{2}} (\cos \theta)^{\frac{2-n}{2}} dy d\theta \\ &= \frac{\Omega_p \Omega_q}{4(2\pi)^{n/2} k c^{n/2k} \varepsilon^{n/2k}} \Gamma\left(\frac{n}{2k}\right) \int_0^{\pi/2} (\sin \theta)^{\frac{p-2}{2}} (\cos \theta)^{\frac{2-n}{2}} d\theta, \\ |\Psi_t^\varepsilon(x)| &\leq \frac{\Omega_p \Omega_q}{8(2\pi)^{n/2} k c^{n/2k} \varepsilon^{n/2k}} \frac{\Gamma\left(\frac{n}{2k}\right) \Gamma\left(\frac{p}{4}\right) \Gamma\left(\frac{4-n}{4}\right)}{\Gamma\left(\frac{4-q}{4}\right)}. \end{aligned}$$

Set

$$u^\varepsilon(x, t) = f(x) * \Psi_t^\varepsilon(x) + g(x) * \Phi_t^\varepsilon(x) \quad (3.7)$$

which is an  $\varepsilon$ -approximation of  $u(x, t)$  in (3.7). For  $\varepsilon \rightarrow 0$ ,  $u^\varepsilon(x, t) \rightarrow u(x, t)$  uniformly. Now

$$u^\varepsilon(x, t) = \int_{\mathbb{R}^n} f(r) \Psi_t^\varepsilon(x-r) dr + \int_{\mathbb{R}^n} g(r) \Phi_t^\varepsilon(x-r) dr.$$

Thus

$$\begin{aligned} |u^\varepsilon(x, t)| &\leq |\Psi_t^\varepsilon(x-r)| \int_{\mathbb{R}^n} |f(r)| dr + |\Phi_t^\varepsilon(x-r)| \int_{\mathbb{R}^n} |g(r)| dr \\ &\leq \frac{\Omega_p \Omega_q}{8(2\pi)^{n/2} k c^{n/2k} \varepsilon^{n/2k}} \frac{\Gamma\left(\frac{n}{2k}\right) \Gamma\left(\frac{p}{4}\right) \Gamma\left(\frac{4-n}{4}\right)}{\Gamma\left(\frac{4-q}{4}\right)} M \\ &\quad + \frac{\Omega_p \Omega_q}{8(2\pi)^{n/2} k c^{n/2k} \varepsilon^{n/2k-1}} \frac{\Gamma\left(\frac{n}{2k}-1\right) \Gamma\left(\frac{p}{4}\right) \Gamma\left(\frac{2-n}{4}\right)}{\Gamma\left(\frac{4-q}{4}\right)} N, \\ \varepsilon^{n/2k} |u^\varepsilon(x, t)| &\leq \frac{\Omega_p \Omega_q}{8(2\pi)^{n/2} k c^{n/2k}} \frac{\Gamma\left(\frac{n}{2k}\right) \Gamma\left(\frac{p}{4}\right) \Gamma\left(\frac{4-n}{4}\right)}{\Gamma\left(\frac{4-q}{4}\right)} M \\ &\quad + \frac{\Omega_p \Omega_q \varepsilon}{8(2\pi)^{n/2} k c^{n/2k}} \frac{\Gamma\left(\frac{n}{2k}-1\right) \Gamma\left(\frac{p}{4}\right) \Gamma\left(\frac{4-n}{4}\right)}{\Gamma\left(\frac{4-q}{4}\right)} N, \end{aligned}$$

where  $M = \int_{\mathbb{R}^n} |f(r)| dr$  and  $N = \int_{\mathbb{R}^n} |g(r)| dr$ . Since  $f$  and  $g$  are absolutely integrable,

$$\lim_{\varepsilon \rightarrow 0} \varepsilon^{n/2k} |u^\varepsilon(x, t)| \leq \frac{\Omega_p \Omega_q}{8(2\pi)^{n/2} k c^{n/2k}} \frac{\Gamma\left(\frac{n}{2k}\right) \Gamma\left(\frac{p}{4}\right) \Gamma\left(\frac{4-n}{4}\right)}{\Gamma\left(\frac{4-q}{4}\right)} = K.$$

It follows that  $u(x, t) = O(\varepsilon^{-n/2k})$  for  $n \neq k$  as  $\varepsilon \rightarrow 0$ .

## DIAMOND OPERATOR RELATED TO BIHARMONIC EQUATION

In particular, if we put  $k = 2$ ,  $n = 1$  and  $p = 0$ , then (3.1) reduces to the solution of the beam equation, see [5, p. 47],

$$u(x, 0) = f(x) \quad \text{and} \quad \frac{\partial}{\partial t} u(x, 0) = g(x),$$

where  $f$  and  $g$  are continuous and absolutely integrable for  $x \in \mathbb{R}^n$ .

Thus we obtain  $u(x, t) = O(\varepsilon^{-1/4})$  which is a solution of such a biharmonic wave equation.

### Acknowledgement

The author would like to thank The Thailand Research Fund and Graduate School, Chiang Mai University, Thailand for financial support.

### References

- [1] G. B. Folland, Introduction to Partial Differential Equation, Princeton University Press, Princeton, New Jersey, 1995.
- [2] A. Kananthai, On the solution of  $n$ -dimensional Diamond operator, Appl. Math. Comput. 88 (1997), 27-37.
- [3] A. Kananthai, On the Fourier transform of the Diamond kernel of Marcel Riesz, Appl. Math. Comput. 101 (1999), 151-158.
- [4] A. Kanathai and K. Nonlaopon, On the generalized heat kernel, Computational Technologies 9(1) (2004), 3-10.
- [5] J. David Logan, An Introduction to Nonlinear Partial Differential Equations, A Wiley-Interscience Publicatin, John Wiley & Sons, Inc., 1997.
- [6] W. Satsanit and A. Kananthai, On the ultra-hyperbolic wave operator, Int. J. Pure Appl. Math., reprint.
- [7] G. Sritantana and A. Kananthai, On the gernalized wave equation related to the beam equation, Journal of Mathematics Analysis and Approximation Theory 1(1) (2006), 23-29.

# The Operator $\otimes$ and Its Spectrum Related to Heat Equation

Wanchak Satsanit and Amnuay Kananthai

Department of Mathematics

Chiangmai University

Chiangmai, 50200, Thailand

E-mail: malamnka@science.cmu.ac.th

## Abstract

In this paper, we study the equation

$$\frac{\partial}{\partial t} u(x, t) + c^2 \otimes u(x, t) = 0$$

with the initial condition

$$u(x, 0) = f(x)$$

for  $x \in \mathbb{R}^n$ -the  $n$ -dimensional Euclidean space. The operator

$$\begin{aligned} \otimes &= \left( \sum_{i=1}^p \frac{\partial^2}{\partial x_i^2} \right)^3 - \left( \sum_{j=p+1}^{p+q} \frac{\partial^2}{\partial x_j^2} \right)^3 \\ &= \left( \sum_{i=1}^p \frac{\partial^2}{\partial x_i^2} - \sum_{j=p+1}^{p+q} \frac{\partial^2}{\partial x_j^2} \right) \left[ \left( \sum_{i=1}^p \frac{\partial^2}{\partial x_i^2} \right)^2 + \left( \sum_{i=1}^p \frac{\partial^2}{\partial x_i^2} \right) \left( \sum_{j=p+1}^{p+q} \frac{\partial^2}{\partial x_j^2} \right) + \left( \sum_{j=p+1}^{p+q} \frac{\partial^2}{\partial x_j^2} \right) \right] \\ &= \frac{3}{4} \diamond \Delta + \frac{1}{4} \square^3 \end{aligned}$$

where

$$\begin{aligned} \Delta &= \frac{\partial^2}{\partial x_1^2} + \frac{\partial^2}{\partial x_2^2} + \dots + \frac{\partial^2}{\partial x_n^2} \\ \square &= \frac{\partial^2}{\partial x_1^2} + \frac{\partial^2}{\partial x_2^2} + \dots + \frac{\partial^2}{\partial x_p^2} - \frac{\partial^2}{\partial x_{p+1}^2} - \frac{\partial^2}{\partial x_{p+2}^2} - \dots - \frac{\partial^2}{\partial x_{p+q}^2} \\ \diamond &= \left( \frac{\partial^2}{\partial x_1^2} + \frac{\partial^2}{\partial x_2^2} + \dots + \frac{\partial^2}{\partial x_p^2} \right)^2 - \left( \frac{\partial^2}{\partial x_{p+1}^2} + \frac{\partial^2}{\partial x_{p+2}^2} + \dots + \frac{\partial^2}{\partial x_{p+q}^2} \right)^2 \end{aligned}$$

$p + q = n$  is the dimension of the Euclidean space  $\mathbb{R}^n$ ,  $u(x, t)$  is an unknown function for  $(x, t) = (x_1, x_2, \dots, x_n, t) \in \mathbb{R}^n \times (0, \infty)$ ,  $f(x)$  is the given generalized function and  $c$  is a positive constant.

On the suitable conditions for  $f$  and  $u$ , we obtain the uniqueness solution of such equation. Moreover, if we put  $q = 0$  we obtain the solution of heat equation

$$\frac{\partial}{\partial t} u(x, t) + c^2 \Delta^3 u(x, t) = 0.$$

**Key Words:** Fourier transform, Tempered distribution, Diamond operator.

## 1 Introduction

It is well known that for the heat equation

$$\frac{\partial}{\partial t} u(x, t) = c^2 \Delta u(x, t) \quad (1.1)$$

with the initial condition

$$u(x, 0) = f(x)$$

where  $\Delta = \sum_{i=1}^n \frac{\partial^2}{\partial x_i^2}$  is the Laplace operator and  $(x, t) = (x_1, x_2, \dots, x_n, t) \in \mathbb{R}^n \times (0, \infty)$ , we obtain

$$u(x, t) = \frac{1}{(4c^2\pi t)^{n/2}} \int_{\mathbb{R}^n} \exp\left(-\frac{|x-y|^2}{4c^2t}\right) f(y) dy \quad (1.2)$$

as the solution of (1.1).

Now, (1.2) can be written  $u(x, t) = E(x, t) * f(x)$  where

$$E(x, t) = \frac{1}{(4c^2\pi t)^{n/2}} \exp\left(-\frac{|x|^2}{4c^2t}\right). \quad (1.3)$$

$E(x, t)$  is called *the heat kernel*, where  $|x|^2 = x_1^2 + x_2^2 + \dots + x_n^2$  and  $t > 0$ , see [1, p208-209].

In 1996, A. Kananthai [2] has introduced the Diamond operator  $\diamond$  defined by

$$\diamond = \left( \sum_{i=1}^p \frac{\partial^2}{\partial x_i^2} \right)^2 - \left( \sum_{j=p+1}^{p+q} \frac{\partial^2}{\partial x_j^2} \right)^2, \quad p + q = n$$

or  $\diamond$  can be written as the product of the operators in the form  $\diamond = \Delta \square = \square \Delta$  where

$\Delta = \sum_{i=1}^n \frac{\partial^2}{\partial x_i^2}$  is the Laplacian and  $\square = \sum_{i=1}^p \frac{\partial^2}{\partial x_i^2} - \sum_{j=p+1}^{p+q} \frac{\partial^2}{\partial x_j^2}$  is the ultra-hyperbolic. The

Fourier transform of the Diamond operator also has been studied and the elementary solution of such operator, see [3].

Next, K. Nonlaopon and A. Kananthai (see [5]) study the equation

$$\frac{\partial}{\partial t} u(x, t) = c^2 \square u(x, t)$$

Now, the purpose of this work is to study the equation

$$\frac{\partial}{\partial t} u(x, t) + c^2 \otimes u(x, t) = 0 \quad (1.4)$$

with the initial condition

$$u(x, 0) = f(x)$$

for  $x \in \mathbb{R}^n$ -the  $n$ -dimensional Euclidean space. The operator

$$\begin{aligned} \otimes &= \left( \sum_{i=1}^p \frac{\partial^2}{\partial x_i^2} \right)^3 - \left( \sum_{j=p+1}^{p+q} \frac{\partial^2}{\partial x_j^2} \right)^3 \\ &= \left( \sum_{i=1}^p \frac{\partial^2}{\partial x_i^2} - \sum_{j=p+1}^{p+q} \frac{\partial^2}{\partial x_j^2} \right) \left[ \left( \sum_{i=1}^p \frac{\partial^2}{\partial x_i^2} \right)^2 + \left( \sum_{i=1}^p \frac{\partial^2}{\partial x_i^2} \right) \left( \sum_{j=p+1}^{p+q} \frac{\partial^2}{\partial x_j^2} \right) + \left( \sum_{j=p+1}^{p+q} \frac{\partial^2}{\partial x_j^2} \right)^2 \right] \\ &= \square (\Delta^2 - \frac{1}{4} (\Delta + \square) (\Delta - \square)) \\ &= \frac{3}{4} \diamond \Delta + \frac{1}{4} \square^3 \end{aligned}$$

$p + q = n$  is the dimension of the Euclidean space  $\mathbb{R}^n$ ,  $u(x, t)$  is an unknown function for  $(x, t) = (x_1, x_2, \dots, x_n, t) \in \mathbb{R}^n \times (0, \infty)$ ,  $f(x)$  is the given generalized function and  $c$  is a positive constant. We obtain  $u(x, t) = E(x, t) * f(x)$  as a solution of (1.4), where

$$E(x, t) = \frac{1}{(2\pi)^n} \int_{\Omega} \exp \left[ c^2 \left[ \left( \sum_{i=1}^p \xi_i^2 \right)^3 - \left( \sum_{j=p+1}^{p+q} \xi_j^2 \right)^3 \right] t + i(\xi, x) \right] d\xi. \quad (1.5)$$

and  $\Omega \subset \mathbb{R}^n$  is the spectrum of  $E(x, t)$  for any fixed  $t > 0$ . The function  $E(x, t)$  is the elementary solution of (1.5).

All properties of  $E(x, t)$  will be studied in details.

Now, if we put  $q = 0$  in (1.4), then (1.4) reduces to the equation

$$\frac{\partial}{\partial t} u(x, t) + c^2 \Delta^3 u(x, t) = 0$$

which is related to the heat equation.

## 2 Preliminaries

**Definition 2.1** Let  $f(x) \in L_1(\mathbb{R}^n)$ -the space of integrable function in  $\mathbb{R}^n$ . The Fourier transform of  $f(x)$  is defined by

$$\widehat{f}(\xi) = \frac{1}{(2\pi)^{n/2}} \int_{\mathbb{R}^n} e^{-i(\xi, x)} f(x) dx \quad (2.1)$$

where  $\xi = (\xi_1, \xi_2, \dots, \xi_n)$  and  $x = (x_1, x_2, \dots, x_n) \in \mathbb{R}^n$ ,  $(\xi, x) = \xi_1 x_1 + \xi_2 x_2 + \dots + \xi_n x_n$  and  $dx = dx_1 dx_2 \dots dx_n$ .

Also, the inverse of Fourier transform is defined by

$$f(x) = \frac{1}{(2\pi)^{n/2}} \int_{\mathbb{R}^n} e^{i(\xi, x)} \widehat{f}(\xi) d\xi. \quad (2.2)$$

If  $f$  is a distribution with compact supports by [6], Theorem 7.4-3, p.187 Eq.(2.1) can be written as

$$\widehat{f}(\xi) = \frac{1}{(2\pi)^{n/2}} \langle f(x), e^{-i(\xi, x)} \rangle. \quad (2.3)$$

**Definition 2.2** The spectrum of the kernel  $E(x, t)$  of (1.5) is the bounded support of the Fourier transform  $\widehat{E}(\xi, t)$  for any fixed  $t > 0$ .

**Definition 2.3** Let  $\xi = (\xi_1, \xi_2, \dots, \xi_n)$  be a point in  $\mathbb{R}^n$  and denote by

$$\Gamma_+ = \{\xi \in \mathbb{R}^n : \xi_1^2 + \xi_2^2 + \dots + \xi_p^2 - \xi_{p+1}^2 - \xi_{p+2}^2 - \dots - \xi_{p+q}^2 > 0 \text{ and } \xi_1 > 0\}$$

the set of an interior of the forward cone, and  $\overline{\Gamma}_+$  denotes the closure of  $\Gamma_+$ .

Let  $\Omega$  be spectrum of  $E(x, t)$  defined by definition 2.2 for any fixed  $t > 0$  and  $\Omega \subset \overline{\Gamma}_+$ . Let  $\widehat{E}(\xi, t)$  be the Fourier transform of  $E(x, t)$  and define

$$\widehat{E}(\xi, t) = \begin{cases} \frac{1}{(2\pi)^{n/2}} \exp \left[ c^2 \left( \left( \sum_{i=1}^p \xi_i^2 \right)^3 - \left( \sum_{j=p+1}^{p+q} \xi_j^2 \right)^3 \right) t \right] & \text{for } \xi \in \Gamma_+, \\ 0 & \text{for } \xi \notin \Gamma_+. \end{cases} \quad (2.4)$$

**Lemma 2.1** (The Fourier transform of  $\otimes \delta$ )

$$\mathcal{F} \otimes \delta = \frac{(-1)^3}{(2\pi)^{n/2}} \left[ (\xi_1^2 + \xi_2^2 + \dots + \xi_p^2)^3 - (\xi_{p+1}^2 + \xi_{p+2}^2 - \dots + \xi_{p+q}^2)^3 \right]$$

where  $\mathcal{F}$  is the Fourier transform defined by Eq.(2.1) and if the norm of  $\xi$  is given by  $\|\xi\| = (\xi_1^2 + \xi_2^2 + \dots + \xi_n^2)^{1/2}$  then

$$|\mathcal{F} \otimes \delta| \leq \frac{3}{(2\pi)^{n/2}} \|\xi\|^6$$

that is  $\mathcal{F} \otimes$  is bounded and continuous on the space  $S'$  of the tempered distribution. Moreover, by Eq.(2.2)

$$\otimes \delta = \mathcal{F}^{-1} \frac{1}{(2\pi)^{n/2}} \left[ (\xi_1^2 + \xi_2^2 + \dots + \xi_p^2)^3 - (\xi_{p+1}^2 + \xi_{p+2}^2 - \dots + \xi_{p+q}^2)^3 \right]$$

**Proof.** By Eq. (2.3)

$$\begin{aligned} \mathcal{F} \otimes \delta &= \frac{1}{(2\pi)^{n/2}} \langle \otimes \delta, e^{-i(\xi, x)} \rangle \\ &= \frac{1}{(2\pi)^{n/2}} \langle \delta, \otimes e^{-i(\xi, x)} \rangle \\ &= \frac{1}{(2\pi)^{n/2}} \left\langle \delta, \left( \frac{3}{4} \diamond \Delta + \frac{1}{4} \square^3 \right) e^{-i(\xi, x)} \right\rangle \\ &= \frac{1}{(2\pi)^{n/2}} \left\langle \delta, \frac{3}{4} \diamond \Delta e^{-i(\xi, x)} \right\rangle + \frac{1}{(2\pi)^{n/2}} \left\langle \delta, \frac{1}{4} \square^3 e^{-i(\xi, x)} \right\rangle \\ &= \frac{1}{(2\pi)^{n/2}} \left\langle \delta, \frac{3}{4} (-1)^2 \left[ \left( \sum_{i=1}^p \xi_i^2 \right)^2 - \left( \sum_{j=p+1}^{p+q} \xi_j^2 \right)^2 \right] (-1) \left( \sum_{i=1}^n \xi_i^2 \right) e^{-i(\xi, x)} \right\rangle \\ &\quad + \frac{1}{(2\pi)^{n/2}} \left\langle \delta, \frac{1}{4} (-1)^3 \left[ \left( \sum_{i=1}^p \xi_i^2 \right) - \left( \sum_{j=p+1}^{p+q} \xi_j^2 \right) \right]^3 e^{-i(\xi, x)} \right\rangle \\ &= \frac{1}{(2\pi)^{n/2}} \left[ \frac{3}{4} (-1)^3 \left[ \left( \sum_{i=1}^p \xi_i^2 \right)^2 - \left( \sum_{j=p+1}^{p+q} \xi_j^2 \right)^2 \right] \left( \sum_{i=1}^n \xi_i^2 \right) \right] \\ &\quad + \frac{1}{(2\pi)^{n/2}} \left( \frac{1}{4} (-1)^3 \left[ \left( \sum_{i=1}^p \xi_i^2 \right) - \left( \sum_{j=p+1}^{p+q} \xi_j^2 \right) \right]^3 \right) \\ &= \frac{(-1)^3}{(2\pi)^{n/2}} \left[ \left( \sum_{i=1}^p \xi_i^2 \right)^3 - \left( \sum_{j=p+1}^{p+q} \xi_j^2 \right)^3 \right] \\ &= \frac{(-1)^3}{(2\pi)^{n/2}} \left[ (\xi_1^2 + \xi_2^2 + \dots + \xi_p^2)^3 - (\xi_{p+1}^2 + \xi_{p+2}^2 + \dots + \xi_{p+q}^2)^3 \right]. \end{aligned}$$

Now,

$$\begin{aligned}
|\mathcal{F} \otimes \delta| &= \frac{1}{(2\pi)^{n/2}} \left| (\xi_1^2 + \xi_2^2 + \dots + \xi_p^2)^3 - (\xi_{p+1}^2 + \xi_{p+2}^2 + \dots + \xi_{p+q}^2)^3 \right| \\
&\leq \frac{1}{(2\pi)^{n/2}} |\xi_1^2 + \dots + \xi_n^2| \left| (\xi_1^2 + \dots + \xi_n^2)^2 + (\xi_1^2 + \dots + \xi_n^2)^2 + (\xi_1^2 + \dots + \xi_n^2)^2 \right| \\
&\leq \frac{3}{(2\pi)^{n/2}} \|\xi\|^6
\end{aligned}$$

where  $\|\xi\| = (\xi_1^2 + \xi_2^2 + \dots + \xi_n^2)^{1/2}$ ,  $\xi_i (i = 1, 2, \dots, n) \in \mathbb{R}$ . Hence we obtain  $\mathcal{F} \otimes \delta$  is bounded and continuous on the space  $\mathcal{S}'$  of the tempered distribution.

Since  $\mathcal{F}$  is 1-1 transformation from the space  $\mathcal{S}'$  of the tempered distribution to the real space  $\mathbb{R}$ , then by Eq.(2.2)

$$\otimes \delta = \mathcal{F}^{-1} \frac{1}{(2\pi)^{n/2}} \left[ (\xi_1^2 + \xi_2^2 + \dots + \xi_p^2)^3 - (\xi_{p+1}^2 + \xi_{p+2}^2 + \dots + \xi_{p+q}^2)^3 \right].$$

That completes the proof.

**Lemma 2.2** Given the operator

$$L = \frac{\partial}{\partial t} + c^2 \left[ \left( \sum_{i=1}^p \frac{\partial^2}{\partial x_i^2} \right)^3 - \left( \sum_{j=p+1}^{p+q} \frac{\partial^2}{\partial x_j^2} \right)^3 \right] \quad (2.5)$$

where

$$\left( \sum_{i=1}^p \frac{\partial^2}{\partial x_i^2} \right)^3 - \left( \sum_{j=p+1}^{p+q} \frac{\partial^2}{\partial x_j^2} \right)^3 = \frac{3}{4} \diamond \Delta + \frac{1}{4} \square^3,$$

$p+q = n$  is the dimension of  $\mathbb{R}^n$ ,  $(x, t) = (x_1, x_2, \dots, x_n, t) \in \mathbb{R}^n \times (0, \infty)$ , and  $c$  is a positive constant. Then we obtain

$$E(x, t) = \frac{1}{(2\pi)^n} \int_{\Omega} \exp \left[ c^2 \left[ \left( \sum_{i=1}^p \xi_i^2 \right)^3 - \left( \sum_{j=p+1}^{p+q} \xi_j^2 \right)^3 \right] t + i(\xi, x) \right] d\xi. \quad (2.6)$$

as a elementary solution of (2.5), where  $\sum_{j=p+1}^{p+q} \xi_j^2 > \sum_{i=1}^p \xi_i^2$ .

**Proof.** Let

$$LE(x, t) = \delta(x, t),$$

where  $E(x, t)$  is the elementary solution of operator  $L$  and  $\delta$  is the Dirac-delta distribution. Thus

$$\frac{\partial}{\partial t} E(x, t) + c^2 \left[ \left( \sum_{i=1}^p \frac{\partial^2}{\partial x_i^2} \right)^3 - \left( \sum_{j=p+1}^{p+q} \frac{\partial^2}{\partial x_j^2} \right)^3 \right] E(x, t) = \delta(x) \delta(t).$$

Take the Fourier transform defined by (2.1) to both sides of the equation, we obtain

$$\frac{\partial}{\partial t} \widehat{E(\xi, t)} - c^2 \left[ \left( \sum_{i=1}^p \xi_i^2 \right)^3 - \left( \sum_{j=p+1}^{p+q} \xi_j^2 \right)^3 \right] \widehat{E(\xi, t)} = \frac{1}{(2\pi)^{n/2}} \delta(t).$$

Thus

$$\widehat{E(\xi, t)} = \frac{H(t)}{(2\pi)^{n/2}} \exp \left[ c^2 \left( \left( \sum_{i=1}^p \xi_i^2 \right)^3 - \left( \sum_{j=p+1}^{p+q} \xi_j^2 \right)^3 \right) t \right]$$

where  $H(t)$  is the Heaviside function. Since  $H(t) = 1$  for  $t > 0$ . Therefore,

$$\widehat{E(\xi, t)} = \frac{1}{(2\pi)^{n/2}} \exp \left[ c^2 \left( \left( \sum_{i=1}^p \xi_i^2 \right)^3 - \left( \sum_{j=p+1}^{p+q} \xi_j^2 \right)^3 \right) t \right]$$

which has been already defined by (2.4). Thus

$$\begin{aligned} E(x, t) &= \frac{1}{(2\pi)^{n/2}} \int_{\mathbb{R}^n} e^{i(\xi, x)} \widehat{E(\xi, t)} d\xi \\ &= \frac{1}{(2\pi)^{n/2}} \int_{\Omega} e^{i(\xi, x)} \widehat{E(\xi, t)} d\xi \end{aligned}$$

where  $\Omega$  is the spectrum of  $E(x, t)$ . Thus from (2.2)

$$E(x, t) = \frac{1}{(2\pi)^n} \int_{\Omega} \exp \left[ c^2 \left[ \left( \sum_{i=1}^p \xi_i^2 \right)^3 - \left( \sum_{j=p+1}^{p+q} \xi_j^2 \right)^3 \right] t + i(\xi, x) \right] d\xi.$$

### 3 Main Results

**Theorem 3.1** *Given the equation*

$$\frac{\partial}{\partial t} u(x, t) + c^2 \otimes u(x, t) = 0 \quad (3.1)$$

*with the initial condition*

$$u(x, 0) = f(x) \quad (3.2)$$

The operator

$$\begin{aligned}
 \otimes &= \left( \sum_{i=1}^p \frac{\partial^2}{\partial x_i^2} \right)^3 - \left( \sum_{j=p+1}^{p+q} \frac{\partial^2}{\partial x_j^2} \right)^3 \\
 &= \left( \sum_{i=1}^p \frac{\partial^2}{\partial x_i^2} - \sum_{j=p+1}^{p+q} \frac{\partial^2}{\partial x_j^2} \right) \left[ \left( \sum_{i=1}^p \frac{\partial^2}{\partial x_i^2} \right)^2 + \left( \sum_{i=1}^p \frac{\partial^2}{\partial x_i^2} \right) \left( \sum_{j=p+1}^{p+q} \frac{\partial^2}{\partial x_j^2} \right) + \left( \sum_{j=p+1}^{p+q} \frac{\partial^2}{\partial x_j^2} \right)^2 \right] \\
 &= \frac{3}{4} \diamond \Delta + \frac{1}{4} \square^3
 \end{aligned}$$

$p+q = n$  is the dimension of Euclidean space  $\mathbb{R}^n$ ,  $k$  is a positive integer,  $u(x, t)$  is an unknown function for  $(x, t) = (x_1, x_2, \dots, x_n, t) \in \mathbb{R}^n \times (0, \infty)$ ,  $f(x)$  is the given generalized function, and  $c$  is a positive constant. Then we obtain

$$u(x, t) = E(x, t) * f(x)$$

as a solution of (3.1) which satisfies (3.2) where  $E(x, t)$  is given by (2.6).

**Proof.** Taking the Fourier transform defined by (2.1) to both sides of (3.1), we obtain

$$\frac{\partial}{\partial t} \hat{u}(\xi, t) - c^2 \left[ \left( \sum_{i=1}^p \xi_i^2 \right)^3 - \left( \sum_{j=p+1}^{p+q} \xi_j^2 \right)^3 \right] \hat{u}(\xi, t) = 0,$$

(see Lemma 2.1). Thus

$$\hat{u}(\xi, t) = K(\xi) \exp \left[ c^2 \left( \left( \sum_{i=1}^p \xi_i^2 \right)^3 - \left( \sum_{j=p+1}^{p+q} \xi_j^2 \right)^3 \right) t \right] \quad (3.3)$$

where  $K(\xi)$  is constant and  $\hat{u}(\xi, 0) = K(\xi)$ .

Now, by (3.2) we have

$$K(\xi) = \hat{u}(\xi, 0) = \hat{f}(\xi) = \frac{1}{(2\pi)^{n/2}} \int_{\mathbb{R}^n} e^{-i(\xi, x)} f(x) dx \quad (3.4)$$

and by the inversion in (2.2), (3.3) and (3.4) we obtain

$$\begin{aligned}
 u(x, t) &= \frac{1}{(2\pi)^{n/2}} \int_{\mathbb{R}^n} e^{i(\xi, x)} \hat{u}(\xi, t) d\xi \\
 &= \frac{1}{(2\pi)^n} \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} e^{i(\xi, x)} e^{-i(\xi, y)} f(y) \exp \left[ c^2 \left( \left( \sum_{i=1}^p \xi_i^2 \right)^3 - \left( \sum_{j=p+1}^{p+q} \xi_j^2 \right)^3 \right) t \right] dy d\xi.
 \end{aligned}$$

Thus

$$u(x, t) = \frac{1}{(2\pi)^n} \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} e^{i(\xi, x-y)} \exp \left[ c^2 \left( \left( \sum_{i=1}^p \xi_i^2 \right)^3 - \left( \sum_{j=p+1}^{p+q} \xi_j^2 \right)^3 \right) t \right] f(y) dy d\xi$$

or

$$u(x, t) = \frac{1}{(2\pi)^n} \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \exp \left[ c^2 \left[ \left( \sum_{i=1}^p \xi_i^2 \right)^3 - \left( \sum_{j=p+1}^{p+q} \xi_j^2 \right)^3 \right] t + i(\xi, x-y) \right] f(y) dy d\xi \quad (3.5)$$

Set

$$E(x, t) = \frac{1}{(2\pi)^n} \int_{\mathbb{R}^n} \exp \left[ c^2 \left[ \left( \sum_{i=1}^p \xi_i^2 \right)^3 - \left( \sum_{j=p+1}^{p+q} \xi_j^2 \right)^3 \right] t + i(\xi, x) \right] d\xi. \quad (3.6)$$

We choose  $\Omega \subset \mathbb{R}^n$  be the spectrum of  $E(x, t)$  and by (2.6), we have

$$\begin{aligned} E(x, t) &= \frac{1}{(2\pi)^n} \int_{\mathbb{R}^n} \exp \left[ c^2 \left[ \left( \sum_{i=1}^p \xi_i^2 \right)^3 - \left( \sum_{j=p+1}^{p+q} \xi_j^2 \right)^3 \right] t + i(\xi, x) \right] d\xi \\ &= \frac{1}{(2\pi)^n} \int_{\Omega} \exp \left[ c^2 \left[ \left( \sum_{i=1}^p \xi_i^2 \right)^3 - \left( \sum_{j=p+1}^{p+q} \xi_j^2 \right)^3 \right] t + i(\xi, x) \right] d\xi. \end{aligned} \quad (3.7)$$

Thus (3.5) can be written in the convolution form

$$u(x, t) = E(x, t) * f(x).$$

Since  $E(x, t)$  exists, then

$$\begin{aligned} \lim_{t \rightarrow 0} E(x, t) &= \frac{1}{(2\pi)^n} \int_{\Omega} e^{i(\xi, x)} d\xi \\ &= \frac{1}{(2\pi)^n} \int_{\mathbb{R}^n} e^{i(\xi, x)} d\xi \\ &= \delta(x), \quad \text{for } x \in \mathbb{R}^n. \end{aligned} \quad (3.8)$$

See [4, p396, Eq.(10.2.19b)].

Thus for the solution  $u(x, t) = E(x, t) * f(x)$  of (3.1), then

$$\lim_{t \rightarrow 0} u(x, t) = u(x, 0) = \delta * f(x) = f(x)$$

which satisfies (3.2).

**Theorem 3.2** The kernel  $E(x, t)$  defined by (3.7) has the following properties :

(1)  $E(x, t) \in C^\infty$ -the space of continuous function for  $x \in \mathbb{R}^n$ ,  $t > 0$  with infinitely differentiable.

$$(2) \left( \frac{\partial}{\partial t} + c^2 \left[ \left( \sum_{i=1}^p \frac{\partial^2}{\partial x_i^2} \right)^3 - \left( \sum_{j=p+1}^{p+q} \frac{\partial^2}{\partial x_j^2} \right)^3 \right] \right) E(x, t) = 0 \quad \text{for } t > 0.$$

(3)  $E(x, t) > 0$  for  $t > 0$ .

$$(4) |E(x, t)| \leq \frac{2^{2-n}}{\pi^{n/2}} \frac{M(t)}{\Gamma(\frac{p}{2})\Gamma(\frac{q}{2})}, \quad \text{for } t > 0,$$

where  $M(t)$  is a function of  $t$  in the spectrum  $\Omega$  and  $\Gamma$  denote the Gamma function. Thus  $E(x, t)$  is bounded for any fixed  $t > 0$ .

$$(5) \lim_{t \rightarrow 0} E(x, t) = \delta.$$

**Proof.**

(1) From (3.7), since

$$\frac{\partial^n}{\partial x^n} E(x, t) = \frac{1}{(2\pi)^n} \int_{\Omega} \frac{\partial^n}{\partial x^n} \exp \left[ c^2 \left[ \left( \sum_{i=1}^p \xi_i^2 \right)^3 - \left( \sum_{j=p+1}^{p+q} \xi_j^2 \right)^3 \right] t + i(\xi, x) \right] d\xi.$$

Thus  $E(x, t) \in C^\infty$  for  $x \in \mathbb{R}^n$ ,  $t > 0$ .

(2) By computing directly, we obtain

$$\left( \frac{\partial}{\partial t} + c^2 \left[ \left( \sum_{i=1}^p \xi_i^2 \right)^3 - \left( \sum_{j=p+1}^{p+q} \xi_j^2 \right)^3 \right] \right) E(x, t) = 0.$$

(3)  $E(x, t) > 0$  for  $t > 0$  is obvious by (3.7).

(4) We have

$$E(x, t) = \frac{1}{(2\pi)^n} \int_{\Omega} \exp \left[ c^2 \left[ \left( \sum_{i=1}^p \xi_i^2 \right)^3 - \left( \sum_{j=p+1}^{p+q} \xi_j^2 \right)^3 \right] t + i(\xi, x) \right] d\xi.$$

$$|E(x, t)| \leq \frac{1}{(2\pi)^n} \int_{\Omega} \exp \left[ c^2 \left[ \left( \sum_{i=1}^p \xi_i^2 \right)^3 - \left( \sum_{j=p+1}^{p+q} \xi_j^2 \right)^3 \right] t \right] d\xi.$$

By changing to bipolar coordinates

$$\xi_1 = r\omega_1, \xi_2 = r\omega_2, \dots, \xi_p = r\omega_p \quad \text{and}$$

$$\xi_{p+1} = s\omega_{p+1}, \xi_{p+2} = s\omega_{p+2}, \dots, \xi_{p+q} = s\omega_{p+q}$$

where  $\sum_{i=1}^p \omega_i^2 = 1$  and  $\sum_{j=p+1}^{p+q} \omega_j^2 = 1$ . Thus

$$|E(x, t)| \leq \frac{1}{(2\pi)^n} \int_{\Omega} \exp [c^2 (s^6 - r^6) t] r^{p-1} s^{q-1} dr ds d\Omega_p d\Omega_q$$

where  $d\xi = r^{p-1} s^{q-1} dr ds d\Omega_p d\Omega_q$ ,  $d\Omega_p$  and  $\Omega_q$  are the elements of surface area of the unit sphere in  $\mathbb{R}^p$  and  $\mathbb{R}^q$  respectively. Since  $\Omega \subset \mathbb{R}^n$  is the spectrum of  $E(x, t)$  and we suppose  $0 \leq r \leq R$  and  $0 \leq s \leq T$  where  $R$  and  $T$  are constants. Thus we obtain

$$\begin{aligned} |E(x, t)| &\leq \frac{\Omega_p \Omega_q}{(2\pi)^n} \int_0^R \int_0^T \exp [c^2 (s^6 - r^6) t] r^{p-1} s^{q-1} ds dr \\ &= \frac{\Omega_p \Omega_q}{(2\pi)^n} M(t) \quad \text{for any fixed } t > 0 \text{ in the spectrum } \Omega \\ &= \frac{2^{2-n}}{\pi^{n/2}} \frac{M(t)}{\Gamma(\frac{p}{2})\Gamma(\frac{q}{2})} \end{aligned}$$

where

$$M(t) = \int_0^R \int_0^T \exp [c^2 (s^6 - r^6) t] r^{p-1} s^{q-1} ds dr$$

is a function of  $t$ ,  $\Omega_p = \frac{2\pi^{p/2}}{\Gamma(\frac{p}{2})}$  and  $\Omega_q = \frac{2\pi^{q/2}}{\Gamma(\frac{q}{2})}$ . Thus, for any fixed  $t > 0$ ,  $E(x, t)$  is bounded.

(5) Obvious by (3.8).

## Acknowledgement

The authors would like to thank The Thailand Research Fund and Graduate School, Chiang Mai University, Thailand for financial support.

## References

- [1] F. John, "Partial Differential Equations", 4<sup>th</sup> Edition, Springer-Verlag, New York (1982).

- [2] A. Kananthai, *On the Solution of the n-Dimensional Diamond Operator*, Applied Mathematics and Computational 88:27-37(1997).
- [3] A. Kananthai, *On the Fourier Transform of the Diamond Kernel of Marcel Riesz*, Applied Mathematics and Computation 101:151-158(1999)..
- [4] R. Haberman, "Elementary Applied Partial Differential Equations", 2<sup>nd</sup> Edition, Prentice-Hall International, Inc. (1983).
- [5] K. Nonlaopon, A. Kananthai, *On the Ultra-hyperbolic heat kernel*, Applied Mathematics Vol.13 No.2 2003,215-225.
- [6] A. H. Zemanian, *Distribution Theory and Transform Analysis*, McGraw-Hill, New York, 1965.

# On the Diamond-wave operator

Amnuay Kananthai

Department of Mathematics

Chiangmai University

Chiangmai, 50200, Thailand

E-mail: malamnka@science.cmu.ac.th

## Abstract

In this paper, we study the solution of the Diamond-wave operator  $L$  which is defined by

$$L = \frac{\partial^2}{\partial t^2} - \diamond$$

where

$$\diamond = \left( \sum_{i=1}^p \frac{\partial^2}{\partial x_i^2} \right)^2 - \left( \sum_{j=p+1}^{p+q} \frac{\partial^2}{\partial x_j^2} \right)^2$$

is the Diamond operator,  $x \in \mathbb{R}^n$ —the  $n$  dimensional Euclidean space,  $t \geq 0$ , and  $p+q = n$  is the dimension of  $\mathbb{R}^n$ . By considering the equation  $Lu(x, t) = 0$  with the suitable initial conditions. We obtained the unique solution  $u(x, t)$  of such equation. Moreover, we obtained the boundedness of  $u(x, t)$  subject to the suitable initial conditions. In particular, if we put  $n = 1$ ,  $p = 1$  and  $q = 0$  we also obtained the solution of the beam equation.

## 1 Introduction

In 1996, A. Kananthai [1] has introduced the Diamond operator  $\diamond$  defined by

$$\diamond = \left( \sum_{i=1}^p \frac{\partial^2}{\partial x_i^2} \right)^2 - \left( \sum_{j=p+1}^{p+q} \frac{\partial^2}{\partial x_j^2} \right)^2, \quad p+q = n$$

or  $\diamond$  can be written as the product of the operators in the form  $\diamond = \square \Delta = \Delta \square$  where

$$\square = \sum_{i=1}^p \frac{\partial^2}{\partial x_i^2} - \sum_{j=p+1}^{p+q} \frac{\partial^2}{\partial x_j^2}$$

is the ultra-hyperbolic operator and

$$\Delta = \sum_{i=1}^n \frac{\partial^2}{\partial x_i^2}$$

is the Laplacian. The Fourier transform of the Diamond operator also has been studied and obtaining the elementary solution of such operator, see [2]. It is well known that the wave equation

$$\frac{\partial^2}{\partial t^2} u(x, t) = c^2 \Delta u(x, t)$$

has been studied widely, particularly, the interesting properties of the solution  $u(x, t)$ . The motivation of this paper is that the operator  $\Delta$  is replaced by  $\diamond$  which is call the Diamond wave equation

$$\frac{\partial^2}{\partial t^2} u(x, t) = c^2 \diamond u(x, t)$$

and by adding the initial conditions

$$u(x, 0) = f(x)$$

and

$$\frac{\partial}{\partial t} u(x, 0) = g(x)$$

where  $f, g \in L^1(\mathbb{R}^n)$ -the space of Lebesgue integrable function, we obtained the uniqueness and boundedness solution  $u(x, t)$  of such equation. In particular, if we put  $n = 1$ ,  $p = 1$  and  $q = 0$  in the Diamond-wave equation reduces to the solution of the beam equation

$$\frac{\partial^2}{\partial t^2} u(x, t) + \frac{\partial^4}{\partial x^4} u(x, t) = 0$$

which is well known equation.

## 2 The solution of the Diamond-wave operator

Given the Diamond wave operator

$$\frac{\partial^2}{\partial t^2} u(x, t) = c^2 \diamond u(x, t) \quad (2.1)$$

with initial conditions

$$u(x, 0) = f(x) \quad \text{and} \quad \frac{\partial}{\partial t} u(x, 0) = g(x) \quad (2.2)$$

where  $\diamond = \left( \sum_{i=1}^p \frac{\partial^2}{\partial x_i^2} \right)^2 - \left( \sum_{j=p+1}^{p+q} \frac{\partial^2}{\partial x_j^2} \right)^2$ ,  $x \in \mathbb{R}^n$ ,  $t \geq 0$ ,  $p + q = n$  and  $f, g \in L^1(\mathbb{R}^n)$ .

We now solving the solution of (2.1) satisfying (2.2) by the method of following steps.

**Step 1** Taking the Fourier transform to both sides of (2.1) where the Fourier transform is defined by

$$\mathfrak{F}f(x) = \widehat{f(\xi)} = \frac{1}{(2\pi)^{n/2}} \int_{\mathbb{R}^n} e^{-i(\xi, x)} f(x) dx \quad (2.3)$$

where  $f \in L^1(\mathbb{R}^n)$  and  $(\xi, x) = \xi_1 x_1 + \xi_2 x_2 + \dots + \xi_n x_n$ . The inverse Fourier transform also defined by

$$f(x) = \mathfrak{F}^{-1} \widehat{f(\xi)} = \frac{1}{(2\pi)^{n/2}} \int_{\mathbb{R}^n} e^{-i(\xi, x)} \widehat{f(\xi)} d\xi. \quad (2.4)$$

By applying (2.3) to both side of (2.1), we obtain

$$\frac{\partial^2}{\partial t^2} \widehat{u(\xi, t)} = \left( [\xi_1^2 + \xi_2^2 + \dots + \xi_p^2]^2 - [\xi_{p+1}^2 + \xi_{p+2}^2 + \dots + \xi_{p+q}^2]^2 \right) \widehat{u(\xi, t)} \quad (2.5)$$

where  $\xi = (\xi_1, \xi_2, \dots, \xi_n) \in \mathbb{R}^n$ . Now, put  $\xi_1^2 + \xi_2^2 + \dots + \xi_p^2 = r^2$ ,  $\xi_{p+1}^2 + \xi_{p+2}^2 + \dots + \xi_{p+q}^2 = s^2$  and let  $s > r$ . Then (2.5) becomes

$$\frac{\partial^2}{\partial t^2} \widehat{u(\xi, t)} + (s^4 - r^4) \widehat{u(\xi, t)} = 0, \quad (2.6)$$

we have the initial conditions

$$u(x, 0) = f(x) \quad \text{and} \quad \frac{\partial}{\partial t} u(x, 0) = g(x).$$

Thus

$$\widehat{u(\xi, 0)} = \widehat{f(\xi)} \quad \text{and} \quad \frac{\partial}{\partial t} \widehat{u(\xi, 0)} = \widehat{g(\xi)}. \quad (2.7)$$

Now, we are solving the solution of (2.6) satisfies (2.7). Then

$$\widehat{u(\xi, t)} = A(\xi) \cos \sqrt{s^4 - r^4} t + B(\xi) \sin \sqrt{s^4 - r^4} t \quad \text{and}$$

$$\frac{\partial}{\partial t} \widehat{u(\xi, t)} = -\sqrt{s^4 - r^4} A(\xi) \sin \sqrt{s^4 - r^4} t + \sqrt{s^4 - r^4} B(\xi) \cos \sqrt{s^4 - r^4} t$$

By (2.7),  $\widehat{u(\xi, 0)} = A(\xi) = \widehat{f(\xi)}$  and  $\frac{\partial}{\partial t} \widehat{u(\xi, 0)} = \sqrt{s^4 - r^4} B(\xi) = \widehat{g(\xi)}$ . Then  $B(\xi) = \frac{\widehat{g(\xi)}}{\sqrt{s^4 - r^4}}$ . Thus the solution of (2.6) satisfies (2.7) is

$$\widehat{u(\xi, t)} = \widehat{f(\xi)} \cos \sqrt{s^4 - r^4} t + \frac{\widehat{g(\xi)}}{\sqrt{s^4 - r^4}} \sin \sqrt{s^4 - r^4} t, \quad (2.8)$$

or in the convolution form

$$u(x, t) = f(x) * \psi(x, t) + g(x) * \phi(x, t). \quad (2.9)$$

Thus (2.9) is a solution of (2.1) where  $\widehat{\phi(\xi, t)} = \frac{1}{\sqrt{s^4 - r^4}} \sin \sqrt{s^4 - r^4} t$  and  $\widehat{\psi(\xi, t)} = \frac{\partial}{\partial t} \widehat{\phi(\xi, t)} = \cos \sqrt{s^4 - r^4} t$ . Since  $\widehat{\phi(\xi, t)}$  and  $\widehat{\psi(\xi, t)}$  can not be Lebesgue integrable, that is  $\widehat{\phi}, \widehat{\psi} \notin L^1(\mathbb{R}^n)$ . Thus we can not find the inverse  $\phi$  and  $\psi$  directly. Thus we can compute the inverse  $\phi$  and  $\psi$  by using the method of  $\epsilon$ -approximation.

**Step 2** The method of  $\epsilon$ -approximation, see [3, P178]. Now, defined  $\widehat{\phi_\epsilon(\xi, t)} = e^{-\epsilon\sqrt{s^4 - r^4}} \widehat{\phi(\xi, t)}$  and  $\widehat{\psi_\epsilon(\xi, t)} = e^{-\epsilon\sqrt{s^4 - r^4}} \widehat{\psi(\xi, t)}$ . Clearly,  $\widehat{\phi_\epsilon(\xi, t)} \rightarrow \widehat{\phi(\xi, t)}, \widehat{\psi_\epsilon(\xi, t)} \rightarrow \widehat{\psi(\xi, t)}$  uniformly as  $\epsilon \rightarrow 0$ , since  $\widehat{\phi_\epsilon}, \widehat{\psi_\epsilon} \in L^1(\mathbb{R}^n)$ , then we can obtain the inverse  $\phi_\epsilon$  and  $\psi_\epsilon$  by applying (2.3) and we obtain  $\phi_\epsilon \rightarrow \phi$  and  $\psi_\epsilon \rightarrow \psi$  as  $\epsilon \rightarrow 0$ . Now, by (2.3) we have

$$\begin{aligned} \phi_\epsilon(x, t) &= \frac{1}{(2\pi)^{n/2}} \int_{\mathbb{R}^n} e^{-i(\xi, x)} \widehat{\phi_\epsilon(\xi, t)} d\xi \\ &= \frac{1}{(2\pi)^{n/2}} \int_{\mathbb{R}^n} e^{-i(\xi, x)} e^{-\epsilon\sqrt{s^4 - r^4}} \frac{\sin \sqrt{s^4 - r^4} t}{\sqrt{s^4 - r^4}} d\xi \end{aligned}$$

and

$$|\phi_\epsilon(x, t)| \leq \frac{1}{(2\pi)^{n/2}} \int_{\mathbb{R}^n} \frac{e^{-\epsilon\sqrt{s^4 - r^4}}}{\sqrt{s^4 - r^4}} d\xi.$$

Now, put  $\xi_1 = rw_1, \xi_2 = rw_2, \dots, \xi_p = rw_p$  and  $\xi_{p+1} = sw_{p+1}, \xi_{p+2} = sw_{p+2}, \dots, \xi_p = sw_{p+q}$ ,  $p+q = n$  where  $w_1^2 + w_2^2 + \dots + w_p^2 = 1$  and  $w_{p+1}^2 + w_{p+2}^2 + \dots + w_{p+q}^2 = 1$ . Thus, by bipolar coordinate

$$\begin{aligned} |\phi_\epsilon(x, t)| &\leq \frac{1}{(2\pi)^{n/2}} \int_0^\infty \int_0^\infty \int_0^s \frac{e^{-\epsilon\sqrt{s^4 - r^4}}}{\sqrt{s^4 - r^4}} r^{p-1} dr s^{q-1} ds d\Omega_p d\Omega_q \\ &= \frac{\Omega_p \Omega_q}{(2\pi)^{n/2}} \int_0^\infty \int_0^s \frac{e^{-\epsilon\sqrt{s^4 - r^4}}}{\sqrt{s^4 - r^4}} r^{p-1} s^{q-1} dr ds \end{aligned}$$

where  $\Omega_p = \frac{2\pi^{p/2}}{\Gamma(\frac{p}{2})}, \Omega_q = \frac{2\pi^{q/2}}{\Gamma(\frac{q}{2})}$  is the surface area of the unit spheres in  $\mathbb{R}^p$  and  $\mathbb{R}^q$  respectively. Now, put  $r^2 = s^2 \sin \theta$ , thus  $0 \leq \theta \leq \frac{\pi}{2}$  we have  $2rdr = s^2 \cos \theta d\theta$ . Then

$$dr = \frac{s^2 \cos \theta}{2r} d\theta = \frac{s^2 \cos \theta}{2s(\sin \theta)^{1/2}} d\theta.$$

Thus

$$\begin{aligned} |\phi_\epsilon(x, t)| &\leq \frac{\Omega_p \Omega_q}{(2\pi)^{n/2}} \int_0^\infty \int_0^{\pi/2} \frac{e^{-\epsilon s^2 \cos \theta}}{s^2 \cos \theta} \cdot \frac{s^{p-1} (\sin \theta)^{\frac{p-1}{2}} s^2 \cos \theta}{2s(\sin \theta)^{1/2}} d\theta s^{q-1} ds \\ &= \frac{\Omega_p \Omega_q}{2(2\pi)^{n/2}} \int_0^\infty \int_0^{\pi/2} e^{-\epsilon s^2 \cos \theta} (\sin \theta)^{\frac{p-2}{2}} s^{p+q-3} d\theta ds \\ &= \frac{\Omega_p \Omega_q}{2(2\pi)^{n/2}} \int_0^{\pi/2} \int_0^\infty e^{-\epsilon s^2 \cos \theta} s^{p+q-3} ds (\sin \theta)^{\frac{p-2}{2}} d\theta \end{aligned}$$

Now, put  $y = \epsilon s^2 \cos \theta$ , thus  $s^2 = \frac{y}{\epsilon \cos \theta}$ ,  $s = \left( \frac{y}{\epsilon \cos \theta} \right)^{\frac{1}{2}}$ , and so

$$ds = \frac{dy}{2s\epsilon \cos \theta} = \frac{dy}{2\epsilon \cos \theta} \left( \frac{\epsilon \cos \theta}{y} \right)^{\frac{1}{2}}.$$

Thus

$$\begin{aligned} \int_0^\infty e^{-\epsilon s^2 \cos \theta} s^{p+q-3} ds &= \int_0^\infty e^{-y} \left( \frac{y}{\epsilon \cos \theta} \right)^{\frac{p+q-3}{2}} \frac{(\epsilon \cos \theta)^{-\frac{1}{2}}}{2\sqrt{y}} dy \\ &= \frac{1}{2} \cdot \frac{1}{(\epsilon \cos \theta)^{\frac{p+q-2}{2}}} \int_0^\infty e^{-y} y^{\frac{p+q-4}{2}} dy \\ &= \frac{1}{2} \cdot \frac{1}{(\epsilon \cos \theta)^{\frac{n-2}{2}}} \int_0^\infty e^{-y} y^{\frac{n-4}{2}} dy, \quad p+q=n \\ &= \frac{1}{2} \cdot \frac{1}{(\epsilon \cos \theta)^{\frac{n-2}{2}}} \Gamma\left(\frac{n-2}{2}\right), \quad n \neq 2. \end{aligned}$$

Thus

$$\begin{aligned} |\phi_\epsilon(x, t)| &\leq \frac{\Omega_p \Omega_q \Gamma(\frac{n-2}{2})}{4(2\pi)^{n/2} \epsilon^{\frac{n-2}{2}}} \int_0^{\pi/2} (\cos \theta)^{\frac{2-n}{2}} (\sin \theta)^{\frac{p-2}{2}} d\theta \\ &= \frac{\Omega_p \Omega_q \Gamma(\frac{n-2}{2})}{8(2\pi)^{n/2} \epsilon^{\frac{n-2}{2}}} \beta\left(\frac{p}{4}, \frac{4-n}{4}\right) \\ &= \frac{\Omega_p \Omega_q \Gamma(\frac{n-2}{2})}{8(2\pi)^{n/2} \epsilon^{\frac{n-2}{2}}} \cdot \frac{\Gamma(\frac{p}{4}) \Gamma(\frac{4-n}{4})}{\Gamma(\frac{4-q}{4})} \end{aligned} \quad (2.10)$$

Now,  $\widehat{\psi_\epsilon(\xi, t)} = e^{-\epsilon \sqrt{s^4 - r^4}} \widehat{\psi(\xi, t)} = e^{-\epsilon \sqrt{s^4 - r^4}} \cos \sqrt{s^4 - r^4} t$ , thus

$$\psi_\epsilon(\xi, t) = \frac{1}{(2\pi)^{n/2}} \int_{\mathbb{R}^n} e^{-i(\xi, x)} \widehat{\psi_\epsilon(\xi, t)} d\xi = \frac{1}{(2\pi)^{n/2}} \int_{\mathbb{R}^n} e^{-i(\xi, x)} e^{-\epsilon \sqrt{s^4 - r^4}} \cos \sqrt{s^4 - r^4} t d\xi,$$

and

$$|\psi_\epsilon(\xi, t)| \leq \frac{1}{(2\pi)^{n/2}} \int_{\mathbb{R}^n} e^{-\epsilon \sqrt{s^4 - r^4}} d\xi.$$

The same process as computing  $|\phi_\epsilon(\xi, t)|$ , we obtain

$$|\psi_\epsilon(\xi, t)| \leq \frac{\Omega_p \Omega_q \Gamma(\frac{n}{2})}{4(2\pi)^{n/2} \epsilon^{\frac{n}{2}}} \cdot \frac{\Gamma(\frac{p}{4}) \Gamma(\frac{4-n}{4})}{\Gamma(\frac{4-q}{4})} \quad (2.11)$$

Now, from (2.9), we define

$$u_\epsilon(x, t) = f(x) * \psi_\epsilon(x, t) + g(x) * \phi_\epsilon(x, t).$$

Thus  $u_\epsilon(x, t) = \int_{\mathbb{R}^n} \psi_\epsilon(y, t) f(x - y) dy + \int_{\mathbb{R}^n} \phi_\epsilon(y, t) g(x - y) dy$

$$\begin{aligned} |u_\epsilon(x, t)| &\leq \int_{\mathbb{R}^n} |\psi_\epsilon(y, t)| |f(x - y)| dy + \int_{\mathbb{R}^n} |\phi_\epsilon(y, t)| |g(x - y)| dy \\ &\leq \frac{\Omega_p \Omega_q}{4(2\pi)^{n/2}} \cdot \frac{\Gamma(\frac{n}{2}) \Gamma(\frac{p}{4}) \Gamma(\frac{4-n}{4})}{\epsilon^{\frac{n}{2}} \Gamma(\frac{4-q}{4})} \int_{\mathbb{R}^n} |f(x - y)| dy \\ &\quad + \frac{\Omega_p \Omega_q}{8(2\pi)^{n/2}} \cdot \frac{\Gamma(\frac{n-2}{2}) \Gamma(\frac{p}{4}) \Gamma(\frac{4-n}{4})}{\epsilon^{\frac{n-2}{2}} \Gamma(\frac{4-q}{4})} \int_{\mathbb{R}^n} |g(x - y)| dy, \end{aligned}$$

by (2.10) and (2.11). Since  $f, g \in L^1(\mathbb{R})$  and let  $M = \int_{\mathbb{R}^n} |f| dy$  and  $N = \int_{\mathbb{R}^n} |g| dy$  where  $M$  and  $N$  are constant. Thus

$$\begin{aligned} |u_\epsilon(x, t)| &\leq \frac{\Omega_p \Omega_q}{4(2\pi)^{n/2}} \cdot \frac{\Gamma(\frac{n}{2}) \Gamma(\frac{p}{4}) \Gamma(\frac{4-n}{4})}{\epsilon^{\frac{n}{2}} \Gamma(\frac{4-q}{4})} M + \frac{\Omega_p \Omega_q}{8(2\pi)^{n/2}} \cdot \frac{\Gamma(\frac{n-2}{2}) \Gamma(\frac{p}{4}) \Gamma(\frac{4-n}{4})}{\epsilon^{\frac{n-2}{2}} \Gamma(\frac{4-q}{4})} N \\ \epsilon^{\frac{n}{2}} |u_\epsilon(x, t)| &\leq \frac{\Omega_p \Omega_q}{4(2\pi)^{n/2}} \cdot \frac{\Gamma(\frac{n}{2}) \Gamma(\frac{p}{4}) \Gamma(\frac{4-n}{4})}{\Gamma(\frac{4-q}{4})} M + \frac{\epsilon \Omega_p \Omega_q}{8(2\pi)^{n/2}} \cdot \frac{\Gamma(\frac{n-2}{2}) \Gamma(\frac{p}{4}) \Gamma(\frac{4-n}{4})}{\Gamma(\frac{4-q}{4})} N \quad (2.12) \\ \lim_{\epsilon \rightarrow 0} \epsilon^{\frac{n}{2}} |u_\epsilon(x, t)| &\leq \frac{\Omega_p \Omega_q}{4(2\pi)^{n/2}} \cdot \frac{\Gamma(\frac{n}{2}) \Gamma(\frac{p}{4}) \Gamma(\frac{4-n}{4})}{\Gamma(\frac{4-q}{4})} M = K \quad \text{say,} \quad (2.13) \end{aligned}$$

where  $K$  is positive constant. Now  $u_\epsilon(x, t) \rightarrow u(x, t)$  as  $\epsilon \rightarrow 0$ . Thus we obtain  $u(x, t) = \mathcal{O}(\epsilon^{\frac{n}{2}})$  as the solution of (2.1) which is bounded by the  $\epsilon$ -approximation. Now, if we put  $n = 1$ ,  $p = 1$  and  $q = 0$  in (2.1) we obtain the one-dimensional beam equation

$$\frac{\partial^2}{\partial t^2} u(x, t) + \frac{\partial^4}{\partial x^4} u(x, t) = 0$$

which has  $u(x, t) = \mathcal{O}(\epsilon^{\frac{1}{2}})$  as a solution.

### 3 The solution of the Diamond-wave operator in numerical form

We can compute the boundedness of  $\epsilon^{\frac{n}{2}} u_\epsilon(x, t)$  from (2.12) and given some  $\epsilon > 0$  and also given the dimension  $n$  and vary  $p$  and  $q$  from  $p + q = n$ . By setting  $\epsilon \rightarrow 0$ , we obtain the solution  $u(x, t) = \mathcal{O}(\epsilon^{\frac{n}{2}})$  that has been shown by the following table.

p	q	$\varepsilon^2  u_\varepsilon(x, t) $			
		$\varepsilon=0.01$	$\varepsilon=0.001$	$\varepsilon=0.0001$	$\varepsilon=0.00001$
1	49	1.91880383307572	1.91844413229468	1.91840816221658	1.91840456520877
2	48	0	0	0	0
3	47	-8.98733613784666	-8.98565136331550	-8.98548288586239	-8.98546603811708
4	46	-24.00499999999999	-24.00049999999999	-24.00005000000000	-24.00000499999999
5	45	-30.0612600515199	-30.05562473928365	-30.05506120806002	-30.05500485493766
6	44	0	0	0	0
7	43	80.88602524061960	80.87086226983924	80.86934597276120	80.86919434305339
8	42	176.03666666666666	176.00366666666667	176.00036666666667	176.00003666666666
9	41	184.6620260307652	184.6274091127423	184.6239474209401	184.6236012517598
10	40	0	0	0	0
11	39	-368.480781651714	-368.411705895937	-368.4047983203597	-368.4041075628019
12	38	-704.1466666666666	-704.0146666666666	-704.0014666666667	-704.0001466666666
13	37	-654.710819563619	-654.588086854265	-654.5758135833299	-654.5745862562364
14	36	0	0	0	0
15	35	1048.752993931801	1048.556393703820	1048.536733681022	1048.534767678742
16	34	1810.662857142857	1810.323428571428	1810.289485714285	1810.286091428571
17	33	1527.658578981790	1527.372202659965	1527.343565027782	1527.340701264564
18	32	0	0	0	0
19	31	-2035.81463527938	-2035.43299954271	-2035.394835969050	-2035.391019611684
20	30	-3218.95619047619	-3218.35276190476	-3218.292419047619	-3218.286384761904
21	29	-2492.49557623343	-2492.02833065571	-2491.981606097946	-2491.976933642169
22	28	0	0	0	0
23	27	2811.363067766763	2810.836046987553	2810.783344909632	2810.778074701840
24	26	4096.853333333333	4096.085333333333	4096.008533333333	4096.000853333333
25	25	2925.973067752327	2925.424562074140	2925.369711506321	2925.364226449539
26	24	0	0	0	0
27	23	-2811.36306776676	-2810.83604698755	-2810.783344909632	-2810.778074701840
28	22	-3781.71076923076	-3781.00184615384	-3780.930953846153	-3780.923864615384
29	21	-2492.49557623343	-2492.02833065571	-2491.981606097946	-2491.976933642169
30	20	0	0	0	0
31	19	2035.814635279387	2035.432999542718	2035.394835969051	2035.391019611684
32	18	2521.140512820512	2520.667897435896	2520.620635897435	2520.615909743588
33	17	1527.658578981790	1527.372202659965	1527.343565027782	1527.340701264564
34	16	0	0	0	0
35	15	-1048.75299393180	-1048.55639370382	-1048.536733681021	-1048.534767678742
36	14	-1186.41906485671	-1186.19665761689	-1186.174416892911	-1186.172192820513
37	13	-654.710819563618	-654.588086854265	-654.5758135833297	-654.5745862562362
38	12	0	0	0	0
39	11	368.4807816517145	368.4117058959373	368.4047983203595	368.4041075628018
40	10	374.6586520600142	374.5884181948082	374.5813948082876	374.5806924696355
41	9	184.6620260307652	184.6274091127424	184.6239474209401	184.6236012517599
42	8	0	0	0	0
43	7	-80.8860252406196	-80.87086226983924	-80.86934597276120	-80.86919434305339
44	6	-71.3635527733360	-71.35017489424919	-71.34883710634051	-71.34870332754964
45	5	-30.0612600515199	-30.05562473928364	-30.05506120806002	-30.05500485493765
46	4	0	0	0	0
47	3	8.98733613784666	8.98565136331550	8.98548288586239	8.98546603811708
48	2	6.20552632811611	6.20436303428247	6.20424670489910	6.20423507196077
49	1	1.91880383307572	1.91844413229468	1.91840816221658	1.91840456520877

$$n = 50, M = N = 1 \text{ and } p + q = 50$$

From the table, the boundedness of  $\epsilon^{n/2}u_\epsilon(x, t)$  is zero for  $q = 4k(k = 1, 2, \dots, 12)$  because  $\Gamma(\frac{4-4k}{4}) = \pm\infty$  which is the denominator of the inequality (2.12). It follows that  $u(x, t)$  is identical to zero at  $q = 4k$  for  $\epsilon \rightarrow 0$ . Similarly, for  $q = 8k - 1, 8k - 2, 8k - 3(k = 1, 2, \dots, 6)$  we obtain  $\epsilon^{n/2}|u_\epsilon(x, t)|$  is bounded by negative numbers because  $\Gamma(\frac{4-4k}{4})$  is negative at such given  $q$ . It follows that  $\epsilon^{n/2}|u_\epsilon(x, t)|$  is not true for such given  $q$ . Moreover, we obtain the symmetry  $p$  and  $q$  of the same boundedness. For example,  $p = 1, q = 49$  symmetry with  $p = 49, q = 1$ ,  $p = 9, q = 41$  symmetry with  $p = 41, q = 9$  and  $p = 17, q = 33$  symmetry with  $p = 33, q = 17$ . From this table, we see that the boundedness of (2.12) tend to (2.13) as  $\epsilon \rightarrow 0$ . That we obtain  $u(x, t) = O(\epsilon^{-n/2})$ .

## References

- [1] A. Kananthai, *On the Solution of the  $n$ -Dimensional Diamond Operator*, Applied Mathematics and Computation 88:27-37 (1997).
- [2] A. Kanathai, *On the Fourier transform of the Diamond Kernel of Marcel Riesz*, Applied Mathematics and Computation 101:151-158 (1999).
- [3] Gerald B. Folland, *Introduction to Partial Differential Equation*, Princeton University, 1995.

# On the Nonlinear heat equation related to the operator $\oplus^k$

Amnuay Kananthai

Department of Mathematics,  
Chiang Mai University,  
Chiang Mai, 50200 Thailand.  
E-mail : malamnka@science.cmu.ac.th

## Abstract

In this paper, we study the nonlinear equation of the form

$$\frac{\partial}{\partial t} u(x, t) - c^2 \oplus^k u(x, t) = f(x, t, u(x, t))$$

where  $\oplus^k$  is the operator iterated  $k$ -times, defined by

$$\oplus^k = \left[ \left( \sum_{i=1}^p \frac{\partial^2}{\partial x_i^2} \right)^4 - \left( \sum_{j=p+1}^{p+q} \frac{\partial^2}{\partial x_j^2} \right)^4 \right]^k$$

where  $p+q = n$  is the dimension of the Euclidean space  $\mathbb{R}^n$ ,  $u(x, t)$  is an unknown for  $(x, t) = (x_1, x_2, \dots, x_n, t) \in \mathbb{R}^n \times (0, \infty)$ ,  $k$  is a positive integer and  $c$  is a positive constant,  $f$  is the given function in nonlinear form depending on  $x, t$  and  $u(x, t)$ . On suitable conditions for  $f, p, q, k$  and the spectrum, we obtain the unique solution  $u(x, t)$  of such equation.

## 1 Introduction

The operator  $\oplus^k$  can be expressed in the form

$$\oplus^k = \left[ \left( \sum_{i=1}^p \frac{\partial^2}{\partial x_i^2} \right)^2 - \left( \sum_{j=p+1}^{p+q} \frac{\partial^2}{\partial x_j^2} \right)^2 \right]^k \left[ \sum_{i=1}^p \frac{\partial^2}{\partial x_i^2} + i \sum_{j=p+1}^{p+q} \frac{\partial^2}{\partial x_j^2} \right]^k \left[ \sum_{i=1}^p \frac{\partial^2}{\partial x_i^2} - i \sum_{j=p+1}^{p+q} \frac{\partial^2}{\partial x_j^2} \right]^k$$

where  $p + q = n$  is the dimension of  $\mathbb{R}^n$ ,  $i = \sqrt{-1}$  and  $k$  is the positive integer. The operator

$$\left[ \left( \sum_{i=1}^p \frac{\partial^2}{\partial x_i^2} \right)^2 - \left( \sum_{j=p+1}^{p+q} \frac{\partial^2}{\partial x_j^2} \right)^2 \right]^k$$

is called the diamond operator iterated  $k$ -times and denoted by  $\diamond^k$ . The such operator is first introduced by A. Kananthai [1]. Moreover, we can find the elementary solution  $K(x)$  of operator  $\oplus^k$ , that is  $\oplus^k K(x) = \delta$ , where  $\delta$  is the Dirac-delta distribution, see[2, p. 226-228].

In this paper, we study the nonlinear equation

$$\frac{\partial}{\partial t} u(x, t) - c^2 \oplus^k u(x, t) = f(x, t, u(x, t)) \quad (1.1)$$

which is in the form of nonlinear heat equation. We consider the equation (1.1) with the following conditions on  $u$  and  $f$  as follows

- (1)  $u(x, t) \in C^{(8k)}(\mathbb{R}^n)$  for any  $t > 0$  where  $C^{(8k)}(\mathbb{R}^n)$  is the space of continuous function with  $8k$ -derivatives.
- (2)  $f$  satisfies the Lipchitz condition,

$$|f(x, t, u) - f(x, t, w)| \leq A|u - w|$$

where  $A$  is constant with  $0 < A < 1$ .

- (3)  $\int_0^\infty \int_{\mathbb{R}^n} |f(x, t, u(x, t))| dx dt < \infty$  for  $x = (x_1, x_2, \dots, x_n) \in \mathbb{R}^n$ ,  $0 < t < \infty$  and  $u(x, t)$  is continuous function on  $\mathbb{R}^n \times (0, \infty)$ .

Under such conditions of  $f$  and  $u$  and for the spectrum of  $E(x, t)$ , we obtain the convolution

$$u(x, t) = E(x, t) * f(x, t, u(x, t))$$

as a unique solution of (1.1) where  $E(x, t)$  is an elementary solution of (1.1).

## 2 Preliminaries

**Definition 2.1** Let  $f(x) \in L_1(\mathbb{R}^n)$  - the space of integrable function in  $\mathbb{R}^n$ . The Fourier transform of  $f(x)$  is defined by

$$\widehat{f(\xi)} = \frac{1}{(2\pi)^{n/2}} \int_{\mathbb{R}^n} e^{-i(\xi, x)} f(x) dx \quad (2.1)$$

where  $\xi = (\xi_1, \xi_2, \dots, \xi_n)$ ,  $x = (x_1, x_2, \dots, x_n) \in \mathbb{R}^n$ ,  $(\xi, x) = \xi_1 x_1 + \xi_2 x_2 + \dots + \xi_n x_n$  is the usual inner product in  $\mathbb{R}^n$  and  $dx = dx_1 dx_2 \dots dx_n$ . Also, the inverse of Fourier transform is defined by

$$f(x) = \frac{1}{(2\pi)^{n/2}} \int_{\mathbb{R}^n} e^{i(\xi, x)} \widehat{f(\xi)} d\xi. \quad (2.2)$$

**Definition 2.2** The spectrum of the kernel  $E(x, t)$  defined by (2.5) is the bounded support of the Fourier transform  $\widehat{E(\xi, t)}$  for any fixed  $t > 0$ .

**Definition 2.3** Let  $\xi = (\xi_1, \xi_2, \dots, \xi_n)$  be a point in  $\mathbb{R}^n$  and write

$$u = \xi_1^2 + \xi_2^2 + \dots + \xi_p^2 - \xi_{p+1}^2 - \xi_{p+2}^2 - \dots - \xi_{p+q}^2, \quad p + q = n.$$

Denote by  $\Gamma_+ = \{\xi \in \mathbb{R}^n : \xi_1 > 0 \text{ and } u > 0\}$  the set of an interior of the forward cone and denote by  $\overline{\Gamma}_+$  the closure of  $\Gamma_+$ . Let  $\Omega$  be the spectrum of  $E(x, t)$  for any fixed  $t > 0$  and  $\Omega \subset \overline{\Gamma}_+$ . Let  $\widehat{E(\xi, t)}$  be the Fourier transform of  $E(x, t)$  and define

$$\widehat{E(\xi, t)} = \begin{cases} \frac{1}{(2\pi)^{n/2}} \exp \left[ c^2 t \left( \left( \sum_{i=1}^p \xi_i^2 \right)^4 - \left( \sum_{j=p+1}^{p+q} \xi_j^2 \right)^4 \right)^k \right], & \text{for } \xi \in \Gamma_+ \\ 0, & \text{for } \xi \notin \Gamma_+ \end{cases} \quad (2.3)$$

**Lemma 2.1** Let  $L$  be the operator defined by

$$L = \frac{\partial}{\partial t} - c^2 \oplus^k \quad (2.4)$$

where  $\oplus^k$  is the operator iterated  $k$ -times defined by

$$\oplus^k = \left[ \left( \sum_{i=1}^p \frac{\partial^2}{\partial x_i^2} \right)^4 - \left( \sum_{j=p+1}^{p+q} \frac{\partial^2}{\partial x_j^2} \right)^4 \right]^k,$$

$p + q = n$  is the dimension of  $\mathbb{R}^n$ ,  $(x_1, x_2, \dots, x_n) \in \mathbb{R}^n$ ,  $t \in (0, \infty)$ ,  $k$  is a positive integer and  $c$  is the positive constant. Then we obtain

$$E(x, t) = \frac{1}{(2\pi)^n} \int_{\Omega} \exp \left[ c^2 t \left( \left( \sum_{i=1}^p \xi_i^2 \right)^4 - \left( \sum_{j=p+1}^{p+q} \xi_j^2 \right)^4 \right)^k + i(\xi, x) \right] d\xi \quad (2.5)$$

as the elementary solution of (2.4) in the spectrum  $\Omega \subset \mathbb{R}^n$  for  $t > 0$ .

**Proof.** Let  $LE(x, t) = \delta(x, t)$  where  $E(x, t)$  is the kernel or the elementary solution of the operator  $L$  and  $\delta$  is the Dirac-delta distribution. Thus

$$\frac{\partial}{\partial t} E(x, t) - c^2 \oplus^k E(x, t) = \delta(x) \delta(t)$$

take the Fourier transform defined by (2.1) to both sides of the equation

$$\frac{\partial}{\partial t} \widehat{E(\xi, t)} - c^2 \left[ \left( \sum_{i=1}^p \xi_i^2 \right)^4 - \left( \sum_{j=p+1}^{p+q} \xi_j^2 \right)^4 \right]^k \widehat{E(\xi, t)} = \frac{1}{(2\pi)^{n/2}} \delta(t).$$

Thus

$$\widehat{E(\xi, t)} = \frac{H(t)}{(2\pi)^{n/2}} \exp \left[ c^2 t \left( \left( \sum_{i=1}^p \xi_i^2 \right)^4 - \left( \sum_{j=p+1}^{p+q} \xi_j^2 \right)^4 \right)^k \right]$$

where  $H(t)$  is the Heaviside function. Since  $H(t) = 1$  for  $t > 0$ ,

$$\widehat{E(\xi, t)} = \frac{1}{(2\pi)^{n/2}} \exp \left[ c^2 t \left( \left( \sum_{i=1}^p \xi_i^2 \right)^4 - \left( \sum_{j=p+1}^{p+q} \xi_j^2 \right)^4 \right)^k \right],$$

so we have

$$E(\xi, t) = \frac{1}{(2\pi)^{n/2}} \int_{\mathbb{R}^n} e^{i(\xi, x)} \widehat{E(\xi, t)} d\xi.$$

By (2.3),

$$E(\xi, t) = \frac{1}{(2\pi)^{n/2}} \int_{\Omega} e^{i(\xi, x)} \widehat{E(\xi, t)} d\xi$$

where  $\Omega$  is the spectrum of  $E(x, t)$ . Thus

$$E(x, t) = \frac{1}{(2\pi)^n} \int_{\Omega} \exp \left[ c^2 t \left( \left( \sum_{i=1}^p \xi_i^2 \right)^4 - \left( \sum_{j=p+1}^{p+q} \xi_j^2 \right)^4 \right)^k + i(\xi, x) \right] d\xi.$$

for  $t > 0$ . □

**Definition 2.4** We can extend  $E(x, t)$  to  $\mathbb{R}^n \times \mathbb{R}$  by setting

$$E(x, t) = \begin{cases} \frac{1}{(2\pi)^n} \int_{\Omega} \exp \left[ c^2 t \left( \left( \sum_{i=1}^p \xi_i^2 \right)^4 - \left( \sum_{j=p+1}^{p+q} \xi_j^2 \right)^4 \right)^k \right] d\xi, & \text{for } t > 0 \\ 0, & \text{for } t \leq 0. \end{cases}$$

**Lemma 2.2** (*The properties of  $E(x, t)$* )

The kernel  $E(x, t)$  defined by (2.5) have the following properties

- (1)  $E(x, t) \in C^\infty$  - the space of continuous function for  $x \in \mathbb{R}^n$ ,  $t > 0$  with infinite differentiable.
- (2)  $\left(\frac{\partial}{\partial t} - c^2 \oplus^k\right) E(x, t) = 0$  for  $t > 0$ .
- (3)  $|E(x, t)| \leq \frac{2^{2-n}}{\pi^{n/2}} \frac{M(t)}{\Gamma(p/2)\Gamma(q/2)}$  for  $t > 0$  where  $M(t)$  is a function of  $t$  in spectrum and  $\Gamma$  denote the Gamma function. Thus  $E(x, t)$  is bounded for a fixed  $t > 0$
- (4)  $\lim_{t \rightarrow 0} E(x, t) = \delta$ .

**Proof.** (1) From (2.5)

$$\frac{\partial^n}{\partial x^n} E(x, t) = \frac{1}{(2\pi)^n} \int_{\Omega} \frac{\partial^n}{\partial x^n} \exp \left[ c^2 t \left( \left( \sum_{i=1}^p \xi_i^2 \right)^4 - \left( \sum_{j=p+1}^{p+q} \xi_j^2 \right)^4 \right)^k + i(\xi, x) \right] d\xi$$

Thus  $E(x, t) \in C^\infty$  for  $x \in \mathbb{R}^n$ ,  $t > 0$ .

(2) By computing directly, we obtain

$$\left( \frac{\partial}{\partial t} - c^2 \oplus^k \right) E(x, t) = 0.$$

(3) We have

$$E(x, t) = \frac{1}{(2\pi)^n} \int_{\Omega} \exp \left[ c^2 t \left( \left( \sum_{i=1}^p \xi_i^2 \right)^4 - \left( \sum_{j=p+1}^{p+q} \xi_j^2 \right)^4 \right)^k + i(\xi, x) \right] d\xi.$$

Thus

$$|E(x, t)| \leq \frac{1}{(2\pi)^n} \int_{\Omega} \exp \left[ c^2 t \left( \left( \sum_{i=1}^p \xi_i^2 \right)^4 - \left( \sum_{j=p+1}^{p+q} \xi_j^2 \right)^4 \right)^k \right] d\xi.$$

By changing to bipolar coordinates

$$\xi_1 = rw_1, \xi_2 = rw_2, \dots, \xi_p = rw_p \text{ and } \xi_{p+1} = sw_{p+1}, \xi_{p+2} = sw_{p+2}, \dots, \xi_{p+q} = sw_{p+q}$$

where

$$\sum_{i=1}^p w_i^2 = 1 \quad \text{and} \quad \sum_{j=p+1}^{p+q} w_j^2 = 1$$

Thus

$$|E(x, t)| \leq \frac{1}{(2\pi)^n} \int_{\Omega} \exp \left[ c^2 t (r^8 - s^8)^k \right] r^{p-1} s^{q-1} dr ds d\Omega_p d\Omega_q$$

where  $d\xi = r^{p-1} s^{q-1} dr ds d\Omega_p d\Omega_q$ ,  $d\Omega_p$  and  $d\Omega_q$  are the elements of surface area of the unit sphere in  $\mathbb{R}^p$  and  $\mathbb{R}^q$  respectively. Since  $\Omega \subset \mathbb{R}^n$  is the spectrum of  $E(x, t)$  and suppose  $0 \leq r \leq R$  and  $0 \leq s \leq L$  where  $R$  and  $L$  are constants. Thus we obtain

$$\begin{aligned} |E(x, t)| &\leq \frac{\Omega_p \Omega_q}{(2\pi)^n} \int_0^R \int_0^L \exp \left[ c^2 t (r^8 - s^8)^k \right] r^{p-1} s^{q-1} dr ds \\ &= \frac{\Omega_p \Omega_q}{(2\pi)^n} M(t) \quad \text{for any fixed } t > 0 \text{ in the spectrum } \Omega \\ &= \frac{2^{2-n}}{\pi^{n/2}} \frac{M(t)}{\Gamma(p/2) \Gamma(q/2)} \end{aligned} \quad (2.6)$$

where  $M(t) = \int_0^R \int_0^L \exp \left[ c^2 t (r^8 - s^8)^k \right] r^{p-1} s^{q-1} dr ds$  is a function for  $t > 0$ ,  $\Omega_p = \frac{2\pi^{p/2}}{\Gamma(p/2)}$  and  $\Omega_q = \frac{2\pi^{q/2}}{\Gamma(q/2)}$ . Thus for any fixed  $t > 0$ ,  $E(x, t)$  is bounded.

(4) From (2.5),

$$\lim_{t \rightarrow 0} E(x, t) = \frac{1}{(2\pi)^n} \int_{\Omega} e^{i(\xi, x)} d\xi = \frac{1}{(2\pi)^n} \int_{\mathbb{R}^n} e^{i(\xi, x)} d\xi = \delta(x),$$

for  $x \in \mathbb{R}^n$ , see [2, p. 396, Eq. (10.2.19b)]. □

### 3 Main Results

**Theorem 3.1** *Given the nonlinear equation*

$$\frac{\partial}{\partial t} u(x, t) - c^2 \oplus^k u(x, t) = f(x, t, u(x, t)) \quad (3.1)$$

for  $(x, t) \in \mathbb{R}^n \times (0, \infty)$ ,  $k$  is a positive number and with the following conditions on  $u$  and  $f$  as follows

- (1)  $u(x, t) \in C^{(8k)}(\mathbb{R}^n)$  for any  $t > 0$  where  $C^{(8k)}(\mathbb{R}^n)$  is the space of continuous function with  $8k$ -derivatives.

(2)  $f$  satisfies the Lipchitz condition,

$$|f(x, t, u) - f(x, t, w)| \leq A|u - w|$$

where  $A$  is constant with  $0 < A < 1$ .

(3)  $\int_0^\infty \int_{\mathbb{R}^n} |f(x, t, u(x, t))| dx dt < \infty$  for  $x = (x_1, x_2, \dots, x_n) \in \mathbb{R}^n$ ,  $0 < t < \infty$  and  $u(x, t)$  is continuous function on  $\mathbb{R}^n \times (0, \infty)$ .

Then obtain the convolution

$$u(x, t) = E(x, t) * f(x, t, u(x, t)) \quad (3.2)$$

as a unique solution of (3.1) for  $x \in \Omega$  where  $\Omega$  is a compact subset of  $\mathbb{R}^n$  and  $0 \leq t \leq T$  with  $T$  is constant and  $E(x, t)$  is an elementary solution defined by (2.5) and also  $u(x, t)$  is bounded for any fixed  $t > 0$ . In particular, if we put  $k = 1$  and  $p = 0$  in (3.1), then (3.1) reduces to the nonlinear equation

$$\frac{\partial}{\partial t} u(x, t) + c^2 \Delta^4 u(x, t) = f(x, t, u(x, t))$$

which is relate to the heat equation.

**Proof.** Convolving both sides of (3.1) with  $E(x, t)$ , that is

$$E(x, t) * \left[ \frac{\partial}{\partial t} u(x, t) - c^2 \oplus^k u(x, t) \right] = E(x, t) * f(x, t, u(x, t))$$

or

$$\left[ \frac{\partial}{\partial t} E(x, t) - c^2 \oplus^k E(x, t) \right] * u(x, t) = E(x, t) * f(x, t, u(x, t)),$$

so

$$\delta(x, t) * u(x, t) = E(x, t) * f(x, t, u(x, t)).$$

Thus

$$\begin{aligned} u(x, t) &= E(x, t) * f(x, t, u(x, t)) \\ &= \int_{-\infty}^{\infty} \int_{\mathbb{R}^n} E(r, s) f(x - r, t - s, u(x - r, t - s)) dr ds \end{aligned}$$

where  $E(r, s)$  is given by definition (2.4). We next show that  $u(x, t)$  is bounded on  $\mathbb{R}^n \times (0, \infty)$ . We have

$$\begin{aligned} |u(x, t)| &\leq \int_{-\infty}^{\infty} \int_{\mathbb{R}^n} |E(r, s)| |f(x - r, t - s, u(x - r, t - s))| dr ds \\ &\leq \frac{2^{2-n} N M(t)}{\pi^{n/2} \Gamma(p/2) \Gamma(q/2)} \quad \text{by condition (3) and (2.6)} \end{aligned}$$

where  $N = \int_{-\infty}^{\infty} \int_{\mathbb{R}^n} |f(x-r, t-s, u(x-r, t-s))| dr ds$ . Thus  $u(x, t)$  is bounded on  $\mathbb{R}^n \times (0, \infty)$ . To show that  $u(x, t)$  is unique. Suppose there is another solution  $w(x, t)$  of (3.1). Let the operator  $L = \frac{\partial}{\partial t} - c^2 \Delta^k$ , then (3.1) can be written in the form  $Lu(x, t) = f(x, t, u(x, t))$ , thus

$$Lu(x, t) - Lw(x, t) = f(x, t, u(x, t)) - f(x, t, w(x, t)).$$

By the condition (2) of the Theorem

$$|Lu(x, t) - Lw(x, t)| \leq A|u(x, t) - w(x, t)|. \quad (3.3)$$

Let  $\Omega \times (0, T]$  be compact subset of  $\mathbb{R}^n \times (0, \infty)$  and  $L : C^{(8k)}(\Omega) \rightarrow C^{(8k)}(\Omega)$  for  $0 \leq t \leq T$ . Now  $(C^{(8k)}(\Omega), \|\cdot\|)$  is a Banach space where  $u(x, t) \in C^{(8k)}(\Omega)$  for  $0 \leq t \leq T$  and  $\|\cdot\|$  is given by  $\|u(x, t)\| = \sup_{x \in \Omega} |u(x, t)|$ . Then from (3.3) with  $0 < A < 1$ ,

$L$  is contraction mapping on  $C^{(8k)}(\Omega)$ . By contraction theorem, see [3, p. 300], we obtain the operator  $L$  has a fixed point and has uniqueness property. Thus  $u(x, t) = w(x, t)$ . It follows that the solution  $u(x, t)$  of (3.1) is unique for  $(x, t) \in \Omega \times (0, T]$  where  $u(x, t)$  is defined by (3.2). In particular, if we put  $k = 1$  and  $p = 0$  in (3.1), then (3.1) reduces to the nonlinear equation

$$\frac{\partial}{\partial t} u(x, t) + c^2 \Delta^4 u(x, t) = f(x, t, u(x, t))$$

which has solution

$$u(x, t) = E(x, t) * f(x, t, u(x, t))$$

where  $E(x, t)$  is defined by (2.5) with  $k = 1$  and  $p = 0$ . □

### Acknowledgement.

The authors would like to thank The Thailand Research Fund for financial support.

## References

- [1] A. Kananthai, *On the solution of n-dimensional Diamond operator*, Applied Mathematics and computation 88(1997):27-37.
- [2] R. Haberman, *Elementary Applied Partial Differential Equation*, 2<sup>nd</sup> Edition, Prentice-Hall International Inc (1983).
- [3] E. Kreyszig, *Introductory Functional Analysis with Applications*, John Wiley & Sons Inc (1978).

## Chapter III

### Fixed Point Theory in Banach Spaces

In this chapter, we divide into two parts. The first part concerns iterative methods for approximating a fixed point and common fixed points of nonlinear mappings. In this part, we introduce a new three-step iteration with errors for nonexpansive nonself-mappings in a uniformly convex Banach space. Weak and strong convergence theorems of the new three-step iteration under certain control conditions are established. We also modify Noor iterations for non-Lipshitzian mappings in Banach spaces and prove weak and strong convergence theorems of the modified Noor iterations under some control conditions. For finding a common fixed point of a finite family of nonexpansive mappings, we introduce a new iterative method for them and prove weak and strong convergence theorems under some suitable control conditions. The second part of this chapter is to find a common element of a fixed point set of nonlinear mappings and the set of solutions of equilibrium problems. Our results improve and extend many results in this area.

The main results of this chapter are written into 6 papers and all of them are published in international journals. Five of them are concerned with fixed point problem and equilibrium problem, while one of them concerning geometric properties of Banach spaces. Here are the list of all of them.

1. S. Thianwan and S. Suantai, **Convergence** Criteria of a New Three-step Iteration with Errors for Nonexpansive- Nonself- Mappings, Computers and Mathematics with Applications 52 (2006) 1107 – 1118.
2. K. Nammanee and S. Suantai, The Modified Noor Iterations with Errors for Non-Lipschitzian Mappings in Banach Spaces, Applied Mathematics and Computation 187 (2007), 669 – 679.
3. N. Petrot and S. Suantai, The Criteria of Strict Monotonicity and Rotundity points in generalized Calderon-Lozanovski Spaces, Nonlinear Analysis ,2009

4. Kiangtunyakarn and S. Suantai, A new mapping for finding common solutions of equilibrium problems and fixed point problems of finite family of nonexpansive mappings, *Nonlinear Analysis : Theory and Methods*
5. and iterative scheme for generalized equilibrium problems and fixed point problems of finite family of nonexpansive mappings, *Nonlinear Analysis : Hybrid Method* , 2009.
6. Kiangtunyakarn and S. Suantai, A new iterative method for common fixed points of a finite family of nonexpansive mappings, *International Journal of Mathematical and Mathematical Sciences*, Vol. 2009, Article ID 391839, 9 pages doi : 10.1155/2009/391839.



ELSEVIER

Available online at [www.sciencedirect.com](http://www.sciencedirect.com)

SCIENCE @ DIRECT®

Computers and Mathematics with Applications 52 (2006) 1107–1118

An International Journal  
**computers &  
mathematics**  
with applications

[www.elsevier.com/locate/camwa](http://www.elsevier.com/locate/camwa)

# Convergence Criteria of a New Three-Step Iteration with Errors for Nonexpansive Nonself-Mappings

SORNSAK THIANWAN<sup>†</sup> AND SUTHEP SUANTAI<sup>‡</sup>

Department of Mathematics  
Faculty of Science, Chiang Mai University  
Chiang Mai, 50200, Thailand  
[sornsakt@nu.ac.th](mailto:sornsakt@nu.ac.th)  
[scmti005@chiangmai.ac.th](mailto:scmti005@chiangmai.ac.th)

(Received December 2005; accepted February 2006)

**Abstract**—A new three-step iteration with errors for nonexpansive nonself-mappings in Banach spaces is introduced and studied. Weak and strong convergence theorems of such iterations are established. The results obtained in this paper extend and improve the several recent results in this area. © 2006 Elsevier Ltd. All rights reserved.

**Keywords**—Nonexpansive nonself-mappings, Completely continuous, Uniformly convex, Opial's condition, Condition (A).

## 1. INTRODUCTION

Let  $X$  be a normed space,  $C$  be a nonempty convex subset of  $X$ ,  $P : X \rightarrow C$  be the nonexpansive retraction of  $X$  onto  $C$ , and  $T : C \rightarrow X$  be a given mapping. Then for a given  $x_1 \in C$ , compute the sequence  $\{x_n\}$ ,  $\{y_n\}$ , and  $\{z_n\}$  by the iterative scheme

$$\begin{aligned} z_n &= P(a_n T x_n + (1 - a_n - \gamma_n)x_n + \gamma_n u_n), \\ y_n &= P(b_n T z_n + c_n T x_n + (1 - b_n - c_n - \mu_n)x_n + \mu_n v_n), \\ x_{n+1} &= P(\alpha_n T y_n + \beta_n T z_n + (1 - \alpha_n - \beta_n - \lambda_n)x_n + \lambda_n w_n), \quad n \geq 1, \end{aligned} \quad (1.1)$$

where  $\{a_n\}$ ,  $\{b_n\}$ ,  $\{c_n\}$ ,  $\{\alpha_n\}$ ,  $\{\beta_n\}$ ,  $\{\gamma_n\}$ ,  $\{\mu_n\}$ ,  $\{\lambda_n\}$  are appropriate sequences in  $[0, 1]$  and  $\{u_n\}$ ,  $\{v_n\}$ , and  $\{w_n\}$  are bounded sequences in  $C$ .

<sup>†</sup>Supported by the Royal Golden Jubilee Project Grant No. PHD/0160/2547 and the Graduate School of Chiang Mai University, Thailand.

<sup>‡</sup>Author to whom all correspondence should be addressed.

The authors would like to thank the Thailand Research Fund (RGJ Project) and the Graduate School of Chiang Mai University for the financial support during the preparation of this paper.

0898-1221/06/\$ - see front matter © 2006 Elsevier Ltd. All rights reserved.  
doi:10.1016/j.camwa.2006.02.012

Typeset by *AMS-TE*

If  $a_n = c_n = \beta_n = \gamma_n = \mu_n = \lambda_n \equiv 0$ , then (1.1) reduces to the iteration scheme defined by Shahzad [1]

$$\begin{aligned} y_n &= P(b_n T x_n + (1 - b_n)x_n), \\ x_{n+1} &= P(\alpha_n T y_n + (1 - \alpha_n)x_n), \quad n \geq 1, \end{aligned} \quad (1.2)$$

where  $\{b_n\}$ ,  $\{\alpha_n\}$  are appropriate sequences in  $[0, 1]$ .

If  $T: C \rightarrow C$ , then the iterative scheme (1.1) reduces to the three-step iterations with errors

$$\begin{aligned} z_n &= a_n T x_n + (1 - a_n - \gamma_n)x_n + \gamma_n u_n, \\ y_n &= b_n T z_n + c_n T x_n + (1 - b_n - c_n - \mu_n)x_n + \mu_n v_n, \\ x_{n+1} &= \alpha_n T y_n + \beta_n T z_n + (1 - \alpha_n - \beta_n - \lambda_n)x_n + \lambda_n w_n, \quad n \geq 1, \end{aligned} \quad (1.3)$$

where  $\{a_n\}$ ,  $\{b_n\}$ ,  $\{c_n\}$ ,  $\{\alpha_n\}$ ,  $\{\beta_n\}$ ,  $\{\gamma_n\}$ ,  $\{\mu_n\}$ ,  $\{\lambda_n\}$  are appropriate sequences in  $[0, 1]$  and  $\{u_n\}$ ,  $\{v_n\}$ , and  $\{w_n\}$  are bounded sequences in  $C$ .

If  $a_n = c_n = \beta_n = \gamma_n = \mu_n = \lambda_n \equiv 0$ , then the iterative scheme (1.3) reduces to the Ishikawa iterative scheme

$$\begin{aligned} y_n &= b_n T x_n + (1 - b_n)x_n, \\ x_{n+1} &= \alpha_n T y_n + (1 - \alpha_n)x_n, \quad n \geq 1, \end{aligned} \quad (1.4)$$

where  $\{b_n\}$ ,  $\{\alpha_n\}$  are appropriate sequences in  $[0, 1]$ .

Fixed-point iteration processes for approximating the fixed point of nonexpansive mapping in Banach spaces have been studied by various authors, using the Mann iteration process (see [2]) or the Ishikawa iteration process (see [3–6]). In 2000, Noor [7] introduced a three-step iterative scheme and studied the approximate solutions of variational inclusion in Hilbert spaces. In 1998, Jung and Kim [8] proved the existence of a fixed point for nonexpansive nonself-mapping in a uniformly convex Banach space with a uniformly Gâteaux differentiable norm. In [5], Tan and Xu introduced a modified Ishikawa process to approximate fixed points of nonexpansive self-mappings defined on nonempty closed convex bounded subsets of a uniformly convex Banach space. In [9], Zhou *et al.* gave criteria for weak convergence theorems of the Ishikawa iterative scheme (1.4) for nonexpansive self-mapping in a uniformly convex Banach space which satisfies Opial's condition, and for strong convergence theorems for nonexpansive self-mapping in a uniformly convex Banach space which satisfies Condition (A). In 2004, Cho, Zhou and Guo [10] defined and studied a new three-step iteration with errors for asymptotically nonexpansive mappings in a uniformly convex Banach space. Suantai [11] defined a new three-step iteration which is an extension of Noor iterations and gave some weak and strong convergence theorems of such iterations for asymptotically nonexpansive mappings in a uniformly convex Banach space. Recently, Shahzad [1] extended Tan and Xu results [5, Theorem 1, p. 305] to the case of nonexpansive nonself-mapping in a uniformly convex Banach space. Inspired and motivated by research going on in this area, we define and study a new three-step iteration with errors for nonexpansive nonself-mapping. This scheme can be viewed as an extension for the two-step iterative schemes of Shahzad [1].

The purpose of this paper is to establish weak and strong convergence results of the iterative scheme (1.1) for nonexpansive nonself-mappings in a uniformly convex Banach space. Our results extend and improve the corresponding ones announced by Shahzad [1], Tan and Xu [5], and others.

Now, we recall the well-known concepts and results.

Recall that a Banach space  $X$  is said to satisfy *Opial's condition* [12] if  $x_n \rightarrow x$  weakly as  $n \rightarrow \infty$  and  $x \neq y$  imply that

$$\limsup_{n \rightarrow \infty} \|x_n - x\| < \limsup_{n \rightarrow \infty} \|x_n - y\|.$$

In the sequel, the following lemmas are needed to prove our main results.

LEMMA 1.1. (See [5, Lemma 1].) Let  $\{a_n\}$ ,  $\{b_n\}$ , and  $\{\delta_n\}$  be sequences of nonnegative real numbers satisfying the inequality

$$a_{n+1} \leq (1 + \delta_n)a_n + b_n, \quad \forall n = 1, 2, \dots$$

If  $\sum_{n=1}^{\infty} \delta_n < \infty$  and  $\sum_{n=1}^{\infty} b_n < \infty$ , then

- (1)  $\lim_{n \rightarrow \infty} a_n$  exists.
- (2)  $\lim_{n \rightarrow \infty} a_n = 0$  whenever  $\liminf_{n \rightarrow \infty} a_n = 0$ .

LEMMA 1.2. (See [13, Lemma 1.4].) Let  $X$  be a uniformly convex Banach space and  $B_r = \{x \in X : \|x\| \leq r\}$ ,  $r > 0$ . Then there exists a continuous, strictly increasing, and convex function  $g : [0, \infty) \rightarrow [0, \infty)$ ,  $g(0) = 0$  such that

$$\|\alpha x + \beta y + \mu z + \lambda w\|^2 \leq \alpha \|x\|^2 + \beta \|y\|^2 + \mu \|z\|^2 + \lambda \|w\|^2 - \alpha \beta g(\|x - y\|),$$

for all  $x, y, z, w \in B_r$ , and all  $\alpha, \beta, \mu, \lambda \in [0, 1]$  with  $\alpha + \beta + \mu + \lambda = 1$ .

LEMMA 1.3. (See [14].) Let  $X$  be a uniformly convex Banach space,  $C$  a nonempty closed convex subset of  $X$ , and  $T : C \rightarrow X$  be a nonexpansive mapping. Then  $I - T$  is demiclosed at 0, i.e., if  $x_n \rightarrow x$  weakly and  $x_n - Tx_n \rightarrow 0$  strongly, then  $x \in F(T)$ , where  $F(T)$  is the set of fixed point of  $T$ .

LEMMA 1.4. (See [11, Lemma 2.7].) Let  $X$  be a Banach space which satisfies Opial's condition and let  $\{x_n\}$  be a sequence in  $X$ . Let  $u, v \in X$  be such that  $\lim_{n \rightarrow \infty} \|x_n - u\|$  and  $\lim_{n \rightarrow \infty} \|x_n - v\|$  exist. If  $\{x_{n_k}\}$  and  $\{x_{m_k}\}$  are subsequences of  $\{x_n\}$  which converge weakly to  $u$  and  $v$ , respectively, then  $u = v$ .

## 2. MAIN RESULTS

Weak and strong convergence theorems of the new three-step iterative scheme (1.1) for nonexpansive nonself-mapping in a uniformly convex Banach space are given in this section. The following lemma is needed.

LEMMA 2.1. Let  $X$  be a uniformly convex Banach space, and let  $C$  be a nonempty closed convex nonexpansive retract of  $X$  with  $P$  as a nonexpansive retraction. Let  $T : C \rightarrow X$  be a nonexpansive nonself-mapping with  $F(T) \neq \emptyset$ . Let  $\{a_n\}$ ,  $\{b_n\}$ ,  $\{c_n\}$ ,  $\{\alpha_n\}$ ,  $\{\beta_n\}$ ,  $\{\gamma_n\}$ ,  $\{\mu_n\}$ , and  $\{\lambda_n\}$  be real sequences in  $[0, 1]$  such that  $a_n + \gamma_n$ ,  $b_n + c_n + \mu_n$ , and  $\alpha_n + \beta_n + \lambda_n$  are in  $[0, 1]$  for all  $n \geq 1$ , and  $\sum_{n=1}^{\infty} \gamma_n < \infty$ ,  $\sum_{n=1}^{\infty} \mu_n < \infty$ ,  $\sum_{n=1}^{\infty} \lambda_n < \infty$ , and let  $\{u_n\}$ ,  $\{v_n\}$ , and  $\{w_n\}$  be bounded sequences in  $C$ . For a given  $x_1 \in C$ , let  $\{x_n\}$ ,  $\{y_n\}$ , and  $\{z_n\}$  be the sequences defined as in (1.1).

- (i) If  $q$  is a fixed point of  $T$ , then  $\lim_{n \rightarrow \infty} \|x_n - q\|$  exists.
- (ii) If  $0 < \liminf_{n \rightarrow \infty} \alpha_n \leq \limsup_{n \rightarrow \infty} (\alpha_n + \beta_n + \lambda_n) < 1$ , then  $\lim_{n \rightarrow \infty} \|Ty_n - x_n\| = 0$ .
- (iii) If either  $0 < \liminf_{n \rightarrow \infty} \beta_n \leq \limsup_{n \rightarrow \infty} (\alpha_n + \beta_n + \lambda_n) < 1$  or  $0 < \liminf_{n \rightarrow \infty} \alpha_n$  and  $0 < \liminf_{n \rightarrow \infty} b_n \leq \limsup_{n \rightarrow \infty} (b_n + c_n + \mu_n) < 1$ , then  $\lim_{n \rightarrow \infty} \|Tz_n - x_n\| = 0$ .
- (iv) If the following conditions:
  - (1)  $0 < \liminf_{n \rightarrow \infty} \alpha_n \leq \limsup_{n \rightarrow \infty} (\alpha_n + \beta_n + \lambda_n) < 1$  and
  - (2) either  $0 < \liminf_{n \rightarrow \infty} \beta_n \leq \limsup_{n \rightarrow \infty} (\alpha_n + \beta_n + \lambda_n) < 1$  and  $\limsup_{n \rightarrow \infty} a_n < 1$  or  $0 < \liminf_{n \rightarrow \infty} b_n \leq \limsup_{n \rightarrow \infty} (b_n + c_n + \mu_n) < 1$
 are satisfied, then  $\lim_{n \rightarrow \infty} \|Tx_n - x_n\| = 0$ .

PROOF. Letting  $q \in F(T)$ , by boundedness of the sequence  $\{u_n\}$ ,  $\{v_n\}$ , and  $\{w_n\}$ , we can put

$$M = \max \left\{ \sup_{n \geq 1} \|u_n - q\|, \sup_{n \geq 1} \|v_n - q\|, \sup_{n \geq 1} \|w_n - q\| \right\}.$$

(i) For each  $n \geq 1$ , we have

$$\begin{aligned} \|x_{n+1} - q\| &= \|P(\alpha_n T y_n + \beta_n T z_n + (1 - \alpha_n - \beta_n - \lambda_n)x_n + \lambda_n w_n) - P(q)\| \\ &= \|\alpha_n T y_n + \beta_n T z_n + (1 - \alpha_n - \beta_n - \lambda_n)x_n + \lambda_n w_n - q\| \\ &\leq \alpha_n \|T y_n - q\| + \beta_n \|T z_n - q\| \\ &\quad + (1 - \alpha_n - \beta_n - \lambda_n) \|x_n - q\| + \lambda_n \|w_n - q\| \\ &\leq \alpha_n \|y_n - q\| + \beta_n \|z_n - q\| + (1 - \alpha_n - \beta_n - \lambda_n) \|x_n - q\| + M \lambda_n, \end{aligned} \quad (2.1)$$

$$\begin{aligned} \|z_n - q\| &= \|P(a_n T x_n + (1 - a_n - \gamma_n)x_n + \gamma_n u_n) - P(q)\| \\ &\leq a_n \|T x_n - q\| + (1 - a_n - \gamma_n) \|x_n - q\| + \gamma_n \|u_n - q\| \\ &\leq a_n \|x_n - q\| + (1 - a_n - \gamma_n) \|x_n - q\| + M \gamma_n \\ &\leq \|x_n - q\| + M \gamma_n, \end{aligned} \quad (2.2)$$

and

$$\begin{aligned} \|y_n - q\| &= \|P(b_n T z_n + c_n T x_n + (1 - b_n - c_n - \mu_n)x_n + \mu_n v_n) - P(q)\| \\ &\leq b_n \|T z_n - q\| + c_n \|T x_n - q\| \\ &\quad + (1 - b_n - c_n - \mu_n) \|x_n - q\| + \mu_n \|v_n - q\| \\ &\leq b_n \|z_n - q\| + c_n \|x_n - q\| + (1 - b_n - c_n - \mu_n) \|x_n - q\| + M \mu_n \\ &\leq b_n \|z_n - q\| + (1 - b_n) \|x_n - q\| + M \mu_n. \end{aligned}$$

From (2.2) we get

$$\begin{aligned} \|y_n - q\| &\leq b_n (\|x_n - q\| + M \gamma_n) + (1 - b_n) \|x_n - q\| + M \mu_n \\ &= \|x_n - q\| + \epsilon_{(1)}^n, \end{aligned} \quad (2.3)$$

where  $\epsilon_{(1)}^n = M b_n \gamma_n + M \mu_n$ . Since  $\sum_{n=1}^{\infty} \gamma_n < \infty$  and  $\sum_{n=1}^{\infty} \mu_n < \infty$ , we have  $\sum_{n=1}^{\infty} \epsilon_{(1)}^n < \infty$ .

From (2.1)-(2.3) we get

$$\begin{aligned} \|x_{n+1} - q\| &\leq \alpha_n (\|x_n - q\| + \epsilon_{(1)}^n) + \beta_n (\|x_n - q\| + M \gamma_n) \\ &\quad + (1 - \alpha_n - \beta_n - \lambda_n) \|x_n - q\| + M \lambda_n \\ &= \alpha_n \|x_n - q\| + \alpha_n \epsilon_{(1)}^n + \beta_n \|x_n - q\| + M \beta_n \gamma_n \\ &\quad + (1 - \alpha_n - \beta_n - \lambda_n) \|x_n - q\| + M \lambda_n \\ &\leq \|x_n - q\| + \epsilon_{(2)}^n, \end{aligned} \quad (2.4)$$

where  $\epsilon_{(2)}^n = \alpha_n \epsilon_{(1)}^n + M \beta_n \gamma_n + M \lambda_n$ . Since  $\sum_{n=1}^{\infty} \epsilon_{(2)}^n < \infty$  we obtained from (2.4) and Lemma 1.1 that  $\lim_{n \rightarrow \infty} \|x_n - q\|$  exists.

(ii) By (i) we have that  $\lim_{n \rightarrow \infty} \|x_n - q\|$  exists for any  $q \in F(T)$ . It follows from (2.2) and (2.3) that  $\{x_n - q\}$ ,  $\{Tx_n - q\}$ ,  $\{z_n - q\}$ ,  $\{Tz_n - q\}$ ,  $\{y_n - q\}$ , and  $\{Ty_n - q\}$  are bounded sequences. This allows us to put

$$K = \max \left\{ M, \sup_{n \geq 1} \|x_n - q\|, \sup_{n \geq 1} \|Tx_n - q\|, \sup_{n \geq 1} \|z_n - q\|, \right. \\ \left. \sup_{n \geq 1} \|Tz_n - q\|, \sup_{n \geq 1} \|y_n - q\|, \sup_{n \geq 1} \|Ty_n - q\| \right\}.$$

Since  $0 < \liminf_{n \rightarrow \infty} \alpha_n \leq \limsup_{n \rightarrow \infty} (\alpha_n + \beta_n + \lambda_n) < 1$ , it follows from (2.2) and (2.3) that

$$\|z_n - q\|^2 \leq \|x_n - q\|^2 + \epsilon_{(3)}^n, \quad (2.5)$$

$$\|y_n - q\|^2 \leq \|x_n - q\|^2 + \epsilon_{(4)}^n, \quad (2.6)$$

where  $\epsilon_{(3)}^n = M^2 \gamma_n^2 + 2MK\gamma_n$ , and  $\epsilon_{(4)}^n = (\epsilon_{(1)}^n)^2 + 2K\epsilon_{(1)}^n$ . Since  $\sum_{n=1}^{\infty} \epsilon_{(3)}^n < \infty$  and  $\sum_{n=1}^{\infty} \epsilon_{(4)}^n < \infty$ , by Lemma 1.2, there is a continuous strictly increasing convex function  $g: [0, \infty) \rightarrow [0, \infty)$ ,  $g(0) = 0$ , such that

$$\|\lambda x + \beta y + \gamma z + \mu w\|^2 \leq \lambda \|x\|^2 + \beta \|y\|^2 + \gamma \|z\|^2 + \mu \|w\|^2 - \lambda \beta g(\|x - y\|), \quad (2.7)$$

for all  $x, y, z, w \in B_K$  and all  $\lambda, \beta, \gamma, \mu \in [0, 1]$  with  $\lambda + \beta + \gamma = 1$ . By (2.5)–(2.7), we have

$$\begin{aligned} \|x_{n+1} - q\|^2 &= \|P(\alpha_n Ty_n + \beta_n Tz_n + (1 - \alpha_n - \beta_n - \lambda_n)x_n + \lambda_n w_n) - P(q)\|^2 \\ &\leq \|\alpha_n(Ty_n - q) + \beta_n(Tz_n - q) \\ &\quad + (1 - \alpha_n - \beta_n - \lambda_n)(x_n - q) + \lambda_n(w_n - q)\|^2 \\ &\leq \alpha_n \|Ty_n - q\|^2 + \beta_n \|Tz_n - q\|^2 \\ &\quad + (1 - \alpha_n - \beta_n - \lambda_n) \|x_n - q\|^2 + \lambda_n \|w_n - q\|^2 \\ &\quad - \alpha_n(1 - \alpha_n - \beta_n - \lambda_n) g(\|Ty_n - x_n\|) \\ &\leq \alpha_n \|y_n - q\|^2 + \beta_n \|z_n - q\|^2 + (1 - \alpha_n - \beta_n - \lambda_n) \|x_n - q\|^2 \\ &\quad + K^2 \lambda_n - \alpha_n(1 - \alpha_n - \beta_n - \lambda_n) g(\|Ty_n - x_n\|) \\ &\leq \alpha_n (\|x_n - q\|^2 + \epsilon_{(4)}^n) + \beta_n (\|x_n - q\|^2 + \epsilon_{(3)}^n) \\ &\quad + (1 - \alpha_n - \beta_n - \lambda_n) \|x_n - q\|^2 + K^2 \lambda_n \\ &\quad - \alpha_n(1 - \alpha_n - \beta_n - \lambda_n) g(\|Ty_n - x_n\|) \\ &= \alpha_n \|x_n - q\|^2 + \alpha_n \epsilon_{(4)}^n + \beta_n \|x_n - q\|^2 + \beta_n \epsilon_{(3)}^n \\ &\quad + (1 - \alpha_n - \beta_n - \lambda_n) \|x_n - q\|^2 + K^2 \lambda_n \\ &\quad - \alpha_n(1 - \alpha_n - \beta_n - \lambda_n) g(\|Ty_n - x_n\|) \\ &\leq \|x_n - q\|^2 + \epsilon_{(5)}^n - \alpha_n(1 - \alpha_n - \beta_n - \lambda_n) g(\|Ty_n - x_n\|), \end{aligned} \quad (2.8)$$

where  $\epsilon_{(5)}^n = \alpha_n \epsilon_{(4)}^n + \beta_n \epsilon_{(3)}^n + K^2 \lambda_n$ . It is worth noting here that  $\sum_{n=1}^{\infty} \epsilon_{(5)}^n < \infty$  since  $\sum_{n=1}^{\infty} \epsilon_{(4)}^n < \infty$ ,  $\sum_{n=1}^{\infty} \epsilon_{(3)}^n < \infty$ , and  $\sum_{n=1}^{\infty} \lambda_n < \infty$ . Since  $0 < \liminf_{n \rightarrow \infty} \alpha_n \leq$

Since  $(\alpha_n + \beta_n + \lambda_n) < 1$ , there exists  $n_0 \in \mathbb{N}$  and  $\delta_1, \delta_2 \in (0, 1)$  such that  $0 < \delta_1 < \alpha_n$  and  $\alpha_n + \beta_n + \lambda_n < \delta_2 < 1$  for all  $n \geq n_0$ . Hence, by (2.8), we have

$$\begin{aligned} (\delta_1 - \delta_2) \sum_{n=n_0}^m g \|Ty_n - x_n\| &< \sum_{n=n_0}^m (\|x_n - q\|^2 - \|x_{n+1} - q\|^2) + \sum_{n=n_0}^m \epsilon_{(5)}^n \\ &= \|x_{n_0} - q\|^2 + \sum_{n=n_0}^m \epsilon_{(5)}^n. \end{aligned} \quad (2.9)$$

Since  $\sum_{n=n_0}^\infty \epsilon_{(5)}^n < \infty$ , by letting  $m \rightarrow \infty$  in (2.9) we get  $\sum_{n=n_0}^\infty g \|Ty_n - x_n\| < \infty$ , and  $\lim_{n \rightarrow \infty} g \|Ty_n - x_n\| = 0$ . Since  $g$  is strictly increasing and continuous at 0 with  $g(0) = 0$ , it follows that  $\lim_{n \rightarrow \infty} \|Ty_n - x_n\| = 0$ .

Now, we assume that  $0 < \liminf_{n \rightarrow \infty} \beta_n \leq \limsup_{n \rightarrow \infty} (\alpha_n + \beta_n + \lambda_n) < 1$ . By (2.7), we

$$\begin{aligned} \|x_n - q\|^2 &\leq \alpha_n \|y_n - q\|^2 + \beta_n \|z_n - q\|^2 + (1 - \alpha_n - \beta_n - \lambda_n) \|x_n - q\|^2 + K^2 \lambda_n \\ &\quad - \beta_n (1 - \alpha_n - \beta_n - \lambda_n) g \|Tz_n - x_n\| \\ &\leq \alpha_n (\|x_n - q\|^2 + \epsilon_{(4)}^n) + \beta_n (\|x_n - q\|^2 + \epsilon_{(3)}^n) \\ &\quad + (1 - \alpha_n - \beta_n - \lambda_n) \|x_n - q\|^2 + K^2 \lambda_n \\ &\quad - \beta_n (1 - \alpha_n - \beta_n - \lambda_n) g \|Tz_n - x_n\| \\ &= \alpha_n \|x_n - q\|^2 + \alpha_n \epsilon_{(4)}^n + \beta_n \|x_n - q\|^2 + \beta_n \epsilon_{(3)}^n \\ &\quad + (1 - \alpha_n - \beta_n - \lambda_n) \|x_n - q\|^2 + K^2 \lambda_n \\ &\quad - \beta_n (1 - \alpha_n - \beta_n - \lambda_n) g \|Tz_n - x_n\| \\ &\leq \|x_n - q\|^2 + \epsilon_{(5)}^n - \beta_n (1 - \alpha_n - \beta_n - \lambda_n) g \|Tz_n - x_n\|, \end{aligned} \quad (2.10)$$

where  $\epsilon_{(5)}^n = \alpha_n \epsilon_{(4)}^n + \beta_n \epsilon_{(3)}^n + K^2 \lambda_n$ . Since  $0 < \liminf_{n \rightarrow \infty} \beta_n \leq \limsup_{n \rightarrow \infty} (\alpha_n + \beta_n + \lambda_n) < 1$ , there exists  $n_0 \in \mathbb{N}$  and  $\delta_1, \delta_2 \in (0, 1)$  such that  $0 < \delta_1 < \beta_n$  and  $\alpha_n + \beta_n + \lambda_n < \delta_2 < 1$  for all  $n \geq n_0$ . Hence, by (2.10), we have  $\epsilon_{(5)}^n = \alpha_n \epsilon_{(4)}^n + \beta_n \epsilon_{(3)}^n + K^2 \lambda_n$ .

$$\begin{aligned} (\delta_1 - \delta_2) \sum_{n=n_0}^m g \|Tz_n - x_n\| &< \sum_{n=n_0}^m (\|x_n - q\|^2 - \|x_{n+1} - q\|^2) + \sum_{n=n_0}^m \epsilon_{(5)}^n \\ &= \|x_{n_0} - q\|^2 + \sum_{n=n_0}^m \epsilon_{(5)}^n. \end{aligned} \quad (2.11)$$

Since  $\sum_{n=n_0}^\infty \epsilon_{(5)}^n < \infty$ , by letting  $m \rightarrow \infty$  in (2.11) we get  $\sum_{n=n_0}^\infty g \|Tz_n - x_n\| < \infty$ , and  $\lim_{n \rightarrow \infty} g \|Tz_n - x_n\| = 0$ . Since  $g$  is strictly increasing and continuous at 0 with  $g(0) = 0$ , it follows that  $\lim_{n \rightarrow \infty} \|Tz_n - x_n\| = 0$ .

Now, we assume that  $0 < \liminf_{n \rightarrow \infty} \alpha_n$  and  $\liminf_{n \rightarrow \infty} b_n \leq \limsup_{n \rightarrow \infty} (b_n + c_n + \mu_n) < 1$ . By (2.7), we have

$$\begin{aligned} \|x_n - q\|^2 &= \|P(b_n Tz_n + c_n Tx_n + (1 - b_n - c_n - \mu_n)x_n + \mu_n v_n) - P(q)\|^2 \\ &\leq \|b_n(Tz_n - q) + c_n(Tx_n - q) + (1 - b_n - c_n - \mu_n)(x_n - q) + \mu_n(v_n - q)\|^2 \\ &\leq b_n \|Tz_n - q\|^2 + c_n \|Tx_n - q\|^2 \end{aligned} \quad (2.12)$$

$$\begin{aligned}
& + (1 - b_n - c_n - \mu_n)\|x_n - q\|^2 + \mu_n\|v_n - q\|^2 \\
& - b_n(1 - b_n - c_n - \mu_n)g\|Tz_n - x_n\| \\
& \leq b_n\|z_n - q\|^2 + c_n\|x_n - q\|^2 + (1 - b_n - c_n - \mu_n)\|x_n - q\|^2 + \mu_n K^2 \\
& \quad - b_n(1 - b_n - c_n - \mu_n)g\|Tz_n - x_n\| \tag{2.12}(\text{cont.}) \\
& \leq b_n\left(\|x_n - q\|^2 + \epsilon_{(3)}^n\right) + c_n\|x_n - q\|^2 + (1 - b_n - c_n - \mu_n)\|x_n - q\|^2 + \mu_n K^2 \\
& \leq b_n\left(\|x_n - q\|^2 + \epsilon_{(3)}^n\right) + c_n\|x_n - q\|^2 + (1 - b_n - c_n - \mu_n)\|x_n - q\|^2 + \mu_n K^2 \\
& \quad - b_n(1 - b_n - c_n - \mu_n)g\|Tz_n - x_n\| \\
& \leq \|x_n - q\|^2 + \epsilon_{(6)}^n - b_n(1 - b_n - c_n - \mu_n)g\|Tz_n - x_n\|,
\end{aligned}$$

where  $\epsilon_{(6)}^n = b_n\epsilon_{(3)}^n + \mu_n K^2$ .

By (2.5), (2.7), and (2.12), we also have

$$\begin{aligned}
\|x_{n+1} - q\|^2 & = \|P(\alpha_n T y_n + \beta_n T z_n + (1 - \alpha_n - \beta_n - \lambda_n)x_n + \lambda_n w_n) - P(q)\|^2 \\
& \leq \|\alpha_n(T y_n - q) + \beta_n(T z_n - q) \\
& \quad + (1 - \alpha_n - \beta_n - \lambda_n)(x_n - q) + \lambda_n(w_n - q)\|^2 \\
& \leq \alpha_n\|y_n - q\|^2 + \beta_n\|z_n - q\|^2 + (1 - \alpha_n - \beta_n - \lambda_n)\|x_n - q\|^2 + K^2\lambda_n \\
& = \alpha_n\left(\|x_n - q\|^2 + \epsilon_{(6)}^n - b_n(1 - b_n - c_n - \mu_n)g\|Tz_n - x_n\|\right) \\
& \quad + \beta_n\left(\|x_n - q\|^2 + \epsilon_{(3)}^n\right) + (1 - \alpha_n - \beta_n - \lambda_n)\|x_n - q\|^2 + K^2\lambda_n \\
& = \alpha_n\|x_n - q\|^2 + \alpha_n\epsilon_{(6)}^n - \alpha_nb_n(1 - b_n - c_n - \mu_n)g\|Tz_n - x_n\| \\
& \quad + \beta_n\|x_n - q\|^2 + \beta_n\epsilon_{(3)}^n + (1 - \alpha_n - \beta_n - \lambda_n)\|x_n - q\|^2 + K^2\lambda_n \\
& \leq \|x_n - q\|^2 + \epsilon_{(7)}^n - \alpha_nb_n(1 - b_n - c_n - \mu_n)g\|Tz_n - x_n\|,
\end{aligned} \tag{2.13}$$

where  $\epsilon_{(7)}^n = \alpha_n\epsilon_{(6)}^n + \beta_n\epsilon_{(3)}^n + K^2\lambda_n$ .

It is worth noting here that  $\sum_{n=1}^{\infty} \epsilon_{(7)}^n < \infty$  since  $\sum_{n=1}^{\infty} \epsilon_{(6)}^n < \infty$ ,  $\sum_{n=1}^{\infty} \epsilon_{(3)}^n < \infty$ , and  $\sum_{n=1}^{\infty} \lambda_n < \infty$ .

By our assumption  $0 < \liminf_{n \rightarrow \infty} \alpha_n$  and  $0 < \liminf_{n \rightarrow \infty} b_n \leq \limsup_{n \rightarrow \infty} (b_n + c_n + \mu_n) < 1$ , there exist  $n_0 \in \mathbb{N}$  and  $\delta_1, \delta_2 \in (0, 1)$  such that  $0 < \delta_1 < \alpha_n$ ,  $0 < \delta_1 < b_n$ , and  $b_n + c_n + \mu_n < \delta_2 < 1$  for all  $n \geq n_0$ . Hence, by (2.13), we have

$$\begin{aligned}
\delta_1^2(1 - \delta_2) \sum_{n=n_0}^m g\|Tz_n - x_n\| & < \sum_{n=n_0}^m (\|x_n - q\|^2 - \|x_{n+1} - q\|^2) + \sum_{n=n_0}^m \epsilon_{(7)}^n \\
& = \|x_{n_0} - q\|^2 + \sum_{n=n_0}^m \epsilon_{(7)}^n.
\end{aligned} \tag{2.14}$$

Since  $\sum_{n=n_0}^{\infty} \epsilon_{(7)}^n < \infty$ , by letting  $m \rightarrow \infty$  in (2.14) we get  $\sum_{n=n_0}^{\infty} g\|Tz_n - x_n\| < \infty$ , and therefore  $\lim_{n \rightarrow \infty} g\|Tz_n - x_n\| = 0$ . Since  $g$  is strictly increasing and continuous at 0 with  $g(0) = 0$ , it follows that  $\lim_{n \rightarrow \infty} \|Tz_n - x_n\| = 0$ .

(iv) Suppose that Conditions (1) and (2) are satisfied. Then by (ii) and (iii), we have

$$\lim_{n \rightarrow \infty} \|Ty_n - x_n\| = 0 \quad \text{and} \quad \lim_{n \rightarrow \infty} \|Tz_n - x_n\| = 0. \quad (2.15)$$

From  $z_n = P(a_nTx_n + (1 - a_n - \gamma_n)x_n + \gamma_nu_n)$  and  $y_n = P(b_nTz_n + c_nTx_n + (1 - b_n - c_n - \mu_n)x_n + \mu_nv_n)$ , we have  $\|z_n - x_n\| \leq a_n\|Tx_n - x_n\| + \gamma_n\|u_n - x_n\|$  and  $\|y_n - x_n\| \leq b_n\|Tz_n - x_n\| + c_n\|Tx_n - x_n\| + \mu_n\|v_n - x_n\|$ .

It follows that

$$\begin{aligned} \|Tx_n - x_n\| &\leq \|Tx_n - Tz_n\| + \|Tz_n - x_n\| \\ &\leq \|x_n - z_n\| + \|Tz_n - x_n\| \\ &\leq a_n\|Tx_n - x_n\| + \gamma_n\|u_n - z_n\| + \|Tz_n - x_n\|, \end{aligned}$$

which implies

$$(1 - a_n)\|Tx_n - x_n\| \leq \gamma_n\|u_n - z_n\| + \|Tz_n - x_n\|.$$

If  $\limsup_{n \rightarrow \infty} a_n < 1$ , this together with (2.15) and  $\lim_{n \rightarrow \infty} \gamma_n = 0$  imply that  $\lim_{n \rightarrow \infty} \|Tx_n - x_n\| = 0$ .

If  $\limsup_{n \rightarrow \infty} (b_n + c_n + \mu_n) < 1$ , there exist a positive integer  $N_0$  and  $\eta \in (0, 1)$  such that

$$c_n \leq b_n + c_n + \mu_n < \eta, \quad \forall n \geq N_0.$$

Then for  $n \geq N_0$ , we have

$$\begin{aligned} \|Tx_n - x_n\| &\leq \|Tx_n - Ty_n\| + \|Ty_n - x_n\| \\ &\leq \|x_n - y_n\| + \|Ty_n - x_n\| \\ &\leq b_n\|Tz_n - x_n\| + c_n\|Tx_n - x_n\| \\ &\quad + \mu_n\|v_n - x_n\| + \|Ty_n - x_n\| \\ &\leq b_n\|Tz_n - x_n\| + \eta\|Tx_n - x_n\| \\ &\quad + \mu_n\|v_n - x_n\| + \|Ty_n - x_n\|. \end{aligned}$$

Hence,

$$(1 - \eta)\|Tx_n - x_n\| \leq b_n\|Tz_n - x_n\| + \mu_n\|v_n - x_n\| + \|Ty_n - x_n\|.$$

This together with (2.15) and the fact that  $\mu_n \rightarrow 0$  as  $n \rightarrow \infty$  imply

$$\lim_{n \rightarrow \infty} \|Tx_n - x_n\| = 0. \quad \blacksquare$$

**THEOREM 2.2.** Let  $X$  be a uniformly convex Banach space, and let  $C$  be a nonempty closed convex nonexpansive retract of  $X$  with  $P$  as a nonexpansive retraction. Let  $T : C \rightarrow X$  be a completely continuous nonexpansive nonself-mapping with  $F(T) \neq \emptyset$ . Let  $\{a_n\}$ ,  $\{b_n\}$ ,  $\{c_n\}$ ,  $\{\alpha_n\}$ ,  $\{\beta_n\}$ ,  $\{\gamma_n\}$ ,  $\{\mu_n\}$ , and  $\{\lambda_n\}$  be sequences of real numbers in  $[0, 1]$  with  $a_n + \gamma_n \in [0, 1]$ ,  $b_n + c_n + \mu_n \in [0, 1]$ , and  $\alpha_n + \beta_n + \lambda_n \in [0, 1]$  for all  $n \geq 1$ , and  $\sum_{n=1}^{\infty} \gamma_n < \infty$ ,  $\sum_{n=1}^{\infty} \mu_n < \infty$ ,  $\sum_{n=1}^{\infty} \lambda_n < \infty$ . If

- (i)  $0 < \min\{\liminf_{n \rightarrow \infty} \alpha_n, \liminf_{n \rightarrow \infty} \beta_n\} \leq \limsup_{n \rightarrow \infty} (\alpha_n + \beta_n + \lambda_n) < 1$ , and  $\limsup_{n \rightarrow \infty} a_n < 1$ , or
- (ii)  $0 < \liminf_{n \rightarrow \infty} \alpha_n \leq \limsup_{n \rightarrow \infty} (\alpha_n + \beta_n + \lambda_n) < 1$ , and  $0 < \liminf_{n \rightarrow \infty} b_n \leq \limsup_{n \rightarrow \infty} (b_n + c_n + \mu_n) < 1$ ,

then the sequences  $\{x_n\}$ ,  $\{y_n\}$ , and  $\{z_n\}$  defined by the iterative scheme (1.1) converge strongly to a fixed point of  $T$ .

PROOF. It follows from Lemma 2.1(i) that  $\{x_n\}$  is bounded. Again by Lemma 2.1, we have

$$\begin{aligned}\lim_{n \rightarrow \infty} \|Ty_n - x_n\| &= 0, \\ \lim_{n \rightarrow \infty} \|Tz_n - x_n\| &= 0, \\ \lim_{n \rightarrow \infty} \|Tx_n - x_n\| &= 0.\end{aligned}\tag{2.16}$$

Since  $T$  is completely continuous and  $\{x_n\}$  is bounded, there exists a subsequence  $\{x_{n_k}\}$  of  $\{x_n\}$  such that  $\{Tx_{n_k}\}$  converges. Hence, by  $\lim_{n \rightarrow \infty} \|Tx_n - x_n\| = 0$ , it follows that  $\{x_{n_k}\}$  converges. Let  $\lim_{n \rightarrow \infty} x_{n_k} = q$ . By continuity of  $T$  and (2.16) we have that  $Tq = q$ , so  $q$  is a fixed point of  $T$ . By Lemma 2.1(i),  $\lim_{n \rightarrow \infty} \|x_n - q\|$  exists. But  $\lim_{k \rightarrow \infty} \|x_{n_k} - q\| = 0$ , so  $\lim_{n \rightarrow \infty} \|x_n - q\| = 0$ . By (2.16), we have

$$\begin{aligned}\|y_n - x_n\| &= \|P(b_nTx_n + c_nTx_n + (1 - b_n - c_n - \mu_n)x_n + \mu_nv_n) - P(x_n)\| \\ &\leq b_n\|Tx_n - x_n\| + c_n\|Tx_n - x_n\| + \mu_n\|v_n - x_n\| \\ &\rightarrow 0, \quad \text{as } n \rightarrow \infty,\end{aligned}$$

and

$$\begin{aligned}\|z_n - x_n\| &= \|P(a_nTx_n + (1 - a_n - \gamma_n)x_n + \gamma_nu_n) - P(x_n)\| \\ &\leq a_n\|Tx_n - x_n\| + \gamma_n\|u_n - x_n\| \\ &\rightarrow 0, \quad \text{as } n \rightarrow \infty.\end{aligned}$$

It follows that  $\lim_{n \rightarrow \infty} y_n = q$  and  $\lim_{n \rightarrow \infty} z_n = q$ .

If  $T$  is a self-mapping, then the iterative scheme (1.1) reduces to that of (1.3) and the following result is directly obtained by Theorem 2.2.

**THEOREM 2.3.** Let  $X$  be a uniformly convex Banach space, and  $C$  a nonempty closed convex subset of  $X$ . Let  $T$  be a completely continuous nonexpansive self-mapping of  $C$  with  $F(T) \neq \emptyset$ . Let  $\{a_n\}$ ,  $\{b_n\}$ ,  $\{c_n\}$ ,  $\{\alpha_n\}$ ,  $\{\beta_n\}$  be sequences of real numbers in  $[0, 1]$  with  $b_n + c_n \in [0, 1]$  and  $\alpha_n + \beta_n \in [0, 1]$  for all  $n \geq 1$ . If

- (i)  $0 < \min\{\liminf_{n \rightarrow \infty} \alpha_n, \liminf_{n \rightarrow \infty} \beta_n\} \leq \limsup_{n \rightarrow \infty} (\alpha_n + \beta_n + \lambda_n) < 1$ , and  $\limsup_{n \rightarrow \infty} a_n < 1$ , or
- (ii)  $0 < \liminf_{n \rightarrow \infty} \alpha_n \leq \limsup_{n \rightarrow \infty} (\alpha_n + \beta_n + \lambda_n) < 1$ , and  $0 < \liminf_{n \rightarrow \infty} b_n \leq \limsup_{n \rightarrow \infty} (b_n + c_n + \mu_n) < 1$ ,

then the sequences  $\{x_n\}$ ,  $\{y_n\}$ , and  $\{z_n\}$  defined by iterations (1.3) converge strongly to a fixed point of  $T$ .

When  $c_n = \beta_n = \gamma_n = \mu_n = \lambda_n \equiv 0$  in Theorem 2.2, the following result is obtained.

**THEOREM 2.4.** Let  $X$  be a uniformly convex Banach space, and let  $C$  be a nonempty closed convex nonexpansive retract of  $X$  with  $P$  as a nonexpansive retraction. Let  $T : C \rightarrow X$  be a completely continuous nonexpansive nonself-mapping with  $F(T) \neq \emptyset$ . Let  $\{a_n\}$ ,  $\{b_n\}$ ,  $\{\alpha_n\}$  be real sequences in  $[0, 1]$  satisfying

- (i)  $0 < \liminf_{n \rightarrow \infty} b_n \leq \limsup_{n \rightarrow \infty} b_n < 1$ , and
- (ii)  $0 < \liminf_{n \rightarrow \infty} \alpha_n \leq \limsup_{n \rightarrow \infty} \alpha_n < 1$ .

For a given  $x_1 \in C$ , define

$$\begin{aligned} z_n &= P(a_n T x_n + (1 - a_n)x_n), \\ y_n &= P(b_n T z_n + (1 - b_n)x_n), \quad n \geq 1, \\ x_{n+1} &= P(\alpha_n T y_n + (1 - \alpha_n)x_n). \end{aligned}$$

Then  $\{x_n\}$ ,  $\{y_n\}$ , and  $\{z_n\}$  converge strongly to a fixed point of  $T$ .

When  $a_n = c_n = \beta_n = \gamma_n = \mu_n = \lambda_n \equiv 0$  in Theorem 2.2, we obtain the following result.

**THEOREM 2.5.** *Let  $X$  be a uniformly convex Banach space, and let  $C$  be a nonempty closed convex nonexpansive retract of  $X$  with  $P$  as a nonexpansive retraction. Let  $T : C \rightarrow X$  be a completely continuous nonexpansive nonself-mapping with  $F(T) \neq \emptyset$ . Let  $\{b_n\}$ ,  $\{\alpha_n\}$  be a real sequence in  $[0, 1]$  satisfying*

- (i)  $0 < \liminf_{n \rightarrow \infty} b_n \leq \limsup_{n \rightarrow \infty} b_n < 1$ , and
- (ii)  $0 < \liminf_{n \rightarrow \infty} \alpha_n \leq \limsup_{n \rightarrow \infty} \alpha_n < 1$ .

For a given  $x_1 \in C$ , define

$$\begin{aligned} y_n &= P(b_n T x_n + (1 - b_n)x_n), \\ x_{n+1} &= P(\alpha_n T y_n + (1 - \alpha_n)x_n), \quad n \geq 1. \end{aligned}$$

Then  $\{x_n\}$  and  $\{y_n\}$  converge strongly to a fixed point of  $T$ .

The mapping  $T : C \rightarrow X$  with  $F(T) \neq \emptyset$  is said to satisfy Condition (A) if there is a nondecreasing function  $f : [0, \infty) \rightarrow [0, \infty)$  with  $f(0) = 0$  and  $f(r) > 0$  for all  $r \in (0, \infty)$  such that for all  $x \in C$ ,

$$\|x - Tx\| \geq f(d(x, F(T))).$$

The following result gives a strong convergence theorem for nonexpansive nonself-mapping in a uniformly convex Banach space satisfying Condition (A).

**THEOREM 2.6.** *Let  $X$  be a uniformly convex Banach space, and let  $C$  be a nonempty closed convex nonexpansive retract of  $X$  with  $P$  as a nonexpansive retraction. Let  $T : C \rightarrow X$  be a nonexpansive nonself-mapping with  $F(T) \neq \emptyset$ . Let  $\{a_n\}$ ,  $\{b_n\}$ ,  $\{c_n\}$ ,  $\{\alpha_n\}$ ,  $\{\beta_n\}$ ,  $\{\gamma_n\}$ ,  $\{\mu_n\}$ , and  $\{\lambda_n\}$  be sequences of real numbers in  $[0, 1]$  with  $a_n + \gamma_n \in [0, 1]$ ,  $b_n + c_n + \mu_n \in [0, 1]$ , and  $a_n + \beta_n + \lambda_n \in [0, 1]$  for all  $n \geq 1$ , and  $\sum_{n=1}^{\infty} \gamma_n < \infty$ ,  $\sum_{n=1}^{\infty} \mu_n < \infty$ ,  $\sum_{n=1}^{\infty} \lambda_n < \infty$ . Suppose that  $T$  satisfies Condition (A). If*

- (i)  $0 < \min\{\liminf_{n \rightarrow \infty} \alpha_n, \liminf_{n \rightarrow \infty} \beta_n\} \leq \limsup_{n \rightarrow \infty} (\alpha_n + \beta_n + \lambda_n) < 1$ , and  $\limsup_{n \rightarrow \infty} a_n < 1$ , or
- (ii)  $0 < \liminf_{n \rightarrow \infty} \alpha_n \leq \limsup_{n \rightarrow \infty} (\alpha_n + \beta_n + \lambda_n) < 1$ , and  $0 < \liminf_{n \rightarrow \infty} b_n \leq \limsup_{n \rightarrow \infty} (b_n + c_n + \mu_n) < 1$ ,

then the sequences  $\{x_n\}$  defined by the iterative scheme (1.1) converge strongly to some fixed point of  $T$ .

**PROOF.** Let  $q \in F(T)$ . Then, as in Lemma 2.1,  $\{x_n\}$  is bounded,  $\lim_{n \rightarrow \infty} \|x_n - q\|$  exists, and

$$\|x_{n+1} - q\| \leq \|x_n - q\| + \epsilon_{(2)}^n,$$

where  $\sum_{n=1}^{\infty} \epsilon_{(2)}^n < \infty$  for all  $n \geq 1$ . This implies that  $d(x_{n+1}, F(T)) \leq d(x_n, F(T)) + \epsilon_{(2)}^n$  and so, by Lemma 1.1,  $\lim_{n \rightarrow \infty} d(x_n, F(T))$  exists. Also, by Lemma 2.1,  $\lim_{n \rightarrow \infty} \|x_n - Tx_n\| = 0$ . Since  $T$  satisfies Condition (A), we conclude that  $\lim_{n \rightarrow \infty} d(x_n, F(T)) = 0$ . Next we show that  $\{x_n\}$  is a Cauchy sequence.

Since  $\lim_{n \rightarrow \infty} d(x_n, F(T)) = 0$  and  $\sum_{n=1}^{\infty} \epsilon_{(2)}^n < \infty$ , given any  $\epsilon < 0$ , there exists a natural number  $n_0$  such that  $d(x_n, F(T)) < \epsilon/4$  and  $\sum_{k=n_0}^n \epsilon_{(2)}^k \epsilon/2$  for all  $n \geq n_0$ . So we can find  $y^* \in F(T)$  such that  $\|x_{n_0} - y^*\| < \epsilon/4$ . For  $n \geq n_0$  and  $m \geq 1$ , we have

$$\begin{aligned} \|x_{n+m} - x_n\| &= \|x_{n+m} - y^*\| + \|x_n - y^*\| \\ &\leq \|x_{n_0} - y^*\| + \|x_{n_0} - y^*\| + \sum_{k=n_0}^n \epsilon_{(2)}^k \\ &< \frac{\epsilon}{4} + \frac{\epsilon}{4} + \frac{\epsilon}{2} = \epsilon. \end{aligned}$$

This shows that  $\{x_n\}$  is a Cauchy sequence and so is convergent since  $X$  is complete. Let  $\lim_{n \rightarrow \infty} x_n = u$ . Then  $d(u, F(T)) = 0$ . It follows that  $u \in F(T)$ . This completes the proof.  $\square$

For  $\alpha_n = c_n = \beta_n = \gamma_n = \mu_n = \lambda_n \equiv 0$ , the iterative scheme (1.1) reduces to that of (1.2) and the following result is directly obtained by Theorem 2.6.

**THEOREM 2.7.** (See [1, Theorem 3.6].) Let  $X$  be a uniformly convex Banach space, and let  $C$  be a nonempty closed convex nonexpansive retract of  $X$  with  $P$  as a nonexpansive retraction. Let  $T: C \rightarrow X$  be a nonexpansive nonself-mapping with  $F(T) \neq \emptyset$ . Let  $\{\alpha_n\}$  and  $\{\beta_n\}$  be sequences in  $[\epsilon, 1 - \epsilon]$  for some  $\epsilon \in (0, 1)$ . Suppose that  $T$  satisfies Condition (A). Then the sequences  $\{z_n\}$  defined by the iterative scheme (1.2) converge strongly to some fixed point of  $T$ .

In the next result, we prove weak convergence of the iterative scheme (1.1) for nonexpansive nonself-mapping in a uniformly convex Banach space satisfying Opial's condition.

**THEOREM 2.8.** Let  $X$  be a uniformly convex Banach space which satisfies Opial's condition, and  $C$  a nonempty closed convex nonexpansive retract of  $X$  with  $P$  as a nonexpansive retraction. Let  $T: C \rightarrow X$  be a nonexpansive nonself-mapping with  $F(T) \neq \emptyset$ . Let  $\{\alpha_n\}$ ,  $\{\beta_n\}$ ,  $\{\gamma_n\}$ ,  $\{\mu_n\}$ ,  $\{\lambda_n\}$  be sequences of real numbers in  $[0, 1]$  with  $\alpha_n + \gamma_n$ ,  $\beta_n + c_n + \mu_n$ , and  $\alpha_n + \beta_n + \lambda_n$  are in  $[0, 1]$  for all  $n \geq 1$ , and  $\sum_{n=1}^{\infty} \gamma_n < \infty$ ,  $\sum_{n=1}^{\infty} \mu_n < \infty$ ,  $\sum_{n=1}^{\infty} \lambda_n < \infty$ . If

- (i)  $0 < \min\{\liminf_{n \rightarrow \infty} \alpha_n, \liminf_{n \rightarrow \infty} \beta_n\} \leq \limsup_{n \rightarrow \infty} (\alpha_n + \beta_n + \lambda_n) < 1$ , and  $\limsup_{n \rightarrow \infty} \alpha_n < 1$ , or
- (ii)  $0 < \liminf_{n \rightarrow \infty} \alpha_n \leq \limsup_{n \rightarrow \infty} (\alpha_n + \beta_n + \lambda_n) < 1$  and  $0 < \liminf_{n \rightarrow \infty} \beta_n \leq \limsup_{n \rightarrow \infty} (\beta_n + c_n + \mu_n) < 1$ ,

then the sequence  $\{x_n\}$ ,  $\{y_n\}$ , and  $\{z_n\}$  defined by the iterative scheme (1.1) converges weakly to a fixed point of  $T$ .

**PROOF.** It follows from Lemma 2.1 that  $\lim_{n \rightarrow \infty} \|Tx_n - x_n\| = 0$  and  $\lim_{n \rightarrow \infty} \|Tz_n - z_n\| = 0$ . Since  $X$  is uniformly convex and  $\{x_n\}$  is bounded, we may assume that  $x_n \rightarrow u$  weakly as  $n \rightarrow \infty$  without loss of generality. By Lemma 1.3, we have  $u \in F(T)$ . Suppose that subsequences  $\{x_{n_k}\}$  and  $\{x_{m_k}\}$  of  $\{x_n\}$  converge weakly to  $u$  and  $v$ , respectively. From Lemma 1.3,  $u, v \in F(T)$ . By Lemma 2.1(i),  $\lim_{n \rightarrow \infty} \|x_n - u\|$  and  $\lim_{n \rightarrow \infty} \|x_n - v\|$  exist. It follows from Lemma 1.4 that  $u = v$ . Therefore  $\{x_n\}$  converges weakly to a fixed point  $u$  of  $T$ . Since  $\|y_n - x_n\| \leq b_n \|Tx_n - x_n\| + c_n \|Tx_n - x_n\| + \mu_n \|v_n - x_n\| \rightarrow 0$  (as  $n \rightarrow \infty$ ) and  $\|z_n - x_n\| \leq a_n \|Tx_n - x_n\| + \gamma_n \|u_n - x_n\| \rightarrow 0$  (as  $n \rightarrow \infty$ ) and  $x_n \rightarrow u$  weakly as  $n \rightarrow \infty$ , it follows that  $y_n \rightarrow u$  and  $z_n \rightarrow u$  weakly as  $n \rightarrow \infty$ .  $\square$

## REFERENCES

1. N. Shahzad, Approximating fixed points of non-self nonexpansive mappings in Banach spaces, *Nonlinear Anal.* **61**, 1031-1039, (2005).
2. W.R. Mann, Mean value methods in iteration, *Proc. Amer. Math. Soc.* **4**, 506-510, (1953).
3. S. Ishikawa, Fixed points and iteration of a nonexpansive mapping in a Banach space, *Proc. Amer. Math. Soc.* **59**, 65-71, (1976).
4. S. Ishikawa, Fixed point by a new iteration, *Proc. Amer. Math. Soc.* **44**, 147-150, (1974).

5. K.K. Tan and H.K. Xu, Approximating fixed points of nonexpansive mappings by the Ishikawa iteration process, *J. Math. Anal. Appl.* **178**, 301–308, (1993).
6. L.C. Zeng, A note on approximating fixed points of nonexpansive mappings by the Ishikawa iteration process, *J. Math. Anal. Appl.* **226**, 245–250, (1998).
7. M. Aslam Noor, New approximation schemes for general variational inequalities, *J. Math. Anal. Appl.* **251**, 217–229, (2000).
8. J.S. Jung and S.S. Kim, Strong convergence theorems for nonexpansive nonself-mappings in Banach spaces, *Nonlinear Anal.* **33**, 321–329, (1998).
9. H. Zhou, R.P. Agarwal, Y.J. Cho and Y.S. Kim, Nonexpansive mappings and iterative methods in uniformly convex Banach spaces, *G. M. J.* **9**, 591–600, (2002).
10. Y.J. Cho, H.Y. Zhou and G. Guo, Weak and strong convergence theorems for three-step iterations with errors for asymptotically nonexpansive mappings, *Computers Math. Applic.* **47**, 707–717, (2004).
11. S. Suantai, Weak and strong convergence criteria of Noor iterations for asymptotically nonexpansive mappings, *J. Math. Anal. Appl.* **311**, 506–517, (2005).
12. Z. Opial, Weak convergence of successive approximations for nonexpansive mappings, *Bull. Amer. Math. Soc.* **73**, 591–597, (1967).
13. K. Nammanee, M. Aslam Noor and S. Suantai, Convergence criteria of modified Noor iterations with errors for asymptotically nonexpansive mappings, *J. Math. Anal. Appl.* **314**, 320–334, (2006).
14. F.E. Browder, Semicontractive and semiaccretive nonlinear mappings in Banach spaces, *Bull. Amer. Math. Soc.* **74**, 660–665, (1968).



# The modified Noor iterations with errors for non-Lipschitzian mappings in Banach spaces

Kamonrat Nammanee<sup>a</sup>, Suthep Suantai<sup>b,\*</sup>

<sup>a</sup> Department of Mathematics, Faculty of Science, Naresuan University, Phitsanulok 65000, Thailand

<sup>b</sup> Department of Mathematics, Faculty of Science, Chiang Mai University, Chiang Mai 50200, Thailand

## Abstract

In this paper, weak and strong convergence theorems are established for the modified Noor iterations for asymptotically nonexpansive mappings in the intermediate sense in a uniformly convex Banach space. Mann and Ishikawa-type iterations are included by the modified Noor iterations with errors. The results obtained in this paper improve the recent ones announced by Schu [J. Schu, Iterative construction of fixed points of asymptotically nonexpansive mappings, *J. Math. Anal. Appl.* 158 (1991) 407–413; J. Schu, Weak and strong convergence to fixed points of asymptotically nonexpansive mappings, *Bull. Austral. Math. Soc.* 43 (1991) 153–159], Xu and Noor [B.L. Xu, M.A. Noor, Fixed point iterations for asymptotically nonexpansive mappings in Banach spaces, *J. Math. Anal. Appl.* 267 (2002) 1–10], et al. [Y.J. Cho, H.Y. Zhou, G. Guo, Weak and strong convergence theorems for three-step iterations with errors for asymptotically nonexpansive mappings, *Comput. Math. Appl.* 47 (2004) 707–717], Suantai [S. Suantai, Weak and strong convergence criteria of Noor Iterations for asymptotically nonexpansive mappings, *J. Math. Anal. Appl.* 311 (2005) 1–10], Nammanee et al. [K. Nammanee, M.A. Noor, S. Suantai, Convergence criteria of modified Noor iterations for asymptotically nonexpansive mappings, *J. Math. Anal. Appl.* 314 (2006) 320–334], and many others.

© 2006 Elsevier Inc. All rights reserved.

**Keywords:** Asymptotically nonexpansive mapping in the intermediate sense; Completely continuous; Modified Noor iteration; Mann condition; Uniformly convex Banach space

## 1. Introduction

The concept of asymptotically nonexpansiveness was introduced by Goebel and Kirk [7] in 1972. Noor [8,9] have introduced the three-step iterations and studied the approximate solutions of set-valued inclusion and variational inequalities in Hilbert spaces. Glowinski and Le Tallec [10] used three-step iterative schemes to find the approximate solutions of the elastoviscoplasticity problem, liquid crystal flow, and eigenvalue computation. It has been shown in [10] that the three-step iterative schemes give better results than the two-step and one-step approximate iterations. In 1998, Haubruge et al. [11] introduced

\* Corresponding author.

E-mail addresses: [kamonratn@nu.ac.th](mailto:kamonratn@nu.ac.th) (K. Nammanee), [scmti005@chiangmai.ac.th](mailto:scmti005@chiangmai.ac.th) (S. Suantai).

<sup>1</sup> Supported by Thailand Research Fund.

convergence analysis of the three-step schemes of Glowinski and Le Tallec [10] and applied these schemes to obtain new spitting-type algorithms for solving variation inequalities, separable convex programming and minimization of a sum of convex functions. They also prove that three-step iterations lead to highly parallelized algorithms under certain conditions. Thus we conclude that three-step schemes play an important and significant part in solving various problems, which arise in pure and applied science.

The concept of asymptotically nonexpansive in the intermediate sense was introduced by Bruck et al. [12]. This concept is a generalization of asymptotically nonexpansiveness. Let  $C$  be a subset of real normed linear space  $X$ , and let  $T$  be a self-mapping on  $C$ . The fixed point set of  $T$ ,  $F(T)$ , is defined by  $F(T) = \{x \in C : Tx = x\}$ .  $T$  is said to be nonexpansive provided  $\|Tx - Ty\| \leq \|x - y\|$  for all  $x, y \in C$ ;  $T$  is called asymptotically nonexpansive if there exists a sequence  $\{k_n\}$ ,  $k_n \geq 1$  with  $\lim_{n \rightarrow \infty} k_n = 1$ , such that

$$\|T^n x - T^n y\| \leq k_n \|x - y\|$$

for all  $x, y \in C$  and each  $n \geq 1$ .

$T$  is called asymptotically nonexpansive in the intermediate sense [12] provided  $T$  is uniformly continuous and

$$\limsup_{n \rightarrow \infty} \sup_{x, y \in C} (\|T^n x - T^n y\| - \|x - y\|) \leq 0.$$

It is known [13] that if  $X$  is a uniformly convex Banach space and  $T$  is asymptotically nonexpansive in the intermediate sense, then  $F(T) \neq \emptyset$ .

The modified Noor iterations with errors is defined as follows.

Let  $X$  be a normed space,  $C$  be a nonempty subset of  $X$ , and  $T : C \rightarrow C$  be a given mapping. Then for a given  $x_0 \in C$ , compute the sequence  $\{x_n\}$ ,  $\{y_n\}$  and  $\{z_n\}$  by the iterative schemes

$$\begin{aligned} z_n &= a_n T^n x_n + (1 - a_n - \gamma_n) x_n + \gamma_n u_n, \\ y_n &= b_n T^n z_n + c_n T^n x_n + (1 - b_n - c_n - \mu_n) x_n + \mu_n v_n, \\ z_{n+1} &= \alpha_n T^n y_n + \beta_n T^n z_n + (1 - \alpha_n - \beta_n - \lambda_n) x_n + \lambda_n w_n, \quad n \geq 1, \end{aligned} \quad (1.1)$$

where  $\{a_n\}$ ,  $\{b_n\}$ ,  $\{c_n\}$ ,  $\{\alpha_n\}$ ,  $\{\beta_n\}$ ,  $\{\gamma_n\}$ ,  $\{\mu_n\}$ ,  $\{\lambda_n\}$  are appropriate sequences in  $[0, 1]$  and  $\{u_n\}$ ,  $\{v_n\}$  and  $\{w_n\}$  are bounded sequences in  $C$ .

The iterative schemes (1.1) are called the modified Noor iterations with errors. Noor iterations include the Mann-Ishikawa iterations as special cases. If  $\gamma_n = \mu_n = \lambda_n \equiv 0$ , then (1.1) reduces to the modified Noor iterations defined by Suantai [5]

$$\begin{aligned} z_n &= a_n T^n x_n + (1 - a_n) x_n, \\ y_n &= b_n T^n z_n + c_n T^n x_n + (1 - b_n - c_n) x_n, \\ z_{n+1} &= \alpha_n T^n y_n + \beta_n T^n z_n + (1 - \alpha_n - \beta_n) x_n, \quad n \geq 1, \end{aligned} \quad (1.2)$$

where  $\{a_n\}$ ,  $\{b_n\}$ ,  $\{c_n\}$ ,  $\{\alpha_n\}$ ,  $\{\beta_n\}$  are appropriate sequences in  $[0, 1]$ .

We note that the usual Ishikawa and Mann iterations are special cases of (1.1) and if  $a_n = b_n = \gamma_n = \mu_n = \lambda_n \equiv 0$ , then (1.1) reduces to the Noor iterations defined by Xu and Noor [3]

$$\begin{aligned} z_n &= a_n T^n x_n + (1 - a_n) x_n, \\ y_n &= b_n T^n z_n + (1 - b_n) x_n, \\ z_{n+1} &= \alpha_n T^n y_n + (1 - \alpha_n) x_n, \quad n \geq 1, \end{aligned} \quad (1.3)$$

where  $\{a_n\}$ ,  $\{b_n\}$ ,  $\{\alpha_n\}$  are appropriate sequences in  $[0, 1]$ .

For  $a_n = c_n = \beta_n = \gamma_n = \mu_n = \lambda_n \equiv 0$ , then (1.1) reduces to the usual Ishikawa iterative schemes

$$\begin{aligned} y_n &= b_n T^n x_n + (1 - b_n) x_n, \\ z_{n+1} &= \alpha_n T^n y_n + (1 - \alpha_n) x_n, \quad n \geq 1, \end{aligned} \quad (1.4)$$

where  $\{b_n\}$ ,  $\{\alpha_n\}$  are appropriate sequences in  $[0, 1]$ .

If  $a_n = b_n = c_n = \beta_n = \gamma_n = \mu_n = \lambda_n \equiv 0$ , then (1.1) reduces to the usual Mann iterative scheme

$$z_{n+1} = \alpha_n T^n x_n + (1 - \alpha_n) x_n, \quad n \geq 1. \quad (1.5)$$

where  $\{\alpha_n\}$  are appropriate sequences in  $[0, 1]$ . See [1,2] for more details about Mann iterative scheme.

The purpose of this paper is to establish several strong convergence theorems for the modified Noor iterations with errors (1.1) for completely continuous asymptotically nonexpansive mappings in the intermediate sense, and weak convergence theorems for asymptotically nonexpansive mappings in the intermediate sense in a uniformly convex Banach space with Opial's condition.

Recall that a Banach space  $X$  is said to satisfy Opial's condition [14] if  $x_n \rightarrow x$  weakly as  $n \rightarrow \infty$  implies that

$$\limsup_{n \rightarrow \infty} \|x_n - x\| < \limsup_{n \rightarrow \infty} \|x_n - y\|.$$

In the sequel, the following lemmas are needed to prove our main results.

**Lemma 1.1** [15, Lemma 1]. Let  $\{a_n\}$ ,  $\{b_n\}$  and  $\{\delta_n\}$  be sequences of nonnegative real numbers satisfying the inequality

$$a_{n+1} \leq (1 + \delta_n)a_n + b_n, \quad \forall n = 1, 2, \dots$$

If  $\sum_{n=1}^{\infty} \delta_n < \infty$  and  $\sum_{n=1}^{\infty} b_n < \infty$ , then

- (1)  $\lim_{n \rightarrow \infty} a_n$  exists.
- (2)  $\lim_{n \rightarrow \infty} a_n = 0$  whenever  $\liminf_{n \rightarrow \infty} a_n = 0$ .

**Lemma 1.2** [4, Lemma 1.6]. Let  $X$  be a uniformly convex Banach space,  $C$  a nonempty closed convex subset of  $X$ , and  $T: C \rightarrow C$  be an asymptotically nonexpansive mapping. Then  $I - T$  is demiclosed at 0, i.e., if  $x_n \rightarrow x$  weakly and  $x_n - Tx_n \rightarrow 0$  strongly, then  $x \in F(T)$ , where  $F(T)$  is the set of fixed point of  $T$ .

**Lemma 1.3** [5, Lemma 2.7]. Let  $X$  be a Banach space which satisfies Opial's condition and let  $\{x_n\}$  be a sequence in  $X$ . Let  $u, v \in X$  be such that  $\lim_{n \rightarrow \infty} \|x_n - u\|$  and  $\lim_{n \rightarrow \infty} \|x_n - v\|$  exist. If  $\{x_{n_k}\}$  and  $\{x_{m_k}\}$  are subsequences of  $\{x_n\}$  which converge weakly to  $u$  and  $v$ , respectively, then  $u = v$ .

**Lemma 1.4** [4, Lemma 1.4]. Let  $X$  be a uniformly convex Banach space and  $B_r = \{x \in X : \|x\| \leq r\}$ ,  $r > 0$ . Then there exists a continuous, strictly increasing, and convex function  $g: [0, \infty) \rightarrow [0, \infty)$ ,  $g(0) = 0$  such that

$$\|\lambda x + \beta y + \gamma z\|^2 \leq \lambda \|x\|^2 + \beta \|y\|^2 + \gamma \|z\|^2 - \lambda \beta g(\|x - y\|)$$

for all  $x, y, z \in B_r$ , and all  $\lambda, \beta, \gamma \in [0, 1]$  with  $\lambda + \beta + \gamma = 1$ .

**Lemma 1.5** [6, Lemma 1.4]. Let  $X$  be a uniformly convex Banach space and  $B_r = \{x \in X : \|x\| \leq r\}$ ,  $r > 0$ . Then there exists a continuous, strictly increasing, and convex function  $g: [0, \infty) \rightarrow [0, \infty)$ ,  $g(0) = 0$  such that

$$\|\alpha x + \beta y + \mu z + \lambda w\|^2 \leq \alpha \|x\|^2 + \beta \|y\|^2 + \mu \|z\|^2 + \lambda \|w\|^2 - \alpha \beta g(\|x - y\|)$$

for all  $x, y, z, w \in B_r$ , and all  $\alpha, \beta, \mu, \lambda \in [0, 1]$  with  $\alpha + \beta + \mu + \lambda = 1$ .

## 2. Main results

In this section, we prove strong convergence theorems for the modified Noor iterations with errors (1.1) for asymptotically nonexpansive mapping in the intermediate sense in a uniformly convex Banach space. In order to prove our main results, the following lemmas are needed.

The next lemma is crucial for proving the main theorems.

**Lemma 2.1.** Let  $X$  be a uniformly convex Banach space, and let  $C$  be a nonempty bounded closed and convex subset of  $X$ . Let  $T: C \rightarrow C$  be an asymptotically nonexpansive mapping in the intermediate sense. Put

$$G_n = \sup_{x, y \in C} (\|T^n x - T^n y\| - \|x - y\|) \vee 0, \quad \forall n \geq 1,$$

so that  $\sum_{n=1}^{\infty} G_n < \infty$ . Let  $\{a_n\}$ ,  $\{b_n\}$ ,  $\{c_n\}$ ,  $\{\alpha_n\}$ ,  $\{\beta_n\}$ ,  $\{\gamma_n\}$ ,  $\{\mu_n\}$  and  $\{\lambda_n\}$  be real sequences in  $[0, 1]$  such that  $a_n + \gamma_n$ ,  $b_n + c_n + \mu_n$  and  $\alpha_n + \beta_n + \lambda_n$  are in  $[0, 1]$  for all  $n \geq 1$ , and  $\sum_{n=1}^{\infty} \gamma_n < \infty$ ,  $\sum_{n=1}^{\infty} \mu_n < \infty$ ,  $\sum_{n=1}^{\infty} \lambda_n < \infty$ .

and let  $\{u_n\}$ ,  $\{v_n\}$  and  $\{w_n\}$  be bounded sequences in  $C$ . For a given  $x_1 \in C$ , let  $\{x_n\}$ ,  $\{y_n\}$  and  $\{z_n\}$  be the sequences defined as in (1.1).

- (i) If  $p \in F(T)$  then  $\lim_{n \rightarrow \infty} \|x_n - p\|$  exists.  
 (ii) If  $0 < \liminf_{n \rightarrow \infty} \alpha_n \leq \limsup_{n \rightarrow \infty} (\alpha_n + \beta_n + \lambda_n) < 1$ , then  $\lim_{n \rightarrow \infty} \|T^n y_n - x_n\| = 0$ .  
 (iii) If  $0 < \liminf_{n \rightarrow \infty} \alpha_n$  and  $0 < \liminf_{n \rightarrow \infty} b_n \leq \limsup_{n \rightarrow \infty} (b_n + c_n + \mu_n) < 1$ , then  $\lim_{n \rightarrow \infty} \|T^n z_n - x_n\| = 0$ .  
 (iv) If  $0 < \liminf_{n \rightarrow \infty} b_n \leq \limsup_{n \rightarrow \infty} (b_n + c_n + \mu_n) < 1$  and  $0 < \liminf_{n \rightarrow \infty} \alpha_n \leq \limsup_{n \rightarrow \infty} (\alpha_n + \beta_n + \lambda_n) < 1$ , then  $\lim_{n \rightarrow \infty} \|T^n x_n - x_n\| = 0$ .

**Proof.** (i) By [13]  $F(T) \neq \emptyset$ . Let  $p \in F(T)$ . Since  $\{G_n\}$ ,  $\{u_n\}$ ,  $\{v_n\}$  and  $\{w_n\}$  are bounded sequences in  $C$ , we put

$$M = \sup_{n \geq 1} G_n \vee \sup_{n \geq 1} \|u_n - p\| \vee \sup_{n \geq 1} \|v_n - p\| \vee \sup_{n \geq 1} \|w_n - p\|.$$

For each  $n \geq 1$ , we note that

$$\begin{aligned} \|z_n - p\| &= \|a_n T^n x_n + (1 - a_n - \gamma_n)x_n + \gamma_n u_n - p\| \\ &\leq (1 - a_n - \gamma_n)\|x_n - p\| + a_n\|T^n x_n - p\| + \gamma_n\|u_n - p\| \\ &\leq a_n\|x_n - p\| + a_n G_n + (1 - a_n - \gamma_n)\|x_n - p\| + \gamma_n\|u_n - p\| \\ &\leq \|x_n - p\| + G_n + M\gamma_n, \end{aligned} \quad (2.1)$$

$$\begin{aligned} \|y_n - p\| &= \|b_n T^n z_n + c_n T^n x_n + (1 - b_n - c_n - \mu_n)x_n + \mu_n v_n - p\| \\ &\leq (1 - b_n - c_n - \mu_n)\|x_n - p\| + b_n\|T^n z_n - p\| \\ &\quad + c_n\|T^n x_n - p\| + \mu_n\|v_n - p\| \\ &\leq (1 - b_n - c_n - \mu_n)\|x_n - p\| + b_n[\|z_n - p\| + G_n] \\ &\quad + c_n[\|x_n - p\| + G_n] + \mu_n\|v_n - p\| \\ &\leq (1 - b_n - c_n - \mu_n)\|x_n - p\| + b_n[(\|x_n - p\| + G_n + M\gamma_n) + G_n] \\ &\quad + c_n[\|x_n - p\| + G_n] + M\mu_n \\ &\leq \|x_n - p\| + 3G_n + M\gamma_n + M\mu_n, \end{aligned} \quad (2.2)$$

$$\begin{aligned} \|x_{n+1} - p\| &= \|\alpha_n T^n y_n + \beta_n T^n z_n + (1 - \alpha_n - \beta_n - \lambda_n)x_n + \lambda_n w_n - p\| \\ &\leq (1 - \alpha_n - \beta_n - \lambda_n)\|x_n - p\| + \alpha_n\|T^n y_n - p\| \\ &\quad + \beta_n\|T^n z_n - p\| + \lambda_n\|w_n - p\| \\ &\leq (1 - \alpha_n - \beta_n - \lambda_n)\|x_n - p\| + \alpha_n[\|y_n - p\| + G_n] \\ &\quad + \beta_n[\|z_n - p\| + G_n] + \lambda_n\|w_n - p\| \\ &\leq (1 - \alpha_n - \beta_n - \lambda_n)\|x_n - p\| \\ &\quad + \alpha_n[(\|x_n - p\| + 3G_n + M\gamma_n + M\mu_n) + G_n] \\ &\quad + \beta_n[(\|x_n - p\| + G_n + M\gamma_n) + G_n] + M\lambda_n \\ &\leq \|x_n - p\| + 6G_n + M\gamma_n + M\mu_n + M\lambda_n. \end{aligned} \quad (2.3)$$

Since  $\sum_{n=1}^{\infty} G_n < \infty$ ,  $\sum_{n=1}^{\infty} \gamma_n < \infty$ ,  $\sum_{n=1}^{\infty} \mu_n < \infty$ , and  $\sum_{n=1}^{\infty} \lambda_n < \infty$ , it follows from Lemma 1.1 that  $\lim_{n \rightarrow \infty} \|x_n - p\|$  exists.

(ii) By [13],  $T$  has a fixed point  $p \in C$ . Choose a number  $r > 0$  such that  $C \subseteq B_r$  and  $C - C \subseteq B_r$ . By Lemma 1.4, there is a continuous, strictly increasing, and convex function  $g_1 : [0, \infty) \rightarrow [0, \infty)$ ,  $g_1(0) = 0$  such that

$$\|\lambda x + \beta y + \gamma z\|^2 \leq \lambda\|x\|^2 + \beta\|y\|^2 + \gamma\|z\|^2 - \lambda\beta g_1(\|x - y\|) \quad (2.4)$$

for all  $x, y, z \in B_r$ , and all  $\lambda, \beta, \gamma \in [0, 1]$  with  $\lambda + \beta + \gamma = 1$ .

It follows from (2.4) that

$$\begin{aligned}
 \|z_n - p\|^2 &= \|a_n T^n x_n + (1 - a_n - \gamma_n)x_n + \gamma_n u_n - p\|^2 \\
 &= \|a_n(T^n x_n - p) + (1 - a_n - \gamma_n)(x_n - p) + \gamma_n(u_n - p)\|^2 \\
 &\leq a_n \|T^n x_n - p\|^2 + (1 - a_n - \gamma_n) \|x_n - p\|^2 + \gamma_n \|u_n - p\|^2 - a_n(1 - a_n - \gamma_n)g_1(\|T^n x_n - x_n\|) \\
 &\leq a_n[\|x_n - p\| + G_n]^2 + (1 - a_n - \gamma_n) \|x_n - p\|^2 + \gamma_n \|u_n - p\|^2 - a_n(1 - a_n - \gamma_n)g_1(\|T^n x_n - x_n\|) \\
 &= a_n[\|x_n - p\|^2 + 2G_n \|x_n - p\| + G_n^2] + (1 - a_n - \gamma_n) \|x_n - p\|^2 + \gamma_n \|u_n - p\|^2 \\
 &\quad - a_n(1 - a_n - \gamma_n)g_1(\|T^n x_n - x_n\|) \\
 &\leq \|x_n - p\|^2 + 2G_n \|x_n - p\| + G_n^2 + M^2 \gamma_n - a_n(1 - a_n - \gamma_n)g_1(\|T^n x_n - x_n\|).
 \end{aligned}$$

By Lemma 1.5, there exists a continuous strictly increasing convex function  $g_2: [0, \infty) \rightarrow [0, \infty)$  such that

$$\|\alpha x + \beta y + \mu z + \lambda w\|^2 \leq \alpha \|x\|^2 + \beta \|y\|^2 + \mu \|z\|^2 + \lambda \|w\|^2 - \alpha \beta g_2(\|x - y\|)$$

for all  $x, y, z, w \in B_r$ , and all  $\alpha, \beta, \mu, \lambda \in [0, 1]$  with  $\alpha + \beta + \mu + \lambda = 1$ . It follows from (2.6) that

$$\begin{aligned}
 \|y_n - p\|^2 &= \|b_n T^n z_n + c_n T^n x_n + (1 - b_n - c_n - \mu_n)x_n + \mu_n v_n - p\|^2 \\
 &= \|b_n(T^n z_n - p) + (1 - b_n - c_n - \mu_n)(x_n - p) + c_n(T^n x_n - p) + \mu_n(v_n - p)\|^2 \\
 &\leq (1 - b_n - c_n - \mu_n) \|x_n - p\|^2 + b_n \|T^n z_n - p\|^2 + c_n \|T^n x_n - p\|^2 \\
 &\quad + \mu_n \|v_n - p\|^2 - b_n(1 - b_n - c_n - \mu_n)g_2(\|T^n z_n - x_n\|) \\
 &\leq (1 - b_n - c_n - \mu_n) \|x_n - p\|^2 + b_n[\|z_n - p\| + G_n]^2 + c_n[\|x_n - p\| + G_n]^2 + \mu_n \|v_n - p\|^2 \\
 &\quad - b_n(1 - b_n - c_n - \mu_n)g_2(\|T^n z_n - x_n\|) \\
 &= (1 - b_n - c_n - \mu_n) \|x_n - p\|^2 + b_n[\|z_n - p\|^2 + 2G_n \|z_n - p\| + G_n^2] \\
 &\quad + c_n[\|x_n - p\|^2 + 2G_n \|x_n - p\| + G_n^2] + \mu_n \|v_n - p\|^2 \\
 &\quad - b_n(1 - b_n - c_n - \mu_n)g_2(\|T^n z_n - x_n\|) \\
 &\leq (1 - b_n - c_n - \mu_n) \|x_n - p\|^2 + b_n(\|x_n - p\|^2 + 2G_n \|x_n - p\| + G_n^2 + M^2 \gamma_n) \\
 &\quad + 2G_n(\|x_n - p\| + G_n + M \gamma_n) + G_n^2 + c_n[\|x_n - p\|^2 + 2G_n \|x_n - p\| + G_n^2] + M^2 \mu_n \\
 &\quad - b_n(1 - b_n - c_n - \mu_n)g_2(\|T^n z_n - x_n\|) \\
 &\leq \|x_n - p\|^2 + 6G_n \|x_n - p\| + 5G_n^2 + M^2(\gamma_n + \mu_n) + 2MG_n \\
 &\quad - b_n(1 - b_n - c_n - \mu_n)g_2(\|T^n z_n - x_n\|),
 \end{aligned}$$

and

$$\begin{aligned}
 \|x_{n+1} - p\|^2 &= \|\alpha_n T^n x_n + \beta_n T^n z_n + (1 - \alpha_n - \beta_n - \lambda_n)x_n + \lambda_n w_n - p\|^2 \\
 &= \|\alpha_n(T^n x_n - p) + (1 - \alpha_n - \beta_n - \lambda_n)(x_n - p) + \beta_n(T^n z_n - p) + \lambda_n(w_n - p)\|^2 \\
 &\leq (1 - \alpha_n - \beta_n - \lambda_n) \|x_n - p\|^2 + \alpha_n \|T^n x_n - p\|^2 + \beta_n \|T^n z_n - p\|^2 \\
 &\quad + \lambda_n \|w_n - p\|^2 - \alpha_n(1 - \alpha_n - \beta_n - \lambda_n)g_2(\|T^n x_n - x_n\|) \\
 &\leq (1 - \alpha_n - \beta_n - \lambda_n) \|x_n - p\|^2 \\
 &\quad + \alpha_n[\|y_n - p\| + G_n]^2 + \beta_n[\|z_n - p\| + G_n]^2 + \lambda_n \|w_n - p\|^2 \\
 &\quad - \alpha_n(1 - \alpha_n - \beta_n - \lambda_n)g_2(\|T^n x_n - x_n\|) \\
 &= (1 - \alpha_n - \beta_n - \lambda_n) \|x_n - p\|^2 + \alpha_n[\|y_n - p\|^2 + 2G_n \|y_n - p\| + G_n^2] \\
 &\quad + \beta_n[\|z_n - p\|^2 + 2G_n \|z_n - p\| + G_n^2] + \lambda_n \|w_n - p\|^2 \\
 &\quad - \alpha_n(1 - \alpha_n - \beta_n - \lambda_n)g_2(\|T^n x_n - x_n\|)
 \end{aligned}$$

$$\begin{aligned}
&\leq (1 - \alpha_n - \beta_n - \lambda_n) \|x_n - p\|^2 \\
&\quad + \alpha_n [\|x_n - p\|^2 + 6G_n \|x_n - p\| + 5G_n^2 + M^2(\gamma_n + \mu_n) + 2MG_n] \\
&\quad + 2G_n (\|x_n - p\| + 3G_n + M\gamma_n + M\mu_n) + G_n^2 \\
&\quad + \beta_n [\|x_n - p\|^2 + 2G_n \|x_n - p\| + G_n^2 + M^2\gamma_n] \\
&\quad + 2G_n (\|x_n - p\| + G_n + M\gamma_n) + G_n^2 + M^2\lambda_n \\
&\quad - \alpha_n (1 - \alpha_n - \beta_n - \lambda_n) g_2(\|T^n y_n - x_n\|) \\
&\leq \|x_n - p\|^2 + 12G_n \|x_n - p\| + 16G_n^2 + M^2(2\gamma_n + \mu_n) + 8MG_n \\
&\quad - \alpha_n (1 - \alpha_n - \beta_n - \gamma_n) g_2(\|T^n y_n - x_n\|),
\end{aligned} \tag{2.8}$$

which imply that

$$\alpha_n (1 - \alpha_n - \beta_n - \lambda) g_2(\|T^n y_n - x_n\|) \leq \|x_n - p\|^2 - \|x_{n+1} - p\|^2 + 12LG_n + 16G_n^2 + M^2(2\gamma_n + \mu_n) + 8MG_n, \tag{2.9}$$

and

$$\alpha_n b_n (1 - b_n - c_n - \mu_n) g_2(\|T^n z_n - x_n\|) \leq \|x_n - p\|^2 - \|x_{n+1} - p\|^2 + 12LG_n + 16G_n^2 + M^2(2\gamma_n + \mu_n) + 8MG_n, \tag{2.10}$$

where  $L = \sup\{\|x_n - p\| : n \geq 1\}$ .

If  $0 < \liminf_{n \rightarrow \infty} \alpha_n \leq \limsup_{n \rightarrow \infty} (\alpha_n + \beta_n + \lambda_n) < 1$ , then there exist a positive integer  $n_0$  and  $\eta, \eta' \in (0, 1)$  such that

$$0 < \eta < \alpha_n \text{ and } \alpha_n + \beta_n + \lambda_n < \eta' < 1 \text{ for all } n \geq n_0.$$

This implies by (2.9) that

$$\begin{aligned}
\eta(1 - \eta') g_2(\|T^n z_n - x_n\|) &\leq \|x_n - p\|^2 - \|x_{n+1} - p\|^2 + 12LG_n + 16G_n^2 + M^2(2\gamma_n + \mu_n) + 8MG_n \\
&\leq \|x_n - p\|^2 - \|x_{n+1} - p\|^2 + 12LG_n + 16MG_n + M^2(2\gamma_n + \mu_n) + 8MG_n \\
&\leq \|x_n - p\|^2 - \|x_{n+1} - p\|^2 + 12KG_n + 5KG_n + M^2(2\gamma_n + \mu_n) + 8KG_n \\
&= \|x_n - p\|^2 - \|x_{n+1} - p\|^2 + 17KG_n + M^2(2\gamma_n + \mu_n),
\end{aligned} \tag{2.11}$$

where  $K = \max\{M, L\}$ , for all  $n \geq n_0$ . It follows from (2.11) that for  $m \geq n_0$

$$\begin{aligned}
\sum_{n=n_0}^m g_2(\|T^n z_n - x_n\|) &\leq \frac{1}{\eta(1 - \eta')} \left( \sum_{n=n_0}^m (\|x_n - p\|^2 - \|x_{n+1} - p\|^2) + \sum_{n=n_0}^m (17KG_n + M^2(2\gamma_n + \mu_n)) \right) \\
&\leq \frac{1}{\eta(1 - \eta')} \left( \|x_{n_0} - p\|^2 + 17K \sum_{n=n_0}^m G_n + M^2 \sum_{n=n_0}^m (2\gamma_n + \mu_n) \right).
\end{aligned} \tag{2.12}$$

Since  $\sum_{n=1}^{\infty} G_n < \infty$ . Let  $m \rightarrow \infty$  in inequality (2.12) we get that  $\sum_{n=n_0}^{\infty} g_2(\|T^n z_n - x_n\|) < \infty$ , and therefore  $\lim_{n \rightarrow \infty} g_2(\|T^n z_n - x_n\|) = 0$ . Since  $g$  is strictly increasing and continuous at 0 with  $g(0) = 0$ , it follows that  $\lim_{n \rightarrow \infty} \|T^n z_n - x_n\| = 0$ .

(iv) If  $0 < \liminf_{n \rightarrow \infty} \alpha_n$  and  $0 < \liminf_{n \rightarrow \infty} b_n \leq \limsup_{n \rightarrow \infty} (b_n + c_n + \mu_n) < 1$ , then by the using a similar method together with inequality (2.10), it can be shown that

$$\lim_{n \rightarrow \infty} \|T^n y_n - x_n\| = 0.$$

(v) If  $0 < \liminf_{n \rightarrow \infty} b_n \leq \limsup_{n \rightarrow \infty} (b_n + c_n + \mu_n) < 1$  and  $0 < \liminf_{n \rightarrow \infty} \alpha_n \leq \limsup_{n \rightarrow \infty} (\alpha_n + \beta_n + \lambda_n) < 1$ , then by (iv) and (iii) we have

$$\lim_{n \rightarrow \infty} \|T^n z_n - x_n\| = 0 \text{ and } \lim_{n \rightarrow \infty} \|T^n y_n - x_n\| = 0. \tag{2.13}$$

From  $y_n = (1 - b_n - c_n - \mu_n)x_n + b_nT^n z_n + c_nT^n x_n + \mu_n v_n$ , we have

$$\begin{aligned}\|y_n - x_n\| &= \|(1 - b_n - c_n - \mu_n)x_n + b_nT^n z_n + c_nT^n x_n + \mu_n v_n - x_n\| \\ &= \|b_n(T^n z_n - x_n) + c_nT^n(x_n - x_n) + \mu_n(v_n - x_n)\| \\ &\leq b_n\|T^n z_n - x_n\| + c_n\|T^n x_n - x_n\| + \mu_n\|x_n - v_n\|.\end{aligned}$$

Thus

$$\begin{aligned}\|T^n x_n - x_n\| &= \|T^n x_n - T^n y_n + T^n y_n - x_n\| \leq \|T^n x_n - T^n y_n\| + \|T^n y_n - x_n\| \\ &\leq \|x_n - y_n\| + G_n + \|T^n y_n - x_n\| \\ &\leq b_n\|T^n z_n - x_n\| + c_n\|T^n x_n - x_n\| + \mu_n\|x_n - v_n\| + G_n + \|T^n y_n - x_n\|,\end{aligned}$$

and so

$$(1 - c_n)\|T^n x_n - x_n\| \leq b_n\|T^n z_n - x_n\| + \mu_n\|x_n - v_n\| + G_n + \|T^n y_n - x_n\|.$$

Since  $\limsup_{n \rightarrow \infty} c_n < 1$ , it follows from (2.13) and  $\sum_{n=1}^{\infty} G_n < \infty$  that

$$\lim_{n \rightarrow \infty} \|T^n x_n - x_n\| = 0. \quad \square$$

**Theorem 2.2.** Let  $X$  be a uniformly convex Banach space, and let  $C$  be a nonempty bounded closed and convex subset of  $X$ . Let  $T$  be a completely continuous asymptotically nonexpansive in the intermediate sense. Put

$$G_n = \sup_{x, y \in C} (\|T^n x - T^n y\| - \|x - y\|) \vee 0 \quad \forall n \geq 1,$$

so that  $\sum_{n=1}^{\infty} G_n < \infty$ . Let  $\{a_n\}$ ,  $\{b_n\}$ ,  $\{c_n\}$ ,  $\{\alpha_n\}$ ,  $\{\beta_n\}$ ,  $\{\gamma_n\}$ ,  $\{\mu_n\}$  and  $\{\lambda_n\}$  be real sequences in  $[0, 1]$  such that  $a_n + \gamma_n$ ,  $b_n + c_n + \mu_n$  and  $\alpha_n + \beta_n + \lambda_n$  are in  $[0, 1]$  for all  $n \geq 1$ , and  $\sum_{n=1}^{\infty} \gamma_n < \infty$ ,  $\sum_{n=1}^{\infty} \mu_n < \infty$ ,  $\sum_{n=1}^{\infty} \lambda_n < \infty$  and let  $\{u_n\}$ ,  $\{v_n\}$  and  $\{w_n\}$  be bounded sequences in  $C$ . For a given  $x_1 \in C$ , let  $\{x_n\}$ ,  $\{y_n\}$  and  $\{z_n\}$  be sequences defined as in (1.1) and

- (i)  $0 < \liminf_{n \rightarrow \infty} \alpha_n \leq \limsup_{n \rightarrow \infty} (\alpha_n + \beta_n + \lambda_n) < 1$  and
- (ii)  $0 < \liminf_{n \rightarrow \infty} b_n \leq \limsup_{n \rightarrow \infty} (b_n + c_n + \mu_n) < 1$ .

Then  $\{x_n\}$ ,  $\{y_n\}$  and  $\{z_n\}$  converge strongly to a fixed point of  $T$ .

**Proof.** By Lemma 2.1, we have

$$\begin{aligned}\lim_{n \rightarrow \infty} \|T^n y_n - x_n\| &= 0, \\ \lim_{n \rightarrow \infty} \|T^n z_n - x_n\| &= 0, \\ \lim_{n \rightarrow \infty} \|T^n x_n - x_n\| &= 0.\end{aligned}$$

It follows from (2.14) that  $\lim_{n \rightarrow \infty} \|y_n - x_n\| = 0$ .

From  $x_{n+1} = (1 - \alpha_n - \beta_n - \lambda_n)x_n + \alpha_n T^n y_n + \beta_n T^n z_n + \lambda_n w_n$ , we have

$$\begin{aligned}\|x_{n+1} - x_n\| &= \|(1 - \alpha_n - \beta_n - \lambda_n)x_n + \alpha_n T^n y_n + \beta_n T^n z_n + \lambda_n w_n - x_n\| \\ &\leq \alpha_n\|T^n y_n - x_n\| + \beta_n\|T^n z_n - x_n\| + \lambda_n\|w_n - x_n\| \rightarrow 0.\end{aligned}$$

And

$$\begin{aligned}\|x_{n+1} - T^n x_{n+1}\| &\leq \|x_{n+1} - x_n\| + \|T^n x_{n+1} - T^n x_n\| + \|T^n x_n - x_n\| \\ &\leq \|x_{n+1} - x_n\| + \|x_{n+1} - x_n\| + G_n + \|T^n x_n - x_n\| \rightarrow 0.\end{aligned}$$

Since

$$\|x_{n+1} - Tx_{n+1}\| \leq \|x_{n+1} - T^{n+1}x_{n+1}\| + \|Tx_{n+1} - T^{n+1}x_{n+1}\|$$

and by uniform continuity of  $T$  and  $\lim_{n \rightarrow \infty} \|T^n x_n - x_n\| = 0$ , it follows that  $\lim_{n \rightarrow \infty} \|x_n - Tx_n\| = 0$ .

Since  $T$  is completely continuous and  $\{x_n\} \subseteq C$  is bounded, there exists a subsequence  $\{x_{n_k}\}$  of  $\{x_n\}$  such that  $\{Tx_{n_k}\}$  converges. Therefore from  $\lim_{n \rightarrow \infty} \|x_n - Tx_n\| = 0$ ,  $\{x_{n_k}\}$  converges. Let  $\lim_{k \rightarrow \infty} x_{n_k} = p$ . By continuity of  $T$  and  $\lim_{n \rightarrow \infty} \|x_n - Tx_n\| = 0$ , we have that  $Tp = p$ , so  $p$  is a fixed point of  $T$ . By Lemma 2.1 (i),  $\lim_{n \rightarrow \infty} \|x_n - p\|$  exists. But  $\lim_{k \rightarrow \infty} \|x_{n_k} - p\| = 0$ . Thus  $\lim_{n \rightarrow \infty} \|x_n - p\| = 0$ . Since  $\|y_n - x_n\| \rightarrow 0$  as  $n \rightarrow \infty$ , and

$$\|z_n - x_n\| = \|a_n T^n x_n + (1 - a_n - \gamma_n)x_n + \gamma_n u_n - x_n\| \leq \|T^n x_n - x_n\| + \gamma_n \|u_n - x_n\| \rightarrow 0 \quad \text{as } n \rightarrow \infty,$$

it follows that  $\lim_{n \rightarrow \infty} y_n = p$  and  $\lim_{n \rightarrow \infty} z_n = p$ .  $\square$

From Theorem 2.2, we have the following results.

**Corollary 2.3** [6, Theorem 2.3]. *Let  $X$  be a uniformly convex Banach space, and  $C$  a nonempty bounded, closed and convex subset of  $X$ . Let  $T$  be a completely continuous asymptotically nonexpansive self-map of  $C$  with  $\{k_n\}$  satisfying  $k_n \geq 1$  and  $\sum_{n=1}^{\infty} (k_n - 1) < \infty$ . Let  $\{a_n\}$ ,  $\{b_n\}$ ,  $\{c_n\}$ ,  $\{\alpha_n\}$ ,  $\{\beta_n\}$ ,  $\{\gamma_n\}$ ,  $\{\mu_n\}$  and  $\{\lambda_n\}$  be sequences of real numbers in  $[0, 1]$  with  $b_n + c_n + \mu_n \in [0, 1]$  and  $\alpha_n + \beta_n + \lambda_n \in [0, 1]$  for all  $n \geq 1$ , and  $\sum_{n=1}^{\infty} \gamma_n < \infty$ ,  $\sum_{n=1}^{\infty} \mu_n < \infty$ ,  $\sum_{n=1}^{\infty} \lambda_n < \infty$  and*

- (i)  $0 < \liminf_{n \rightarrow \infty} b_n \leq \limsup_{n \rightarrow \infty} (b_n + c_n + \mu_n) < 1$ , and
- (ii)  $0 < \liminf_{n \rightarrow \infty} \alpha_n \leq \limsup_{n \rightarrow \infty} (\alpha_n + \beta_n + \lambda_n) < 1$ .

*Let  $\{x_n\}$ ,  $\{y_n\}$  and  $\{z_n\}$  be the sequences defined by the modified Noor iterations with errors (1.1). Then  $\{x_n\}$ ,  $\{y_n\}$  and  $\{z_n\}$  converge strongly to a fixed point of  $T$ .*

**Corollary 2.4** [5, Theorem 2.3]. *Let  $X$  be a uniformly convex Banach space, and  $C$  a nonempty bounded, closed and convex subset of  $X$ . Let  $T$  be a completely continuous asymptotically nonexpansive self-map of  $C$  with  $\{k_n\}$  satisfying  $k_n \geq 1$  and  $\sum_{n=1}^{\infty} (k_n - 1) < \infty$ . Let  $\{a_n\}$ ,  $\{b_n\}$ ,  $\{c_n\}$ ,  $\{\alpha_n\}$ ,  $\{\beta_n\}$  be sequences of real numbers in  $[0, 1]$  with  $b_n + c_n \in [0, 1]$  and  $\alpha_n + \beta_n \in [0, 1]$  for all  $n \geq 1$ , and*

- (i)  $0 < \liminf_{n \rightarrow \infty} b_n \leq \limsup_{n \rightarrow \infty} (b_n + c_n) < 1$ , and
- (ii)  $0 < \liminf_{n \rightarrow \infty} \alpha_n \leq \limsup_{n \rightarrow \infty} (\alpha_n + \beta_n) < 1$ .

*Let  $\{x_n\}$ ,  $\{y_n\}$  and  $\{z_n\}$  be the sequences defined by the three-step iterative scheme (1.2). Then  $\{x_n\}$ ,  $\{y_n\}$  and  $\{z_n\}$  converge strongly to a fixed point of  $T$ .*

For  $c_n = \beta_n \equiv 0$  in Theorem 2.2, we obtain the following result.

**Corollary 2.5** [3, Theorem 2.1]. *Let  $X$  be a uniformly convex Banach space, and let  $C$  be a bounded, closed and convex subset of  $X$ . Let  $T$  be a completely continuous asymptotically nonexpansive self-map of  $C$  with  $\{k_n\}$  satisfying  $k_n \geq 1$  and  $\sum_{n=1}^{\infty} (k_n - 1) < \infty$ . Let  $\{a_n\}$ ,  $\{b_n\}$ ,  $\{\alpha_n\}$  be real sequences in  $[0, 1]$  satisfying*

- (i)  $0 < \liminf_{n \rightarrow \infty} b_n \leq \limsup_{n \rightarrow \infty} b_n < 1$ , and
- (ii)  $0 < \liminf_{n \rightarrow \infty} \alpha_n \leq \limsup_{n \rightarrow \infty} \alpha_n < 1$ .

*For a given  $x_1 \in C$ , define*

$$\begin{aligned} z_n &= a_n T^n x_n + (1 - a_n)x_n, \\ y_n &= b_n T^n z_n + (1 - b_n)x_n, \quad n \geq 1, \\ x_{n+1} &= \alpha_n T^n y_n + (1 - \alpha_n)x_n. \end{aligned}$$

*Then  $\{x_n\}$ ,  $\{y_n\}$  and  $\{z_n\}$  converge strongly to a fixed point of  $T$ .*

When  $a_n = c_n = \beta_n \equiv 0$  in Theorem 2.2, we can obtain Ishikawa-type convergence result.

**Corollary 2.6.** Let  $X$  be a uniformly convex Banach space, and let  $C$  be a bounded, closed and convex subset of  $X$ . Let  $T$  be a completely continuous asymptotically nonexpansive self-map of  $C$  with  $\{k_n\}$  satisfying  $\sum_{n=1}^{\infty} (k_n - 1) < \infty$ . Let  $\{b_n\}, \{\alpha_n\}$  be a real sequence in  $[0, 1]$  satisfying

- (i)  $0 < \liminf_{n \rightarrow \infty} b_n \leq \limsup_{n \rightarrow \infty} b_n < 1$ , and
- (ii)  $0 < \liminf_{n \rightarrow \infty} \alpha_n \leq \limsup_{n \rightarrow \infty} \alpha_n < 1$ .

For a given  $x_1 \in C$ , define

$$\begin{aligned} y_n &= b_n T^n z_n + (1 - b_n)x_n, \\ x_{n+1} &= \alpha_n T^n y_n + (1 - \alpha_n)x_n, \quad n \geq 1. \end{aligned}$$

Then  $\{x_n\}$  and  $\{y_n\}$  converge strongly to a fixed point of  $T$ .

In the next result, we prove weak convergence of the modified Noor iterations with errors for an asymptotically nonexpansive mapping in a uniformly convex Banach space satisfying Opial's condition.

**Theorem 2.7.** Let  $X$  be a uniformly convex Banach space which satisfies Opial's condition, and let  $C$  be a nonempty bounded, closed and convex subset of  $X$ . Let  $T$  be an asymptotically nonexpansive in the intermediate sense. Put

$$G_n = \sup_{x, y \in C} (\|T^n x - T^n y\| - \|x - y\|) \vee 0, \quad \forall n \geq 1,$$

so that  $\sum_{n=1}^{\infty} G_n < \infty$ . For a given  $x_1 \in C$ , let  $\{x_n\}, \{y_n\}$  and  $\{z_n\}$  be the sequences defined as in (1.1).

- (i)  $0 < \liminf_{n \rightarrow \infty} \alpha_n \leq \limsup_{n \rightarrow \infty} (\alpha_n + \beta_n + \lambda_n) < 1$  and
- (ii)  $0 < \liminf_{n \rightarrow \infty} b_n \leq \limsup_{n \rightarrow \infty} (b_n + c_n + \mu_n) < 1$ .

Then  $\{x_n\}$  converges weakly to a fixed point of  $T$ .

**Proof.** It follows from Theorem 2.2 that  $\lim_{n \rightarrow \infty} \|Tx_n - x_n\| = 0$ . Since  $X$  is uniformly convex and  $C$  is bounded, we may assume that  $x_n \rightarrow u$  weakly as  $n \rightarrow \infty$ , without loss of generality. By Lemma 2.1,  $u \in F(T)$ . Suppose that subsequences  $\{x_{n_k}\}$  and  $\{x_{m_k}\}$  of  $\{x_n\}$  converge weakly to  $u$  and  $v$ , respectively. By Lemma 1.2,  $u, v \in F(T)$ . By Lemma 2.1 (i),  $\lim_{n \rightarrow \infty} \|x_n - u\|$  and  $\lim_{n \rightarrow \infty} \|x_n - v\|$  exist. It follows from Lemma 1.3 that  $u = v$ . Therefore  $\{x_n\}$  converges weakly to a fixed point of  $T$ .  $\square$

From Theorem 2.7, we have the following results.

**Corollary 2.8** [6, Theorem 2.8]. Let  $X$  be a uniformly convex Banach space which satisfies Opial's condition, and let  $C$  be a nonempty closed, bounded and convex subset of  $X$ . Let  $T$  be an asymptotically nonexpansive self-map of  $C$  with  $\{k_n\}$  satisfying  $k_n \geq 1$  and  $\sum_{n=1}^{\infty} (k_n - 1) < \infty$ . Let  $\{a_n\}, \{b_n\}, \{c_n\}, \{\alpha_n\}, \{\beta_n\}, \{\mu_n\}, \{\lambda_n\}$  be sequences of real numbers in  $[0, 1]$  with  $a_n + \gamma_n, b_n + c_n + \mu_n$  and  $\alpha_n + \beta_n + \lambda_n$  are in  $[0, 1]$  for all  $n \geq 1$ , and  $\sum_{n=1}^{\infty} \mu_n < \infty, \sum_{n=1}^{\infty} \lambda_n < \infty$  and

- (i)  $0 < \liminf_{n \rightarrow \infty} b_n \leq \limsup_{n \rightarrow \infty} (b_n + c_n + \mu_n) < 1$ , and
- (ii)  $0 < \liminf_{n \rightarrow \infty} \alpha_n \leq \limsup_{n \rightarrow \infty} (\alpha_n + \beta_n + \lambda_n) < 1$ .

Let  $\{x_n\}$  be the sequence defined by modified Noor iterations with errors (1.1). Then  $\{x_n\}$  converges weakly to a fixed point of  $T$ .

**Corollary 2.9** [5, Theorem 2.3]. Let  $X$  be a uniformly convex Banach space which satisfies Opial's condition, and let  $C$  be a nonempty bounded, closed and convex subset of  $X$ . Let  $T$  be an asymptotically nonexpansive self-map of  $C$  with  $\{k_n\}$  satisfying  $k_n \geq 1$  and  $\sum_{n=1}^{\infty} (k_n - 1) < \infty$ . Let  $\{a_n\}, \{b_n\}, \{c_n\}, \{\alpha_n\}, \{\beta_n\}$  be sequences of real numbers in  $[0, 1]$  with  $b_n + c_n \in [0, 1]$  and  $\alpha_n + \beta_n \in [0, 1]$  for all  $n \geq 1$ , and

- (ii)  $0 < \liminf_{n \rightarrow \infty} b_n \leq \limsup_{n \rightarrow \infty} (b_n + c_n) < 1$ , and  
 (iii)  $0 < \liminf_{n \rightarrow \infty} \alpha_n \leq \limsup_{n \rightarrow \infty} (\alpha_n + \beta_n) < 1$ .

Let  $\{x_n\}$  be the sequence defined by three-step iterative scheme (1.2). Then  $\{x_n\}$  converges weakly to a fixed point of  $T$ .

When  $c_n = \beta_n \equiv 0$  in Theorem 2.7, we obtain the following result.

**Corollary 2.10.** Let  $X$  be a uniformly convex Banach space which satisfies Opial's condition, and  $C$  a nonempty bounded, closed and convex subset of  $X$ . Let  $T$  be an asymptotically nonexpansive self-map of  $C$  with  $\{k_n\}$  satisfying  $k_n \geq 1$  and  $\sum_{n=1}^{\infty} (k_n - 1) < \infty$ . Let  $\{a_n\}$ ,  $\{b_n\}$ ,  $\{\alpha_n\}$  be sequences of real numbers in  $[0, 1]$  and

- (i)  $0 < \liminf_{n \rightarrow \infty} b_n \leq \limsup_{n \rightarrow \infty} b_n < 1$ , and  
 (ii)  $0 < \liminf_{n \rightarrow \infty} \alpha_n \leq \limsup_{n \rightarrow \infty} \alpha_n < 1$ .

Let  $\{x_n\}$ ,  $\{y_n\}$  and  $\{z_n\}$  be the sequences defined by

$$\begin{aligned} z_n &= a_n T^n x_n + (1 - a_n) x_n, \\ y_n &= b_n T^n z_n + (1 - b_n) x_n, \quad n \geq 1, \\ x_{n+1} &= \alpha_n T^n y_n + (1 - \alpha_n) x_n. \end{aligned}$$

Then  $\{x_n\}$  converges weakly to a fixed point of  $T$ .

When  $a_n = c_n = \beta_n \equiv 0$  in Theorem 2.7, we obtain Ishikawa-type weak convergence theorem as follows:

**Corollary 2.11.** Let  $X$  be a uniformly convex Banach space which satisfies Opial's condition, and  $C$  a nonempty bounded, closed and convex subset of  $X$ . Let  $T$  be an asymptotically nonexpansive self-map of  $C$  with  $\{k_n\}$  satisfying  $k_n \geq 1$  and  $\sum_{n=1}^{\infty} (k_n - 1) < \infty$ . Let  $\{b_n\}$ ,  $\{\alpha_n\}$  be sequences of real numbers in  $[0, 1]$  such that

- (i)  $0 < \liminf_{n \rightarrow \infty} b_n \leq \limsup_{n \rightarrow \infty} b_n < 1$ , and  
 (ii)  $0 < \liminf_{n \rightarrow \infty} \alpha_n \leq \limsup_{n \rightarrow \infty} \alpha_n < 1$ .

Let  $\{x_n\}$  and  $\{y_n\}$  be the sequences defined by

$$\begin{aligned} y_n &= b_n T^n x_n + (1 - b_n) x_n, \\ x_{n+1} &= \alpha_n T^n y_n + (1 - \alpha_n) x_n, \quad n \geq 1. \end{aligned}$$

Then  $\{x_n\}$  converges weakly to a fixed point of  $T$ .

## Acknowledgement

The author would like to thank the Thailand Research Fund for their financial support.

## References

- [1] J. Schu, Iterative construction of fixed points of asymptotically nonexpansive mappings, *J. Math. Anal. Appl.* 158 (1991) 407–413.
- [2] J. Schu, Weak and strong convergence to fixed points of asymptotically nonexpansive mappings, *Bull. Austral. Math. Soc.* 43 (1991) 151–159.
- [3] T. Xu, M.A. Noor, Fixed point iterations for asymptotically nonexpansive mappings in Banach spaces, *J. Math. Anal. Appl.* 267 (2002) 444–453.
- [4] H.Y. Zhou, G. Guo, Weak and strong convergence theorems for three-step iterations with errors for asymptotically nonexpansive mappings, *Comput. Math. Appl.* 47 (2004) 707–717.
- [5] S. Suantai, Weak and strong convergence criteria of Noor Iterations for asymptotically nonexpansive mappings, *J. Math. Anal. Appl.* 311 (2005) 506–517.
- [6] K. Nammanee, M.A. Noor, S. Suantai, Convergence criteria of modified Noor iterations with errors for asymptotically nonexpansive mappings, *J. Math. Anal. Appl.* 314 (2006) 320–334.

- [7] K. Goebel, W.A. Kirk, A fixed point theorem for asymptotically nonexpansive mappings, *Proc. Amer. Math. Soc.* 35 (1972) 171–175.
- [8] M.A. Noor, New approximation schemes for general variational inequalities, *J. Math. Anal. Appl.* 251 (2000) 217–229.
- [9] M.A. Noor, Three-step iterative algorithms for multivalued quasi-variational inclusions, *J. Math. Anal. Appl.* 255 (2001) 101–113.
- [10] R. Glowinski, P. Le Tallec, *Augmented Lagrangian and Operator-splitting Methods in Nonlinear Mechanics*, SIAM, Philadelphia, 1989.
- [11] S. Haubruge, V.H. Nguyen, J.J. Strodiot, Convergence analysis and applications of the Glowinski–Le Tallec splitting scheme for finding a zero of the sum of two maximal monotone operators, *J. Optim. Theory Appl.* 97 (1998) 645–673.
- [12] R.E. Bruck, T. Kuczumow, S. Reich, Convergence of iterates of asymptotically nonexpansive mappings in Banach spaces with uniform Opial property, *Colloq. Math.* 65 (1993) 169–179.
- [13] W.A. Kirk, Fixed point theorems for non-Lipschitzian mappings of asymptotically nonexpansive type, *Israel J. Math.* 15 (1974) 346.
- [14] Z. Opial, Weak convergence of successive approximations for nonexpansive mappings, *Bull. Amer. Math. Soc.* 73 (1967) 293–297.
- [15] K.K. Tan, H.K. Xu, Approximating fixed points of nonexpansive mapping by the Ishikawa iteration process, *J. Math. Anal. Appl.* 178 (1993) 301–308.

## The criteria of strict monotonicity and rotundity points in generalized Calderón–Lozanovskiĭ spaces<sup>☆</sup>

Narin Petrot<sup>a,\*</sup>, Suthep Suantai<sup>b</sup>

<sup>a</sup> Department of Mathematics, Faculty of Science, Naresuan University, Phitsanulok, 65000, Thailand

<sup>b</sup> Department of Mathematics, Faculty of Science, Chiang Mai University, Chiang Mai, 50200, Thailand

Received 20 December 2007; accepted 28 February 2008

**Abstract** In this paper, some basic properties of the general modular space are proven. Criteria for strictly monotone points, extreme points and rotundity points in generalized Calderón–Lozanovskiĭ spaces are obtained. Consequently, the sufficient and necessary conditions for convexity properties of such spaces are given.

© 2008 Elsevier Ltd. All rights reserved.

**MSC:** 46B20; 46B30; 46C05; 46E30

**Keywords:** Orlicz–Orlicz function; Generalized Calderón–Lozanovskiĭ spaces; Point of lower(upper) monotonicity; Extreme point; *SU*-point; Rotundity point

### 1. Introduction

Throughout the paper  $\mathbb{R}$ ,  $\mathbb{R}^+$  and  $\mathbb{N}$  denote the sets of reals, nonnegative reals and natural numbers, respectively. Let  $X$  be a vector space, a function  $\rho : X \rightarrow [0, \infty]$  is called a *modular* if it satisfies the following conditions:

- (M1)  $\rho(0) = 0$  and  $x = 0$  whenever  $\rho(\lambda x) = 0$  for any  $\lambda > 0$ ;
- (M2)  $\rho(\alpha x) = \rho(x)$  for all scalar  $\alpha$  with  $|\alpha| = 1$ ;
- (M3)  $\rho(\alpha x + \beta y) \leq \rho(x) + \rho(y)$ , for all  $x, y \in X$  and all  $\alpha, \beta \geq 0$  with  $\alpha + \beta = 1$ .

Replacing (M3) by

- (M3')  $\rho(\alpha x + \beta y) \leq \alpha\rho(x) + \beta\rho(y)$ , for all  $x, y \in X$  and all  $\alpha, \beta \geq 0$  with  $\alpha + \beta = 1$ ,

the modular  $\rho$  is called *convex modular*. Moreover, for arbitrary  $x \in X$  we define

$$\rho(x) := \inf \left\{ \lambda > 0 : \rho\left(\frac{x}{\lambda}\right) < \infty \right\}.$$

When  $\rho(x) = \infty$  by the definition.

<sup>☆</sup> This research was supported by the Thailand Research Fund (Project No. MRG4980167).

\* Corresponding author.

E-mail addresses: [narinp@nu.ac.th](mailto:narinp@nu.ac.th) (N. Petrot), [scmti005@chiangmai.ac.th](mailto:scmti005@chiangmai.ac.th) (S. Suantai).

For any modular  $\rho$  on  $X$ , the space

$$X_\rho = \{x \in X : \rho(\lambda x) \rightarrow 0 \text{ as } \lambda \rightarrow 0^+\},$$

is called the modular space. If  $\rho$  is a convex modular, the functional

$$\|x\|_\rho = \inf \left\{ \lambda > 0 : \rho\left(\frac{x}{\lambda}\right) \leq 1 \right\},$$

is a norm on  $X_\rho$ , which is called the Luxemburg norm (see [35]). A modular  $\rho$  is called right-continuous (left-continuous) [continuous] if  $\lim_{\lambda \rightarrow 1^+} \rho(\lambda x) = \rho(x)$  for all  $x \in X_\rho$  ( $\lim_{\lambda \rightarrow 1^-} \rho(\lambda x) = \rho(x)$  for all  $x \in X_\rho$ ) [it is both right- and left-continuous].

**Remark 1.1.** If  $\rho$  is a convex modular and  $\rho(\lambda_0 x) < \infty$  for some  $x \in X_\rho$  and  $\lambda_0 > 0$ , then  $\rho$  is right-continuous at  $\lambda x$  for any  $\lambda \in [0, \lambda_0]$  and left-continuous at  $\lambda x$  for any  $\lambda \in (0, \lambda_0]$ . Indeed, this follows from the fact that the function  $f(t) = \rho(tx)$  is convex on  $\mathbb{R}^+$  and has finite values on the interval  $[0, \lambda_0]$  so it is a continuous function on  $[0, \lambda_0]$ .

A triple  $(T, \Sigma, \mu)$  stands for a nonatomic, positive, complete and  $\sigma$ -finite measure space, while  $L^0 = L^0(T, \Sigma, \mu)$  denotes the space of all (equivalence classes of)  $\sigma$ -measurable functions  $x : T \rightarrow \mathbb{R}$ . In what follows we will identify measurable functions which differ only on a set of measure zero. For  $x, y \in L^0$ , we write  $x \leq y$  if  $x(t) \leq y(t)$  for  $\mu$ -a.e.  $t \in T$  and the notion  $x < y$  is used for  $x \leq y$  and  $x \neq y$ . Moreover, for any  $x \in L^0$ , we denote by  $|x|$  the absolute value of  $x$ , i.e.  $|x|(t) = |x(t)|$  for  $\mu$ -a.e.  $t \in T$ .

By  $E$  we denote a Köthe space over the measure space  $(T, \Sigma, \mu)$ , i.e.  $E \subset L^0$  which satisfies the following conditions:

- (i) if  $x \in E$ ,  $y \in L^0$  and  $|y| \leq |x|$  for  $\mu$ -a.e. then  $y \in E$  and  $\|y\|_E \leq \|x\|_E$ ,
- (ii) there exists a function  $x$  in  $E$  which is strictly positive on the whole  $T$ .

A function  $\varphi : T \times \mathbb{R} \rightarrow [0, \infty)$  is said to be a Musielak–Orlicz function if  $\varphi(t, \cdot)$  is a nonzero function, it vanishes at zero, it is convex and even for  $\mu$ -a.e.  $t \in T$  and  $\varphi(\cdot, u)$  as well as  $\varphi^{-1}(\cdot, u)$  are  $\Sigma$ -measurable functions for any  $u \in \mathbb{R}^+$ , where  $\varphi^{-1}(t, \cdot)$  is the generalized inverse function of  $\varphi(t, \cdot)$  defined on  $[0, \infty)$  by

$$\varphi^{-1}(t, u) = \inf\{v \geq 0 : \varphi(t, v) > u\}$$

for each  $t \in T$  (see [35]). For Musielak–Orlicz function  $\varphi$  we define a measurable function with respect to  $t \in T$  by

$$a(t) = \sup\{u \geq 0 : \varphi(t, u) = 0\},$$

see [6, page 175].

**Remark 1.2.** Let  $\varphi : T \times \mathbb{R} \rightarrow [0, \infty)$  be a Musielak–Orlicz function. Then

- (i)  $\varphi^{-1}(t, \cdot)$  vanishes only at zero;
- (ii)  $\varphi(t, \varphi^{-1}(t, u)) = u$  for all  $u \in [0, \infty)$  and

$$\varphi^{-1}(t, \varphi(t, u)) = \begin{cases} 0, & \text{if } u \in [0, a(t)], \\ u, & \text{if } u \in (a(t), \infty); \end{cases}$$

for  $\mu$ -a.e.  $t \in T$ .

Given any Musielak–Orlicz function  $\varphi$ , we define on  $L^0$  a convex modular  $\varrho_\varphi$  by

$$\varrho_\varphi(x) = \begin{cases} \|\varphi \circ x\|_E & \text{if } \varphi \circ x \in E, \\ \infty & \text{otherwise;} \end{cases}$$

and the generalized Calderón–Lozanovskii space is defined by

$$E_\varphi = \{x \in L^0 : \varphi \circ \lambda x \in E \text{ for some } \lambda > 0\}.$$

Then  $E_\varphi = (E_\varphi, \|\cdot\|_\varphi)$  becomes a normed space, where  $\|\cdot\|_\varphi$  denotes for the Luxemburg norm induced by  $\varrho_\varphi$  (see [4,9]).

For the investigations of generalized Calderón–Lozanovskii space we refer to [8–10,27].

In the case when  $\varphi$  is an Orlicz function, i.e. there is a set  $A \in \Sigma$  with  $\mu(A) = 0$  such that  $\varphi(t_1, \cdot) = \varphi(t_2, \cdot)$  for all  $t_1, t_2 \in T \setminus A$ , these Calderón–Lozanovskii spaces were investigated in [3,4,30] and the investigations were continued in the papers [5,11,15,17,20,26,28,29,32–34,36,37].

We say a Musielak–Orlicz function  $\varphi$  satisfies the condition  $\Delta_2^E$  if there exist a set  $A \in \Sigma$  with  $\mu(A) = 0$ , a constant  $K > 0$  and a nonnegative function  $h \in E$  such that the inequality

$$\varphi(t, 2u) \leq K\varphi(t, u) + h(t)$$

holds for all  $t \in T \setminus A$  and  $u \in \mathbb{R}$  (see [35] when  $E = L^1$  and [9] in general).

**Lemma 1.3 ([9, Lemma 5]).** The property that  $\|x\|_\varphi = 1$  if and only if  $\varrho_\varphi(x) = 1$  holds true for any  $x \in E_\varphi$  if and only if  $\varphi \in \Delta_2^E$ .

**Lemma 1.4 ([19, Lemma 1]).** For any Musielak–Orlicz function  $\varphi$  the inequality

$$\varphi(t, x+v) \geq \varphi(t, u) + \varphi(t, a(t) + v)$$

holds for  $\mu$ -a.e.  $t \in T$  and any  $u \geq a(t)$ ,  $v \geq 0$ .

**Lemma 1.5 ([9, Corollary 7]).** If  $\varphi \in \Delta_2^E$  then  $\mu(\{t \in T : a(t) > 0\}) = 0$ .

By  $S(E)$ ,  $B(E)$  and  $E^+ (= \{x \in E : x \geq 0\})$  we denote the unit sphere, the closed unit ball and the positive cone in the Köthe space  $E$ . For any  $x \in E$ , define  $\text{supp } x = \{t \in T : x(t) \neq 0\}$ .

A point  $x \in E^+$  is called a point of *upper monotonicity* (UM-point for short) if for every  $y \in E^+ \setminus \{0\}$  we have  $\|x\|_E < \|x+y\|_E$ . A point  $x \in E^+ \setminus \{0\}$  is called a point of *lower monotonicity* (LM-point for short) if for every  $y \in E^+ \setminus \{0\}$ , such that  $y < x$ , we have  $\|x-y\|_E < \|x\|_E$ . If every point of  $S(E^+)$  is a UM-point (or an LM-point), then we say that the space  $E$  is *strictly monotone*. It is easy to see that  $x \in E^+ \setminus \{0\}$  in any Köthe space  $E$  is a LM-point (LM-point) if and only if  $x/\|x\|$  is a UM-point (LM-point). Therefore, it is enough to formulate the criteria of monotonicity for points in  $S(E^+)$  only.

A point  $x \in S(E)$  is said to be an *extreme point* of  $B(E)$  ( $x \in \text{ext } B(E)$  for short) if for any  $y, z \in B(E)$  such that  $x = y+z$  we have  $y = z$ . If any point of  $S(E)$  is an extreme point of  $B(E)$ , we say that the space  $E$  is *rotund* (see [29]).

A point  $x \in S(E)$  is called a *strong U-point* (SU-point for short) of  $B(E)$  if for any  $y \in S(E)$  with  $\|x+y\|_E = 2$ , we have  $x = y$ . It is obvious that a Banach space  $E$  is rotund if and only if any  $x \in S(E)$  is an SU-point, but the notions of an extreme point and an SU-point are different (see [7]).

It is well known that rotundity properties of Banach spaces have applications in various branches of mathematics, namely, Fixed point Theory, Approximation Theory, Ergodic Theory, and many others. Moreover, if the focus of the study is Banach lattices, then there are strong relationships between rotundity properties and monotonicity properties (see [2,13,14,16,18,21,24,25]). Specially, in [17,20] the local rotundity and local monotonicity structures of a certain Banach lattice, namely Calderón–Lozanovskii spaces, were studied. The results of our paper will be a generalization of these excellent papers [17,20] by considering Orlicz function with parameter called Musielak–Orlicz function instead of Orlicz function. Of course, some ideas from those papers are also applied in our paper. However, because of the different properties among functions, in many parts of the proofs of our results new methods and techniques are designed.

Let us note that if  $E$  has the Fatou property, i.e. for any  $x \in L^0$  and  $(x_n)_{n=1}^\infty$  in  $E$  such that  $0 \leq x_n \nearrow x$   $\mu$ -a.e. and  $\sup_n \|x_n\|_E < \infty$  we have that  $x \in E$  and  $\|x\|_E = \lim_{n \rightarrow \infty} \|x_n\|_E$  (see [1,23,31]), then  $E_\varphi$  also has this property and moreover, the modular  $\varrho_\varphi$  is left-continuous (see [9, Theorem 12]). Consequently,  $E_\varphi$  is a Banach space. In the whole paper we will assume that  $E$  is a Köthe space with the Fatou property. Moreover, we will denote  $\varphi(t) = \varphi(t, x(t))$  for each  $t \in T$ .

The paper is organized as follows. In Section 2 we give some basic auxiliary results of general modular space and in Section 3 is devoted to the strictly monotone points of  $E_\varphi$ . We study rotundity points of  $E_\varphi$  in Section 4. Finally, in Section 5 we give a characterization of rotundity structure in  $E_\varphi$ .

## 2. Auxiliary lemmas

We start by proving some facts in any modular space.

**Lemma 2.1.** Let  $X_\rho$  be a modular space generated by a convex modular  $\rho$  and  $x, y \in B(X_\rho)$ . If  $\xi(x) < 1$  then  $\xi\left(\frac{x+y}{2}\right) < 1$ .

**Proof.** Since  $\xi(x) < 1$ , we take a real number  $a \in (\xi(x), 1)$  and put  $\varepsilon = \frac{1-a}{1+a}$ . Then  $\varepsilon > 0$  and  $\frac{(1+\varepsilon)a}{2} + \frac{1+\varepsilon}{2} = 1$ . Thus,

$$\begin{aligned} \rho\left((1+\varepsilon)\left(\frac{x+y}{2}\right)\right) &= \rho\left(\frac{1+\varepsilon}{2} \cdot x + \frac{1+\varepsilon}{2} \cdot y\right) \\ &= \rho\left(\frac{(1+\varepsilon)a}{2} \cdot \frac{x}{a} + \frac{1+\varepsilon}{2} \cdot y\right) \\ &\leq \frac{(1+\varepsilon)a}{2} \rho\left(\frac{x}{a}\right) + \frac{1+\varepsilon}{2} \rho(y) < \infty, \end{aligned}$$

which implies that  $\xi\left(\frac{x+y}{2}\right) < 1$ . This completes the proof.  $\square$

**Lemma 2.2.** Let  $X_\rho$  be the modular space generated by a convex modular  $\rho$  and  $x \in B(X_\rho)$  be such that  $\xi(x) < 1$ . If  $y$  is any element in  $B(X_\rho)$  satisfying  $\left\|\frac{x+y}{2}\right\|_\rho = 1$ , then  $\rho\left(\frac{x+y}{2}\right) = 1$ .

**Proof.** By  $\xi(x) < 1$  and Lemma 2.1, we have  $\xi\left(\frac{x+y}{2}\right) < 1$ . Put  $I = \left[0, \frac{1}{\xi\left(\frac{x+y}{2}\right)}\right)$  and define a function  $f: I \rightarrow \mathbb{R}$  by  $f(t) = \rho\left(t\frac{x+y}{2}\right)$ . Then  $f$  is a convex function and has finite values on  $I$ , which imply that  $f$  is a continuous function on  $I$ . Assuming that  $\rho\left(\frac{x+y}{2}\right) < 1$ , there exists a  $\lambda > 1$  such that  $\rho\left(\lambda\frac{x+y}{2}\right) < 1$  whence  $\left\|\frac{x+y}{2}\right\|_\rho \leq \frac{1}{\lambda} < 1$ , contradiction.  $\square$

We close this section by giving a basic result on the generalized Calderón–Lozanovskiĭ space as follows:

**Lemma 2.3.** For any  $x \in E_\varphi$  and any measurable partition  $\{T_i\}_{i=1}^n$  of  $T$  we have,

$$\xi(x) = \max_{1 \leq i \leq n} \{\xi(x \chi_{T_i})\}.$$

**Proof.** Put  $\alpha = \max_{1 \leq i \leq n} \{\xi(x \chi_{T_i})\}$ , then it is obvious that  $\alpha \leq \xi(x)$ . We now show that the converse inequality holds. If not, then a real number  $\beta \in (\alpha, \xi(x))$  can be found and consequently,

$$\varrho_\varphi\left(\frac{x}{\beta}\right) = \left\|\varphi\left(\frac{x}{\beta}\right)\right\|_E = \left\|\sum_{i=1}^n \varphi\left(\frac{x}{\beta} \chi_{T_i}\right)\right\|_E \leq \sum_{i=1}^n \left\|\varphi\left(\frac{x}{\beta} \chi_{T_i}\right)\right\|_E = \sum_{i=1}^n \varrho_\varphi\left(\frac{x}{\beta} \chi_{T_i}\right) < \infty,$$

which contradicts the definition of the number  $\xi(x)$ .  $\square$

## 3. Points of monotonicity in $E_\varphi$

In this section, we give some criteria for upper and lower monotonicity points in  $E_\varphi$ .

**Theorem 3.1.** A point  $x \in S(E_\varphi^+)$  is upper monotone if and only if

- (i)  $\varrho_\varphi(x) = 1$ ;
- (ii)  $\mu(\{t \in T : x(t) < a(t)\}) = 0$ ;
- (iii)  $\varphi \circ x$  is an upper monotone point of  $E$ .

**Proof.** Necessity. If condition (i) does not hold, then  $\varrho_\varphi(x) =: r < 1$ . Let  $D$  be a subset of  $A$  such that  $\mu(D) > 0$  and  $\varphi \circ x|_D$  is a nonnegative measurable function defined by

$$\varphi \circ x|_D = \varphi^{-1} \left( t, \frac{1-r}{\|\chi_D\|_E} \right) \chi_D(t).$$

Then  $\varphi \circ x|_D = \frac{1-r}{\|\chi_D\|_E} \chi_D$  which implies  $\varphi \circ u \in E$ , and moreover,

$$\|\varphi \circ x|_D\|_E = \left\| \frac{(1-r)}{\|\chi_D\|_E} \chi_D \right\|_E = 1-r.$$

Thus,  $\varphi \circ x|_D$  is not an upper monotone point. There exist a real number  $\lambda > 0$  and a measurable function  $y > 0$  with  $\text{supp } y = D$  satisfying

$$\varphi(t, x(t)) + y(t) \leq \varphi(t, x(t)) + \varphi(t, u(t)), \quad y(t) \leq \lambda$$

for a.e.  $t \in T$ . On the other hand, an ascending sequence  $(T_n)_{n=1}^\infty$  such that  $\bigcup_n T_n = T$  and  $\sup_{t \in T_n} \varphi(t, u) < \infty$  for every  $n \in \mathbb{N}$  and  $u \in \mathbb{R}^+$  can be found (see [22]), which allows us to obtain a nonnegative real number  $d_\lambda$  such that

$$d_\lambda = \sup\{\varphi(t, \lambda) : t \in D\}.$$

Consequently,  $\varphi \circ y \leq d_\lambda \chi_D$  which implies that  $y \in E_\varphi$ . Moreover,

$$\begin{aligned} \|\varphi \circ (x+y)\|_E &= \|\varphi \circ x \chi_{T \setminus D} + \varphi \circ (x+y) \chi_D\|_E \leq \|\varphi \circ x \chi_{T \setminus D} + \varphi \circ x \chi_D + \varphi \circ u\|_E \\ &= \|\varphi \circ x + \varphi \circ u\|_E \leq \|\varphi \circ x\|_E + \|\varphi \circ u\|_E = r + (1-r) = 1. \end{aligned}$$

Thus,  $\|x+y\|_\varphi \leq 1$  and therefore,  $x$  is not an upper monotone point.

Suppose that (ii) is not satisfied. Then the set  $A = \{t \in T : x(t) < a(t)\}$  has a positive measure. Let us define  $\varphi \circ (x+y) = \varphi \circ x \chi_{T \setminus A} + \varphi \circ (x+y) \chi_A$  for all  $t \in T$ . We see that  $y \in E_\varphi^+ \setminus \{0\}$  and

$$\begin{aligned} \|\varphi \circ (x+y)\|_E &= \|\varphi \circ x \chi_{T \setminus A} + \varphi \circ (x+y) \chi_A\|_E \\ &= \|\varphi \circ x \chi_{T \setminus A} + \varphi \circ a \chi_A\|_E \\ &= \|\varphi \circ x \chi_{T \setminus A}\|_E \leq \varrho_\varphi(x) \leq 1. \end{aligned}$$

Thus,  $\|x+y\|_\varphi \leq 1$ . But, since  $y \in E_\varphi^+ \setminus \{0\}$  the fact that  $\|x+y\|_\varphi \geq \|x\|_\varphi = 1$  is always true, we obtain  $\|x+y\|_\varphi = 1$ . This means that  $x$  is not an upper monotone point.

It remains to show the necessity of condition (iii). Let us assume that  $x \in S(E_\varphi^+)$  is an upper monotone point. Since the necessity of (i) has been proved, we may assume that  $\varphi \circ x \in S(E)$  and suppose that condition (iii) is not satisfied, then there exists  $y \in E^+ \setminus \{0\}$  such that  $\|\varphi \circ x + y\|_E = 1$ . Let us define  $z \in E_\varphi^+ \setminus \{0\}$  by  $z(t) = \varphi^{-1}(t, y(t))$  for all  $t \in T$ . Then there exists a nonnegative measurable function  $h$  such that  $\text{supp } h \subset \text{supp } z$  and

$$\varphi(t, x(t)) + h(t) \leq \varphi(t, x(t)) + \varphi(t, z(t)), \quad h(t) \leq \lambda$$

for a.e.  $t \in T$ . Thus  $h \in E_\varphi$  and

$$\|\varphi \circ (x+h)\|_E \leq \|\varphi \circ x + \varphi \circ z\|_E = \|\varphi \circ x + y\|_E = 1,$$

which implies that  $\|x+h\|_\varphi = 1$ . This contradicts the upper monotonicity of  $x$  and the proof is completed.

**Sufficiency.** Let  $x \in S(E_\varphi^+)$  and assume that conditions (i)–(iii) are satisfied. Let  $y \in E^+ \setminus \{0\}$  be given. In view of Lemma 3.4, condition (ii) gives

$$\varphi(t, x(t)) + y(t) \geq \varphi(t, x(t)) + \varphi(t, a(t) + y(t))$$

for a.e.  $t \in T$ . Since  $\mu(\{t \in T : \varphi(t, a(t) + y(t)) > 0\}) > 0$  and  $\varphi \circ x$  is an upper monotone point in  $E$ , we have

$$\|\varphi \circ (x+y)\|_E = \|\varphi \circ (x+y)\|_E \geq \|\varphi \circ x + \varphi \circ (a+y)\|_E > \|\varphi \circ x\|_E = \varrho_\varphi(x) = 1,$$

thus,  $\|x+y\|_\varphi > 1$ . This completes the proof.  $\square$

**Theorem 3.2.** A point  $x \in S(E_\varphi^+)$  is a lower monotone point if and only if

$$\varrho_\varphi(x) < 1.$$

- (ii)  $\mu(\{t \in \text{supp } x : x(t) \leq a(t)\}) = 0$ ;
- (iii)  $\varphi \circ x$  is a lower monotone point of  $E$ .

**Proof. Necessity.** Let  $x \in S(E^+)$  be a lower monotone point. Suppose that condition (i) is not satisfied, i.e.  $\xi(x) = 1$ . Take  $A, B \in \Sigma$ , both of positive measure, such that  $A \cap B = \emptyset$  and  $A \cup B = \text{supp } x$ . Thus by Lemma 2.3 we obtain  $\xi(x\chi_A) = 1$  or  $\xi(x\chi_B) = 1$ . Without loss of generality we may assume that  $\xi(x\chi_A) = 1$ , and it would be  $\xi(x - x\chi_B) = \xi(x\chi_A) = 1$ . This implies  $\|x - x\chi_B\|_\varphi \geq 1$ , a contradiction.

If condition (ii) does not hold, then the set  $A = \{t \in \text{supp } x : x(t) \leq a(t)\}$  has positive measure. By (i), the necessity of which has been already proved, we have  $\xi(x) < 1$ , and consequently  $\varrho_\varphi(x) = 1$  by Lemma 2.2. Define  $y(t) = x(t)\chi_A(t)$ , then we have  $0 < y < x$ , and

$$\varrho_\varphi(x - y) = \|\varphi \circ x\chi_{T \setminus A}\|_E = \|\varphi \circ x\|_E = \varrho_\varphi(x) = 1.$$

This implies that  $\|x - y\|_\varphi = 1$ , a contradiction.

Now we will show that condition (iii) holds. By (i), we have  $\varphi \circ x \in S(E)$ . Let us take  $y \in E$  such that  $0 < y < \varphi \circ x$  and choose a measurable function  $z$  such that  $0 < z < x$  with  $\varphi \circ x - y \leq \varphi \circ (x - z)$ . Since  $x$  is a lower monotone point, we have

$$\|\varphi \circ x - y\|_E \leq \|\varphi \circ (x - z)\|_E = \varrho_\varphi(x - z) \leq \|x - z\|_\varphi < 1.$$

This shows that  $\varphi \circ x$  is then a lower monotone point of  $E$ .

**Sufficiency.** Let  $x \in S(E^+)$ ,  $y \in E^+ \setminus \{0\}$  be such that  $y < x$  and conditions (i)–(iii) are satisfied. Obviously,  $y \subset \text{supp } x$  which together with condition (ii) imply that for  $z = \varphi \circ x - \varphi \circ (x - y)$  we have  $z > 0$ . Moreover, by condition (i), we have  $\varrho_\varphi(x) = 1$ . Since  $\varphi \circ x$  is a lower monotone point of  $E$  and  $z \leq \varphi \circ x$ , so

$$\varrho_\varphi(x - y) = \|\varphi \circ (x - y)\|_E = \|\varphi \circ x - z\|_E < \|\varphi \circ x\|_E = \varrho_\varphi(x) = 1.$$

Using Eq. (3.1) together with  $\xi(x - y) < 1$  (by condition (i)) and the continuity of  $\varrho_\varphi$ , in light of Lemma 2.2, we have  $\|x - y\|_\varphi < 1$ . This completes the proof.  $\square$

#### 4. Points of rotundity in $E_\varphi$

We will study the points of rotundity, such as extreme point and  $SU$ -point in this Section. We begin with the following definition:

A point  $x \in S(E^+)$  is said to be an extreme point of  $B(E^+)$  ( $x \in \text{ext} B(E^+)$  for short) if for any  $x, y \in S(E^+)$  such that  $x = (y + z)/2$ , we have  $y = z = x$ .

**Lemma 4.1** ([17, Lemma 4]). In any Köthe space  $E$ ,  $x \in S(E)$  is an extreme point of  $B(E)$  if and only if  $|x|$  is a UM-point of  $E$  and  $|x| \in \text{ext } B(E^+)$ .

**Theorem 4.2.** A point  $x \in S(E_\varphi)$  is an extreme point of  $B(E_\varphi)$  if and only if

- (i)  $\varrho_\varphi(x) = 1$ ;
- (ii)  $\mu(\{t \in T : |x(t)| < a(t)\}) = 0$ ;
- (iii)  $\varphi \circ |x|$  is a UM-point;
- (iv) if  $u, v \in S(E)$  satisfy  $\frac{u+v}{2} = \varphi \circ |x|$  then either

$$u = v \quad \text{or} \quad \varphi \circ \left( \frac{y + z}{2} \right) < \frac{1}{2}(\varphi \circ y + \varphi \circ z),$$

where  $y(t) = \varphi^{-1}(t, |u(t)|)$ ,  $z(t) = \varphi^{-1}(t, |v(t)|)$  for all  $t \in T$ .

**Proof. Sufficiency.** Assume that conditions (i)–(iv) are satisfied. Let  $x \in S(E_\varphi)$  and  $y, z \in B(E_\varphi)$  be such that  $2x = y + z$ . We shall show that  $y = z$ . First, we will show that

$$\varphi \circ |x|(t) = \varphi \circ \frac{|y + z|}{2}(t) = \varphi \circ \left[ \frac{|y| + |z|}{2} \right](t) = \frac{1}{2}[\varphi \circ |y|(t) + \varphi \circ |z|(t)]$$

□. Note that, we always have

$$\varphi \circ \frac{|y+z|}{2}(t) \leq \varphi \circ \left[ \frac{|y|+|z|}{2} \right](t) \leq \frac{1}{2} [\varphi \circ |y|(t) + \varphi \circ |z|(t)]$$

□. Let  $A = \{t \in T : \varphi \circ |x|(t) < \frac{1}{2} [\varphi \circ |y|(t) + \varphi \circ |z|(t)]\}$ . If  $\mu(A) > 0$  then by conditions (i) and (iii)

$$\begin{aligned} \varphi \circ |x| &= \|\varphi \circ |x|\|_E < \left\| \frac{1}{2} \varphi \circ |y| + \frac{1}{2} \varphi \circ |z| \right\|_E \\ &\leq \frac{1}{2} (\|\varphi \circ |y|\|_E + \|\varphi \circ |z|\|_E) \leq 1, \end{aligned}$$

□. Contradiction. Consequently, Eq. (4.1) holds.

□. Let  $t \in T : \varphi(t, \cdot)$  is a convex and even function. It is clear that  $\mu(T \setminus C_\varphi) = 0$ . Next for each  $t \in C_\varphi$  we have  $\hat{y}(t) = \varphi^{-1}(t, \varphi(t, |y(t)|))$  and  $\hat{z}(t) = \varphi^{-1}(t, (\varphi(t, |z(t)|)))$ . Using condition (ii) together with the right of Remark 1.2(ii), we have  $\hat{y}(t) = |y(t)|$  and  $\hat{z}(t) = |z(t)|$  for  $\mu$ -a.e.  $t \in C_\varphi$ . Consequently, by condition (iv) we conclude that  $\varphi \circ |y|(t) = \varphi \circ |z|(t)$  for  $\mu$ -a.e.  $t \in C_\varphi$ . We claim that  $|y| = |z|$ . Put  $B = \{t \in T : |y|(t) \neq |z|(t)\}$  and suppose that  $\mu(B) > 0$ . Thus, since  $\varphi(t, \cdot)$  is an injective function on the set  $C_\varphi$  we should have

$$|y(t)| \leq a(t) \quad \text{and} \quad |y(t)| \wedge |z(t)| < a(t) \quad (4.2)$$

□. So

$$= \frac{1}{2} [\varphi \circ |y|(t) + \varphi \circ |z|(t)] = 0$$

□. Combining this equation with Eq. (4.2) and the assumption that  $2x = y + z$  we obtain  $|x(t)| < |a(t)|$  for  $t \in B$ . This contradicts condition (ii). Hence, we have the claim. Finally, by condition (ii) and the fact that  $\varphi(t, \cdot)$  is a function on  $[a(t), \infty)$  for all  $t \in C_\varphi$ , in view of Eq. (4.1), we obtain that  $|y(t) + z(t)| = |y(t)| + |z(t)|$ . This together with  $|y(t)| = |z(t)|$  for  $\mu$ -a.e.  $t \in T$  implies that  $y = z$ .

□. Let  $x \in S(E_\varphi)$  be an extreme point of  $B(E_\varphi)$ . By, Lemma 4.1 we obtain that  $|x|$  is a *UM*-point in  $E_\varphi$ . Then, from 3.1 we have  $x(t) \geq a(t)$  for  $\mu$ -a.e.  $t \in T$ ,  $\varrho_\varphi(x) = 1$  and  $\varphi \circ x$  is an upper monotone point of  $E$ . It remains only to prove that if  $x \in \text{ext } B(E_\varphi)$  then condition (iv) holds. If not, there are  $u, v \in S(E)$  such

$$\varphi \circ \left[ \frac{y+z}{2} \right](t) = \frac{1}{2} [\varphi \circ y(t) + \varphi \circ z(t)] = \frac{u(t) + v(t)}{2} = \varphi \circ |x|(t),$$

□. where  $y(t), z(t)$  are defined in condition (iv). Clearly,  $y, z \in S(E_\varphi)$  with  $y \neq z$ . Consequently,  $|x| \notin \text{ext } B(E_\varphi)$ . Finally, Lemma 4.1 yields that  $x \notin \text{ext } B(E_\varphi)$ . □

□. A point  $x \in S(E^+)$  is called a *strong U-point* (an *SU-point* for short) of  $B(E^+)$  if for any  $y \in S(E^+)$  with  $x \neq y$ , we have  $x = y$ .

□. (see page 387)). If a point  $x \in S(E^+)$  is an *SU-point* of  $B(E^+)$ , then  $x$  is a *LM-point* of  $E$  and  $x$  is an *LM-point* of  $E$ .

□. (Lemma 7)). A point  $x \in S(E)$  is an *SU-point* of  $B(E)$  if and only if  $|x|$  is an *SU-point* of  $B(E^+)$ .

□. Let  $E$  be a strictly monotone Köthe space and  $x \in S(E_\varphi)$ . Then  $x$  is an *SU-point* of  $B(E_\varphi)$  if and only

□.  $\mu(\{t \in T : |x|(t) \leq a(t)\}) = 0$ ;

□.  $E^+$  satisfies  $\|u + \varphi \circ |x|\|_E = 2$  then either

$$x = u + |x| \quad \text{or} \quad \varphi \circ \left( \frac{|x| + y}{2} \right) < \frac{1}{2} (\varphi \circ |x| + \varphi \circ y),$$

$$= \varphi^{-1}(t, u(t)) \text{ for all } t \in T.$$

**Proof. Necessity.** Assume that  $x$  is an  $SU$ -point of  $B(E_\varphi)$ . Applying Lemma 4.4, Remark 4.3 and Theorem 3.2 we see that the remainder is condition (iii). Suppose the converse, that is, there are  $u \in S(E^+)$  such that  $\|u + \varphi \circ |x|\|_E = 1$ ,  $u \neq \varphi \circ |x|$  and  $\varphi \circ \left(\frac{|x|+y}{2}\right) = \frac{1}{2}[\varphi \circ |x| + \varphi \circ y]$ , where  $y(t)$  is defined as in condition (iii). Then,

$$\varrho_\varphi(y) = \|\varphi \circ y\|_E = \|u\|_E = 1,$$

and consequently,

$$\begin{aligned} 2 &= \|u + \varphi \circ |x|\|_E = \|\varphi \circ y + \varphi \circ |x|\|_E \\ &\leq \|\varphi \circ y\|_E + \|\varphi \circ |x|\|_E \\ &\leq \varrho_\varphi(y) + \varrho_\varphi(x) \leq 2. \end{aligned}$$

This implies that

$$\begin{aligned} \varrho_\varphi\left(\frac{|x|+y}{2}\right) &= \left\|\varphi \circ \left(\frac{x+y}{2}\right)\right\|_E \\ &= \frac{1}{2} [\|\varphi \circ |x| + \varphi \circ y\|_E] \\ &= \frac{1}{2} [\|\varphi \circ |x|\|_E + \|\varphi \circ y\|_E] \\ &= \frac{1}{2} [\varrho_\varphi(|x|) + \varrho_\varphi(y)] = 1, \end{aligned}$$

so  $\left\|\frac{|x|+y}{2}\right\|_\varphi = 1$ . Since  $u \neq \varphi \circ |x|$ , we have  $|x| \neq y$ , which implies that  $|x|$  is not an  $SU$ -point of  $B(E_\varphi)$ . The

Lemma 4.4 finishes the proof of the necessity.

**Sufficiency.** Let  $y \in S(E_\varphi)$  be such that

$$\left\|\frac{x+y}{2}\right\|_\varphi = 1.$$

We shall show that  $x = y$ . Combining Eq. (4.3) with condition (i), and applying Lemma 2.2, we get  $\varrho_\varphi\left(\frac{x+y}{2}\right) = 1$ . This gives

$$\begin{aligned} 1 &= \varrho_\varphi\left(\frac{x+y}{2}\right) = \left\|\varphi \circ \left(\frac{x+y}{2}\right)\right\|_E \\ &\leq \frac{1}{2} \|\varphi \circ x + \varphi \circ y\|_E \\ &\leq \frac{1}{2} [\varrho_\varphi(x) + \varrho_\varphi(y)] \\ &\leq 1, \end{aligned}$$

whence

$$\|\varphi \circ x + \varphi \circ y\|_E = 2.$$

Using this equation together with the strict monotonicity of  $E$ , the fact  $\varrho_\varphi\left(\frac{x+y}{2}\right) = 1$  and the convexity of  $\|\cdot\|_E$  on  $\mathbb{R}$  for all  $t \in C_\varphi$ , where  $C_\varphi$  defined as in Theorem 4.2 it is easy to see that

$$\varphi \circ \left(\frac{|x|+|y|}{2}\right)(t) = \frac{\varphi \circ |x|(t) + \varphi \circ |y|(t)}{2}$$

for  $\mu$ -a.e.  $t \in C_\varphi$ . Put  $u(t) = \varphi \circ |y|(t)$  for all  $t \in T$ . Then  $u \in E^+$  and  $\|u\|_E = \|\varphi \circ y\|_E = \varrho_\varphi(y) = 1$  by Eq. (4.4). Moreover, by virtue of condition (iii), Eqs. (4.5) and (4.6) imply that  $\varphi \circ |x|(t) = \varphi \circ |y|(t)$  for  $\mu$ -a.e.  $t \in C_\varphi$ . Since  $\mu(\{t \in \text{supp } x : |x|(t) \leq a(t)\}) = 0$  and  $\varphi(t, \cdot)$  is an injective function on the interval  $[a(t), b(t)]$ ,  $\mu$ -a.e.  $t \in C_\varphi$  we get  $|x|(t) = |y|(t)$  for  $\mu$ -a.e.  $t \in T$ . Then  $|x+y| \leq |x| + |y| = 2|x|$ . If  $|x+y| < |x| + |y|$  then

$\varphi(|x|) + \varphi(|y|) < 1$  (since  $|x|$  is an  $LM$ -point of  $E_\varphi$  by Theorem 3.2). This contradicts Eq. (4.3) and proves that  $|x| = |y|$ . Combining this equality with  $|x| = |y|$ , we get  $x = y$ .  $\square$

#### Rotundity of $E_\varphi$

In this final section we present a result concerning the rotundity structure of  $E_\varphi$ .

**Theorem 5.1.** *Let  $E$  be a Köthe space and  $\varphi$  be a Musielak–Orlicz function. Then  $E_\varphi \in (R)$  if and only if*

(i)  $E \in (SM)$ ;

(ii)  $\varphi \in \Delta_2^E$ ;

(iii)  $\forall u, v \in S(E^+)$  with  $u \neq v$  then either

$$\left\| \frac{u+v}{2} \right\|_E < 1 \quad \text{or} \quad \varphi \circ \left( \frac{x+y}{2} \right) < \frac{1}{2}(\varphi \circ x + \varphi \circ y),$$

where  $x(t) = \varphi^{-1}(t, u(t))$  and  $y(t) = \varphi^{-1}(t, v(t))$  for all  $t \in T$ .

**Proof.** *Necessity.* Suppose on the contrary that  $E_\varphi \in (R)$  and  $E \notin (SM)$ . Then an element  $u \in S(E^+)$  which is not a  $UM$ -point can be found. Put  $x(t) = \varphi^{-1}(t, u(t))$ . Then  $\varrho_\varphi(x) = \|\varphi \circ x\|_E = \|u\|_E = 1$ , so  $x \in S(E_\varphi)$  and hence  $x \in \text{ext } B(E_\varphi)$ . However,  $\varphi \circ x$  is not a  $UM$ -point in  $E$ , thus Theorem 4.2 yields a contradiction.

Suppose that  $E_\varphi \in (R)$  and  $\varphi \notin \Delta_2^E$ . By Lemma 1.3, there exists  $x \in S(E_\varphi)$  with  $\varrho_\varphi(x) < 1$ . By  $E_\varphi \in (R)$ ,  $x \in \text{ext } B(E_\varphi)$  and Theorem 4.2 yields a contradiction.

Suppose that condition (iii) is not satisfied. Then there are  $u, v \in S(E^+)$  with  $u \neq v$  such that  $\|u+v\|_E = 2$  and  $\varphi \circ \left( \frac{x+y}{2} \right) = \frac{1}{2}(\varphi \circ x + \varphi \circ y) = \frac{u+v}{2}$ , where  $x(t), y(t)$  are defined in condition (iii). Putting  $z = \frac{x+y}{2}$ , we have  $\varphi \circ z = \frac{u+v}{2}$ , thus  $z \in \text{ext } B(E_\varphi)$ . Since  $x \in \text{ext } B(E_\varphi)$ , Theorem 4.2 yields a contradiction.

*Sufficiency.* Let  $x \in S(E_\varphi)$  be arbitrary. We shall show that  $x \in \text{ext } B(E_\varphi)$ , by proving that conditions (i)–(iv) in Theorem 4.2 are satisfied. First, by  $\varphi \in \Delta_2^E$  we have  $\varrho_\varphi(x) = 1$  and  $|x(t)| \geq a(t)$  for  $\mu$ -a.e.  $t \in T$  by Lemmas 1.3 and 1.2, respectively. Next,  $\varphi \circ |x|$  is a  $UM$ -point in  $E$ , because  $E \in (SM)$ . Finally, we will show that condition (iv) in Theorem 4.2 holds. Let  $u, v \in S(E)$  be such that  $\frac{u+v}{2} = \varphi \circ |x|$ . By condition (iii) in our assumptions, we get  $\varphi \circ \left( \frac{z+y}{2} \right) < \frac{1}{2}(\varphi \circ y + \varphi \circ z)$ , where  $\varphi \circ y = u$  and  $\varphi \circ z = v$ , which means that condition (iv) from Theorem 4.2 is satisfied. Hence, our theorem is proved.  $\square$

Observe that, if  $E = L^1$  then  $E_\varphi = \{x \in L^0 : \int_T \varphi(t, \lambda x(t)) d\mu < \infty \text{ for some } \lambda > 0\} =: L^\varphi$ , which is called the Musielak–Orlicz space. Therefore, a direct consequence of Theorem 5.1, we have the following result.

**Corollary 5.2.** *Let  $\varphi$  be a Musielak–Orlicz function and  $L^\varphi$  be the Musielak–Orlicz space generated by  $\varphi$ . Then  $L^\varphi \in (R)$  if and only if*

(i)  $L^1 \in (SM)$ ;

(ii)  $\forall u, v \in S(L_1^+)$  with  $u \neq v$  then

$$\varphi \circ \left( \frac{z+y}{2} \right) < \frac{1}{2}(\varphi \circ x + \varphi \circ y),$$

where  $x(t) = \varphi^{-1}(t, u(t))$  and  $y(t) = \varphi^{-1}(t, v(t))$  for all  $t \in T$ .

**Proof.** Since  $L^1 \in (SM)$  and for any  $u, v \in S(L_1^+)$  we must have  $\|\frac{u+v}{2}\|_{L_1} = 1$ , thus, the conclusion of Corollary 5.2 follows directly from Theorem 5.1. This completes the proof.  $\square$

**Theorem 5.3.** *Rotundity properties of Musielak–Orlicz space,  $L^\varphi$ , equipped with the Luxemburg norm were given in [12], in terms of the strict convexity of Musielak–Orlicz function  $\varphi$ . Since condition (ii) in Corollary 5.2 is that  $\varphi(t, \cdot)$  is a strictly convex Musielak–Orlicz function for  $\mu$ -a.e.  $t \in T$ , therefore, Corollary 5.2 gives a result*

## Acknowledgement

The authors are thankful to the referees for their valuable suggestions that helped to improve the presentation specially Lemma 2.2 and Theorem 5.1.

## References

- [1] C.D. Aliprantis, O. Burkinshaw, Positive operator, in: Pure and Applied Math., Academic Press Inc., 1985.
- [2] M.A. Akcoglu, L. Sucheston, On uniform monotonicity of norms and ergodic theorems in function spaces, *Re. Circ. Mat. Palermo* 2 (Suppl. 8) (1985) 325–335.
- [3] E.I. Bereznoi, M. Mastylo, On Calderón–Lozanovskii construction, *Bull. Pol. Acad. Sci. Math.* 37 (1989) 23–32.
- [4] A.P. Calderón, Intermediate spaces and interpolation, the complex method, *Studia Math.* 24 (1964) 113–190.
- [5] J. Cerdà, H. Hudzik, M. Mastylo, On the geometry of some Calderón–Lozanovskii interpolation spaces, *Indag. Math.* 6 (1) (1995) 35–40.
- [6] S. Chen, Geometry of Orlicz spaces, *Dissertationes Math.* 356 (1996).
- [7] Y. Cui, H. Hudzik, C. Meng, On some local geometry of Orlicz sequence spaces equipped with the Luxemburg norm, *Acta Math. Hungar.* 100 (1–2) (1998) 143–154.
- [8] F. Foralewski, On some geometric properties of generalized Calderón–Lozanovskii spaces, *Acta Math. Hungar.* 80 (1–2) (1998) 55–60.
- [9] P. Foralewski, H. Hudzik, Some basic properties of generalized Calderón–Lozanovskii spaces, *Collect. Math.* 48 (4–6) (1997) 523–530.
- [10] P. Foralewski, H. Hudzik, On some geometrical and topological properties of generalized Calderón–Lozanovskii sequence spaces, *Houston J. Math.* 25 (3) (1999) 523–542.
- [11] P. Foralewski, P. Kolwicz, Local uniform rotundity in Calderón–Lozanovskii spaces, *J. Convex Anal.* 14 (2) (2007) 395–412.
- [12] H. Hudzik, Strict convexity of Musielak–Orlicz spaces with Luxemburg’s norm, *Bull. Acad. Polon. Sci. Math.* 29 (5–6) (1981) 225–230.
- [13] H. Hudzik, Geometry of some classes of Banach function spaces, in: *Proceedings of the International Symposium on Banach and Function Spaces*, Yokohama Publisher, Kitakyushu, Japan, 2003, pp. 17–57.
- [14] H. Hudzik, A. Kamińska, Monotonicity properties of Lorentz spaces, *Proc. Amer. Math. Soc.* 123 (9) (1995) 2715–2721.
- [15] H. Hudzik, A. Kamińska, M. Mastylo, Geometric properties of some Calderón–Lozanovskii spaces and Orlicz–Lorentz spaces, *Houston J. Math.* 22 (1996) 639–663.
- [16] H. Hudzik, A. Kamińska, M. Mastylo, Monotonicity and rotundity properties in Banach lattices, *Rocky Mountain J. Math.* 30 (3) (2000) 933–950.
- [17] H. Hudzik, P. Kolwicz, A. Narloch, Local rotundity structure of Calderón–Lozanovskii spaces, *Indag. Math. (NS)* 17 (3) (2006) 375–380.
- [18] H. Hudzik, W. Kurc, Monotonicity properties of Musielak–Orlicz spaces and dominated best approximation in Banach lattices, *J. Approx. Theory* 95 (1998) 353–368.
- [19] H. Hudzik, X. Liu, T. Wang, Points of monotonicity in Musielak–Orlicz function spaces endowed with the Luxemburg norm, *Arch. Math.* 73 (2004) 534–545.
- [20] H. Hudzik, A. Narloch, Local monotonicity structure of Calderón–Lozanovskii spaces, *Indag. Math. (NS)* 15 (1) (2004) 1–12.
- [21] H. Hudzik, A. Narloch, Relationships between monotonicity and complex rotundity properties with some consequences, *Math. Slovaca* 55 (2005) 289–306.
- [22] A. Kamińska, Some convexity properties of Musielak–Orlicz spaces of Bochner type, *Rend. Circ. Mat. Palermo Suppl., Serie I* 30 (1996) 63–73.
- [23] L.V. Kantorovitch, G.P. Akilov, *Functional Analysis*, Nauka, Moscow, 1977 (in Russian).
- [24] W. Kurc, Strictly and uniformly monotone Musielak–Orlicz spaces and applications to best approximation, *J. Approx. Theory* 68 (2000) 173–187.
- [25] W. Kurc, Strictly and uniformly monotone sequential Musielak–Orlicz spaces, *Collect. Math.* 50 (1) (1999) 1–17.
- [26] P. Kolwicz, On property  $(\beta)$  in Banach lattices, Calderón–Lozanovskii and Orlicz–Lorentz spaces, *Proc. Indian Acad. Sci. (Math. Sci.)* 111 (2001) 319–336.
- [27] P. Kolwicz,  $P$ -convexity of Calderón–Lozanovskii spaces of Bochner type, *Acta Math. Hungar.* 91 (1–2) (2001) 115–130.
- [28] P. Kolwicz, Rotundity properties in Calderón–Lozanovskii spaces, *Houston J. Math.* 31 (3) (2005) 883–912.
- [29] P. Kolwicz, R. Pluciennik, On uniform rotundity in every direction in Calderón–Lozanovskii spaces, *J. Convex Anal.* 14 (3) (2007) 425–430.
- [30] G.Ya. Lozanovskii, A remark on an interpolation theorem of Calderón, *Funktsional. Anal. Prilozhen.* 6 (1972) 333–334.
- [31] J. Lindenstrauss, L. Tzafriri, *Classical Banach Spaces II*, Springer-Verlag, Berlin, Heidelberg, New York, 1979.
- [32] L. Maligranda, Calderón–Lozanovskii space and interpolation of operators, *Semesterbericht Functionalanalysis*, Tübingen 8 (1982) 100–104.
- [33] L. Maligranda, Orlicz Spaces and Interpolation, *Sem. Math.* 5 (1989) Campinas.
- [34] M. Mastylo, Interpolation of linear operators in Calderón–Lozanovskii spaces, *Comment. Math. Prace Mat.* 26 (1986) 247–256.
- [35] J. Musielak, Orlicz Spaces and Modular Spaces, in: *Lecture Notes in Math.*, vol. 1034, Springer, 1983.
- [36] Y. Raynaud, On duals of Calderón–Lozanovskii intermediate space, *Studia Math.* 124 (1997) 9–36.
- [37] Y. Raynaud, Ultrapowers of Calderón–Lozanovskii interpolation space, *Indag. Math. (NS)* 9 (1) (1998) 65–105.



Contents lists available at ScienceDirect

Nonlinear Analysis

journal homepage: [www.elsevier.com/locate/na](http://www.elsevier.com/locate/na)

# A new mapping for finding common solutions of equilibrium problems and fixed point problems of finite family of nonexpansive mappings

Arid Kangtunyakarn, Suthep Suantai\*

Department of Mathematics, Faculty of Science, Chiang Mai University, Chiang Mai, 50200, Thailand

## ARTICLE INFO

### Article history:

Received 16 December 2008

Accepted 3 March 2009

### Keywords:

Nonexpansive mappings

Strongly positive operator

Equilibrium problem

Viscosity approximation method

Fixed point

## ABSTRACT

In this paper, we introduce and study a new mapping generated by a finite family of nonexpansive mappings and finite real numbers and introduce a general iterative method concerning the new mappings for finding a common element of the set of solutions of an equilibrium problem and of the set of common fixed points of a finite family of nonexpansive mappings in a Hilbert space. Then, we prove a strong convergence theorem of the proposed iterative method for a finite family of nonexpansive mappings to the unique solution of variational inequality which is the optimality condition for a minimization problem. Our main result can be applied to obtain strong convergence of the general iterative methods which are modifications of those in [G. Marino, H.K. Xu, A general iterative method for nonexpansive mappings in Hilbert spaces, *J. Math. Anal. Appl.* 318 (1) (2006) 43–52; S. Plubtieng, R. Punpaeng, A general iterative method for equilibrium problems and fixed point problems in Hilbert spaces, *J. Math. Anal. Appl.* 336 (1) (2007) 455–469; S. Takahashi, W. Takahashi, Viscosity approximation methods for equilibrium problems and fixed point problems in Hilbert spaces, *J. Math. Anal. Appl.* 331 (1) (2007) 506–515] to a common element of the set of solutions of an equilibrium problem and the set of fixed points of a nonexpansive mapping.

© 2009 Elsevier Ltd. All rights reserved.

## 1. Introduction

Let  $H$  be a real Hilbert space and let  $C$  be a nonempty closed convex subset of  $H$ . A mapping  $T$  of  $H$  into itself is called nonexpansive if  $\|Tx - Ty\| \leq \|x - y\|$  for all  $x, y \in H$ . We denote by  $F(T)$  the set of fixed points of  $T$  (i.e.  $F(T) = \{x \in H : Tx = x\}$ ). Goebel and Kirk [1] showed that  $F(T)$  is always closed convex, and also nonempty provided  $T$  has a bounded trajectory.

A bounded linear operator  $A$  on  $H$  is called strongly positive with coefficient  $\bar{\gamma}$  if there is a constant  $\bar{\gamma} > 0$  with the property

$$\langle Ax, x \rangle \geq \bar{\gamma} \|x\|^2.$$

Let  $\{T_i\}_{i=1}^N$  be a finite family of nonexpansive mappings with  $F := \bigcap_{i=1}^N F(T_i) \neq \emptyset$ . Many authors (see [2–7]) introduced iterative methods for finding an element of  $F$  which is an optimal point for the minimization problem. For  $n > N$ ,  $T_n$  is understood as  $T_{(n \bmod N)}$  with the mod function taking values in  $\{1, 2, \dots, N\}$ . Let  $u$  be a fixed element of  $H$ . In 2003, Xu [8] proved that the sequence  $\{x_n\}$  generated by

$$x_{n+1} = (1 - \epsilon_n A)T_{n+1}x_n + \epsilon_n u$$

\* Corresponding author. Tel.: +66 53 943327; fax: +66 53 892280.

E-mail addresses: [beawrock@hotmail.com](mailto:beawrock@hotmail.com) (A. Kangtunyakarn), [scmti005@chiangmai.ac.th](mailto:scmti005@chiangmai.ac.th) (S. Suantai).

converges strongly to the solution of the quadratic minimization problem

$$\min_{x \in F} \frac{1}{2} \langle Ax, x \rangle - \langle x, u \rangle$$

under suitable hypotheses on  $\{\epsilon_n\}$  and under the additional hypothesis,

$$F = F(T_1 T_2 \dots T_N) = F(T_N T_1 \dots T_{N-1}) = \dots = F(T_2 T_3 \dots T_N T_1).$$

In 2000, Moudafi [9] introduced the viscosity approximation method for nonexpansive mappings. Let  $f$  be a contraction on  $H$  and  $x_0 \in H$ , define a sequence  $\{x_n\}$  recursively by

$$x_{n+1} = (1 - \sigma_n)Tx_n + \sigma_n f(x_n), \quad n \geq 0, \quad (1.1)$$

where  $\{\sigma_n\}$  is a sequence in  $(0, 1)$ . He proved that under the certain appropriate conditions imposed on  $\{\sigma_n\}$ , the sequence  $\{x_n\}$  generated by (1.1) strongly converges to the unique solution  $x^*$  in  $C$  of the variational inequality

$$\langle (I - f)x^*, x - x^* \rangle \geq 0, \quad \forall x \in C. \quad (1.2)$$

In 2006, Marino and Xu [10] introduced the following general iterative method:

$$x_{n+1} = (1 - \alpha_n A)Tx_n + \alpha_n \gamma f(x_n), \quad n \geq 0, \quad (1.3)$$

where  $\{\alpha_n\}$  is a sequence in  $(0, 1)$  satisfying the following conditions:

- (C1)  $\alpha_n \rightarrow 0$ ;
- (C2)  $\sum_{n=0}^{\infty} \alpha_n = \infty$ ;
- (C3) either  $\sum_{n=0}^{\infty} |\alpha_{n+1} - \alpha_n| < \infty$  or  $\lim_{n \rightarrow \infty} \frac{\alpha_{n+1}}{\alpha_n} = 1$ .

They proved the following theorem:

**Theorem 1.1.** Let  $\{x_n\}$  be generated by algorithm (1.3) with the sequence  $\{\alpha_n\}$  of parameters satisfying conditions (C1)–(C3). Then  $\{x_n\}$  converges strongly to  $x^*$  where  $x^*$  is the unique solution of the following variation inequality:

$$\langle (A - \gamma f)x^*, x^* - z \rangle \leq 0, \quad \forall z \in F(T).$$

Equivalently, we have  $P_{F(T)}(I - A + \gamma f)x^* = x^*$ .

Let  $G : C \times C \rightarrow \mathbb{R}$  be a bifunction. The equilibrium problem for  $G$  is to determine its equilibrium points, i.e. the set

$$EP(G) = \{x \in C : G(x, y) \geq 0, \forall y \in C\}. \quad (1.4)$$

Many problems in physics, optimization, and economics are seeking some elements of  $EP(G)$ , see [11,12]. Several iterative methods have been proposed to solve the equilibrium problem, see, for instance, [4,12–15]. In 2005, Combettes and Hirstoaga [12] introduced some iterative schemes of finding the best approximation to the initial data when  $EP(G)$  is nonempty and proved the strong convergence theorem.

Also in [12] Combettes and Hirstoaga, following [11] define  $S_r : H \rightarrow C$  by

$$S_r(x) = \left\{ z \in C : G(z, y) + \frac{1}{r} \langle y - z, z - x \rangle \geq 0 \forall y \in C \right\}. \quad (1.5)$$

They prove that under suitable hypotheses  $G, S_r$  is single-valued and firmly nonexpansive with  $F(S_r) = EP(G)$ .

In 2007, Takahashi and Takahashi [15] proved the following theorem:

**Theorem 1.2.** Let  $C$  be a nonempty closed convex subset of  $H$ . Let  $G$  be a bifunction from  $C \times C$  to  $\mathbb{R}$  satisfying

- (A1)  $G(x, x) = 0 \forall x \in C$ ;
- (A2)  $G$  is monotone, i.e.  $G(x, y) + G(y, x) \leq 0 \forall x, y \in C$ ;
- (A3)  $\forall x, y, z \in C$ ,

$$\lim_{t \rightarrow 0^+} G(tz + (1-t)x, y) \leq G(x, y).$$

- (A4)  $\forall x \in C, y \mapsto G(x, y)$  is convex and lower semicontinuous;

and let  $S$  be a nonexpansive mapping of  $C$  into  $H$  such that  $F(S) \cap EP(G) \neq \emptyset$ . Let  $f$  be a contraction of  $H$  into itself and let  $\{x_n\}$  and  $\{u_n\}$  be sequences generated by  $x_1 \in H$  and

$$G(u_n, y) + \frac{1}{r_n} \langle y - u_n, u_n - x_n \rangle \geq 0, \quad \forall y \in C$$

$$x_{n+1} = \alpha_n f(x_n) + (1 - \alpha_n) S u_n$$

for all  $n \in \mathbb{N}$ , where  $\{\alpha_n\} \subset [0, 1]$  and  $\{r_n\} \subset (0, 1)$  satisfy (C1)–(C3) and  $\liminf_{n \rightarrow \infty} r_n > 0$  and  $\sum_{n=1}^{\infty} |r_{n+1} - r_n| < \infty$ . Then  $\{x_n\}$  and  $\{u_n\}$  converge strongly to  $z \in F(S) \cap EP(G)$ , where  $z = P_{F(S) \cap EP(G)} f(z)$ .

In 2007, Plubtieng and Punpaeng [13] introduced a general iterative method for finding a common element of  $EP(G)$  and  $F(S)$ . They proved the following theorem.

**Theorem 1.3.** Let  $H$  be a real Hilbert space, let  $G$  be a bifunction from  $H \times H \rightarrow \mathbb{R}$  satisfying (A1)–(A4) and let  $S$  be a nonexpansive mapping on  $H$  such that  $F(S) \cap EP(F) \neq \emptyset$ . Let  $f$  be a contraction of  $H$  into itself with  $\alpha \in (0, 1)$  and let  $A$  be a strongly positive bounded linear operator on  $H$  with coefficients  $\bar{\gamma} > 0$  and  $0 < \gamma < \frac{\bar{\gamma}}{\alpha}$ . Let  $\{x_n\}$  be a sequence generated by  $x_1 \in H$

$$\begin{cases} G(u_n, y) + \frac{1}{r_n} \langle y - u_n, u_n - x_n \rangle \geq 0 & \forall y \in H, \\ x_{n+1} = \alpha_n \gamma f(x_n) + (I - \alpha_n A) S u_n, & \forall n \in \mathbb{N}, \end{cases}$$

where  $u_n = S_{r_n} x_n$ ,  $\{r_n\} \subset (0, 1)$  and  $\{\alpha_n\} \subset [0, 1]$  satisfy (C1)–(C3)  $\liminf_{n \rightarrow \infty} r_n > 0$  and  $\sum_{n=1}^{\infty} |r_{n+1} - r_n| < \infty$ . Then  $\{x_n\}$  and  $\{u_n\}$  converge strongly to  $z \in F(S) \cap EP(F)$  which solves the variational inequality:

$$\langle (A - \gamma f)z, x - z \rangle \geq 0, \quad \forall x \in F(S) \cap EP(G).$$

Equivalently, we have  $P_{F(S) \cap EP(G)}(I - A + \gamma f)z = z$ .

**Question 1.** Are the conditions (C1) and (C2) in Theorems 1.2 and 1.3 sufficient for strong convergence of the sequence  $\{x_n\}$ ?

In 1999, Atsushiba and Takahashi [16] defined the mapping  $W_n$  as follows:

$$\begin{aligned} U_{n,1} &= \lambda_{n,1} T_1 + (1 - \lambda_{n,1}) I, \\ U_{n,2} &= \lambda_{n,2} T_2 U_{n,1} + (1 - \lambda_{n,2}) I, \\ U_{n,3} &= \lambda_{n,3} T_3 U_{n,2} + (1 - \lambda_{n,3}) I, \\ &\vdots \\ U_{n,N-1} &= \lambda_{n,N-1} T_N - 1 U_{n,N-2} + (1 - \lambda_{n,N-1}) I, \\ W_n &= U_{n,N} = \lambda_{n,N} T_N U_{n,N-1} + (1 - \lambda_{n,N}) I, \end{aligned} \quad (1.6)$$

where  $\{\lambda_{n,i}\}_i^N \subseteq [0, 1]$ . This mapping is called the  $W$ -mapping generated by  $T_1, T_2, \dots, T_N$  and  $\lambda_{n,1}, \lambda_{n,2}, \dots, \lambda_{n,N}$ . In 2000 Takahashi and Shimoji [14] proved that if  $X$  is strictly convex Banach space, then  $F(W_n) = \bigcap_{i=1}^N F(T_i)$ , where  $0 < \lambda_{n,i} < 1, i = 1, 2, \dots, N$ .

Very recently, Colao, Marino and Xu [17], introduced a new general iterative method for finding a common element of the set of solutions of equilibrium problem and the set of common fixed points of finite family of nonexpansive mappings in a Hilbert space. They proved that under some sufficient suitable conditions, the sequences  $\{u_n\}$  and  $\{x_n\}$  generated by  $x_1 \in H$  and

$$\begin{cases} G(u_n, y) + \frac{1}{r_n} \langle y - u_n, u_n - x_n \rangle \geq 0, & \forall y \in H, \\ x_{n+1} = \epsilon_n \gamma f(x_n) + \beta x_n + [(1 - \beta)I - \epsilon_n A] W_n u_n \end{cases} \quad (1.7)$$

converge strongly to a point  $x^* \in F$  which is an equilibrium point for  $G$  and is the unique solution of the variational inequality,

$$\langle (A - \gamma f)x^*, x - x^* \rangle \geq 0 \quad \forall x \in F \cap EP(G). \quad (1.8)$$

Motivated by Atsushiba and Takahashi [16], Plubtieng and Punpaeng [13], Colao, Marino and Xu [17], we introduce a new mapping and apply it to the iteration scheme (1.7) to obtain strong convergence to a common element of  $EP(G)$  and  $F$ .

Let  $X$  be a real Banach space and  $C$  a nonempty closed convex subset of  $X$ . For a finite family of nonexpansive mappings  $T_1, T_2, \dots, T_N$  and sequence  $\{\lambda_{n,i}\}_i^N$  in  $[0, 1]$ , we define the mapping  $K_n : C \rightarrow C$  as follows:

$$\begin{aligned} U_{n,1} &= \lambda_{n,1} T_1 + (1 - \lambda_{n,1}) I, \\ U_{n,2} &= \lambda_{n,2} T_2 U_{n,1} + (1 - \lambda_{n,2}) U_{n,1}, \\ U_{n,3} &= \lambda_{n,3} T_3 U_{n,2} + (1 - \lambda_{n,3}) U_{n,2}, \\ &\vdots \\ U_{n,N-1} &= \lambda_{n,N-1} T_N - 1 U_{n,N-2} + (1 - \lambda_{n,N-1}) U_{n,N-2}, \\ K_n &= U_{n,N} = \lambda_{n,N} T_N U_{n,N-1} + (1 - \lambda_{n,N}) U_{n,N-1}. \end{aligned} \quad (1.9)$$

For  $x_1 \in H$ , let  $\{u_n\}$  and  $\{x_n\}$  be the sequence defined by

$$\begin{cases} G(u_n, y) + \frac{1}{r_n} \langle y - u_n, u_n - x_n \rangle \geq 0 & \forall y \in C, \\ x_{n+1} = \epsilon_n \gamma f(x_n) + \beta x_n + ((1 - \beta)I - \epsilon_n A) K_n u_n. \end{cases} \quad (1.10)$$

Please cite this article in press as: A. Kangtanyakarn, S. Suantai, A new mapping for finding common solutions of equilibrium problems and fixed point problems of finite family of nonexpansive mappings, Nonlinear Analysis (2009), doi:10.1016/j.na.2009.03.003

In this paper, we prove that if  $X$  is strictly convex, then  $F(K_n) = \bigcap_{i=1}^N F(T_i)$  where  $0 < \lambda_i < 1$  for every  $i = 1, \dots, N-1$  and  $0 < \lambda_N \leq 1$ , and under the conditions (C1) and (C2) and some other suitable conditions, the sequences  $\{x_n\}$  and  $\{u_n\}$  strongly converge to a point  $x^* = P_{F \cap EP(G)}(I - (A - \gamma f))x^*$ , where  $P_{F \cap EP(G)} : H \rightarrow F \cap EP(G)$  is the metric projection of  $H$  onto  $F \cap EP(G)$ .

## 2. Preliminaries

In this section, we give some useful lemmas that will be used for the main result in the next section.

Let  $C$  be closed convex subset of a Hilbert space  $H$ , let  $P_C$  be the metric projection of  $H$  onto  $C$  i.e., for  $x \in H$ ,  $P_C x$  satisfies the property

$$\|x - P_C x\| = \min_{y \in C} \|x - y\|.$$

The following characterizes the projection  $P_C$ .

**Lemma 2.1** (See [18]). Given  $x \in H$  and  $y \in C$ . Then  $P_C x = y$  if and only if there holds the inequality

$$\langle x - y, y - z \rangle \geq 0 \quad \forall z \in C.$$

**Lemma 2.2** (See [8]). Let  $\{s_n\}$  be a sequence of nonnegative real numbers satisfying

$$s_{n+1} = (1 - \alpha_n)s_n + \alpha_n \beta_n, \quad \forall n \geq 0$$

where  $\{\alpha_n\}, \{\beta_n\}$  satisfy the conditions

$$(1) \quad \{\alpha_n\} \subset [0, 1], \quad \sum_{n=1}^{\infty} \alpha_n = \infty;$$

$$(2) \quad \limsup_{n \rightarrow \infty} \beta_n \leq 0 \quad \text{or} \quad \sum_{n=1}^{\infty} |\alpha_n \beta_n| < \infty.$$

Then  $\lim_{n \rightarrow \infty} s_n = 0$ .

**Lemma 2.3** (See [19]). Let  $\{x_n\}$  and  $\{z_n\}$  be bounded sequences in a Banach space  $X$  and let  $\{\beta_n\}$  be a sequence in  $[0, 1]$  with  $0 < \liminf_{n \rightarrow \infty} \beta_n \leq \limsup_{n \rightarrow \infty} \beta_n < 1$ . Suppose

$$x_{n+1} = \beta_n x_n + (1 - \beta_n) z_n$$

for all integer  $n \geq 0$  and

$$\limsup_{n \rightarrow \infty} (\|z_{n+1} - z_n\| - \|x_{n+1} - x_n\|) \leq 0.$$

Then  $\lim_{n \rightarrow \infty} \|x_n - z_n\| = 0$ .

**Lemma 2.4** (See [10]). Let  $A$  be a strongly positive linear bounded operator on a Hilbert space  $H$  with coefficient  $\bar{\gamma}$  and  $0 < \rho \leq \|A\|^{-1}$ . Then  $\|I - \rho A\| \leq 1 - \rho \bar{\gamma}$ .

**Lemma 2.5** (See [12]). Let  $C$  be a nonempty closed convex subset of a Hilbert space  $H$  and  $G : C \times C \rightarrow \mathbb{R}$  satisfy

- (A1)  $G(x, x) = 0 \quad \forall x \in C$ ;
- (A2)  $G$  is monotone, i.e.  $G(x, y) + G(y, x) \leq 0 \quad \forall x, y \in C$ ;
- (A3)  $\forall x, y, z \in C$ ,

$$\lim_{t \rightarrow 0^+} G(tz + (1-t)x, y) \leq G(x, y);$$

- (A4)  $\forall x \in C, y \mapsto G(x, y)$  is convex and lower semicontinuous.

For  $x \in H$  and  $r > 0$ , set  $S_r : H \rightarrow C$  to be

$$S_r(x) = \left\{ z \in C : G(z, y) + \frac{1}{r} \langle y - z, z - x \rangle \geq 0, \forall y \in C \right\}.$$

Then  $S_r$  is well defined and the following hold:

- (1)  $S_r$  is single-valued;
- (2)  $S_r$  is firmly nonexpansive, i.e.

$$\|S_r(x) - S_r(y)\|^2 \leq \langle S_r(x) - S_r(y), x - y \rangle \quad \forall x, y \in H;$$

- (3)  $F(S_r) = EP(G)$ ;
- (4)  $EP(G)$  is closed and convex.

**Lemma 2.6** (See [18]). *Demiclosedness principle. Assume that  $T$  is a nonexpansive self-mapping of closed convex subset  $C$  of a Hilbert space  $H$ . If  $T$  has a fixed point, then  $I - T$  is demiclosed. That is, whenever  $\{x_n\}$  is a sequence in  $C$  weakly converging to some  $x \in C$  and the sequence  $\{(I - T)x_n\}$  strongly converges to some  $y$  it follows that  $(I - T)x = y$ . Here,  $I$  is the identity mapping of  $H$ .*

**Lemma 2.7.** Let  $H$  be a real Hilbert space. Then, for all  $x, y \in H$ ,

$$\|x + y\|^2 \leq \|x\|^2 + 2\langle y, x + y \rangle.$$

**Lemma 2.8** (See [20]). In a strictly convex Banach space  $E$ , if

$$\|x\| = \|y\| = \|\lambda x + (1 - \lambda)y\|$$

for all  $x, y \in E$  and  $\lambda \in (0, 1)$ , then  $x = y$ .

**Definition 2.1.** Let  $C$  be a nonempty convex subset of a real Banach space. Let  $\{T_i\}_{i=1}^N$  be a finite family of nonexpansive mappings of  $C$  into itself, and let  $\lambda_1, \dots, \lambda_N$  be real numbers such that  $0 \leq \lambda_i \leq 1$  for every  $i = 1, \dots, N$ . We define a mapping  $K : C \rightarrow C$  as follows:

$$\begin{aligned} U_1 &= \lambda_1 T_1 + (1 - \lambda_1)I, \\ U_2 &= \lambda_2 T_2 U_1 + (1 - \lambda_2)U_1, \\ U_3 &= \lambda_3 T_3 U_2 + (1 - \lambda_3)U_2, \\ &\vdots \\ U_{N-1} &= \lambda_{N-1} T_{N-1} U_{N-2} + (1 - \lambda_{N-1})U_{N-2}, \\ K &= U_N = \lambda_N T_N U_{N-1} + (1 - \lambda_N)U_{N-1}. \end{aligned} \tag{2.1}$$

Such a mapping  $K$  is called the  $K$ -mapping generated by  $T_1, \dots, T_N$  and  $\lambda_1, \dots, \lambda_N$ .

**Lemma 2.9.** Let  $C$  be a nonempty closed convex subset of a strictly convex Banach space. Let  $\{T_i\}_{i=1}^N$  be a finite family of nonexpansive mappings of  $C$  into itself with  $\bigcap_{i=1}^N F(T_i) \neq \emptyset$  and let  $\lambda_1, \dots, \lambda_N$  be real numbers such that  $0 < \lambda_i < 1$  for every  $i = 1, \dots, N - 1$  and  $0 < \lambda_N \leq 1$ . Let  $K$  be the  $K$ -mapping generated by  $T_1, \dots, T_N$  and  $\lambda_1, \dots, \lambda_N$ . Then  $F(K) = \bigcap_{i=1}^N F(T_i)$ .

**Proof.** It is easy to see that  $\bigcap_{i=1}^N F(T_i) \subset F(K)$ . Let  $x_0 \in F(K)$  and  $x^* \in \bigcap_{i=1}^N F(T_i)$ . By the definition of  $K$ , we have

$$\begin{aligned} \|x_0 - x^*\| &= \|Kx_0 - x^*\| = \|\lambda_N(T_N U_{N-1}x_0 - x^*) + (1 - \lambda_N)(U_{N-1}x_0 - x^*)\| \\ &\leq \lambda_N \|T_N U_{N-1}x_0 - x^*\| + (1 - \lambda_N) \|U_{N-1}x_0 - x^*\| \\ &\leq \lambda_N \|U_{N-1}x_0 - x^*\| + (1 - \lambda_N) \|U_{N-1}x_0 - x^*\| \\ &= \|U_{N-1}x_0 - x^*\| \\ &= \|\lambda_{N-1}(T_{N-1} U_{N-2}x_0 - x^*) + (1 - \lambda_{N-1})(U_{N-2}x_0 - x^*)\| \\ &\leq \lambda_{N-1} \|T_{N-1} U_{N-2}x_0 - x^*\| + (1 - \lambda_{N-1}) \|U_{N-2}x_0 - x^*\| \\ &\leq \lambda_{N-1} \|U_{N-2}x_0 - x^*\| + (1 - \lambda_{N-1}) \|U_{N-2}x_0 - x^*\| \\ &= \|U_{N-2}x_0 - x^*\| \\ &\vdots \\ &\leq \|U_1x_0 - x^*\| \\ &= \|\lambda_1(T_1x_0 - x^*) + (1 - \lambda_1)(x_0 - x^*)\| \\ &\leq \lambda_1 \|T_1x_0 - x^*\| + (1 - \lambda_1) \|x_0 - x^*\| \\ &\leq \lambda_1 \|x_0 - x^*\| + (1 - \lambda_1) \|x_0 - x^*\| \\ &= \|x_0 - x^*\|. \end{aligned} \tag{2.2}$$

This implies that  $\|x_0 - x^*\| = \|\lambda_1(T_1x_0 - x^*) + (1 - \lambda_1)(x_0 - x^*)\|$  and  $\|x_0 - x^*\| = \|T_1x_0 - x^*\|$ .

By Lemma 2.8, we have  $T_1x_0 = x_0$ , that is  $x_0 \in F(T_1)$ .

It follows that  $U_1x_0 = x_0$ .

By (2.2), we have

$$\begin{aligned} \|x_0 - x^*\| &= \|U_2x_0 - x^*\| = \|\lambda_2(T_2U_1x_0 - x^*) + (1 - \lambda_2)(U_1x_0 - x^*)\| \\ &= \|\lambda_2(T_2x_0 - x^*) + (1 - \lambda_2)(x_0 - x^*)\|. \end{aligned}$$

Again by (2.2) together with  $U_1x_0 = x_0$ , we have

$$\begin{aligned}\|x_0 - x^*\| &= \lambda_2 \|T_2 U_1 x_0 - x^*\| + (1 - \lambda_2) \|U_1 x_0 - x^*\| \\ &= \lambda_2 \|T_2 x_0 - x^*\| + (1 - \lambda_2) \|x_0 - x^*\|,\end{aligned}$$

which implies  $\|x_0 - x^*\| = \|T_2 x_0 - x^*\|$ .

By Lemma 2.8, we have  $T_2 x_0 = x_0$ .

It follows that  $U_2 x_0 = x_0$ .

By using the same argument, we can conclude that  $T_i x_0 = x_0$  and  $U_i x_0 = x_0$  for  $i = 1, 2, \dots, N - 1$ .

This implies that  $0 = x_0 - x_0 = \lambda_N (T_N x_0 - x_0)$ .

It follows that  $x_0 \in F(T_N)$ . Therefore  $x_0 \in \bigcap_{i=1}^N F(T_i)$ .  $\square$

**Lemma 2.10.** Let  $C$  be a nonempty closed convex subset of a Banach space. Let  $\{T_i\}_{i=1}^N$  be a finite family of nonexpansive mappings of  $C$  into itself and  $\{\lambda_{n,i}\}_{i=1}^N$  sequences in  $[0, 1]$  such that  $\lambda_{n,i} \rightarrow \lambda_i$ , as  $n \rightarrow \infty$ , ( $i = 1, 2, \dots, N$ ). Moreover, for every  $n \in \mathbb{N}$  let  $K$  and  $K_n$  be the  $K$ -mappings generated by  $T_1, T_2, \dots, T_N$  and  $\lambda_1, \lambda_2, \dots, \lambda_N$ , and  $T_1, T_2, \dots, T_N$  and  $\lambda_{n,1}, \lambda_{n,2}, \dots, \lambda_{n,N}$  respectively. Then, for every  $x \in C$ , we have

$$\lim_{n \rightarrow \infty} \|K_n x - Kx\| = 0.$$

**Proof.** Let  $x \in C$  and  $U_k$  and  $U_{n,k}$  be generated by  $T_1, T_2, \dots, T_N$  and  $\lambda_1, \lambda_2, \dots, \lambda_N$ , and  $T_1, T_2, \dots, T_N$  and  $\lambda_{n,1}, \lambda_{n,2}, \dots, \lambda_{n,N}$  respectively. Note that

$$\begin{aligned}\|U_{n,1}x - U_1x\| &= \|(\lambda_{n,1} - \lambda_1)T_1x - (\lambda_{n,1} - \lambda_1)x\| \\ &\leq |\lambda_{n,1} - \lambda_1| \|T_1x - x\|.\end{aligned}$$

For  $k \in \{2, 3, \dots, N\}$ , we have

$$\begin{aligned}\|U_{n,k}x - U_kx\| &= \|\lambda_{n,k}T_kU_{n,k-1}x + (1 - \lambda_{n,k})U_{n,k-1}x - \lambda_kT_kU_{k-1}x - (1 - \lambda_k)U_{k-1}x\| \\ &= \|\lambda_{n,k}T_kU_{n,k-1}x + \lambda_{n,k}T_kU_{k-1}x - \lambda_{n,k}T_kU_{k-1}x + \lambda_{n,k}U_{k-1}x - \lambda_{n,k}U_{k-1}x \\ &\quad + (1 - \lambda_{n,k})U_{n,k-1}x - \lambda_kT_kU_{k-1}x - (1 - \lambda_k)U_{k-1}x\| \\ &= \|\lambda_{n,k}(T_kU_{n,k-1}x - T_kU_{k-1}x) + (\lambda_{n,k} - \lambda_k)T_kU_{k-1}x - (1 - \lambda_{n,k})U_{k-1}x \\ &\quad + (\lambda_k - \lambda_{n,k})U_{k-1}x + (1 - \lambda_{n,k})U_{n,k-1}x\| \\ &\leq \lambda_{n,k} \|T_kU_{n,k-1}x - T_kU_{k-1}x\| + |\lambda_{n,k} - \lambda_k| \|T_kU_{k-1}x\| \\ &\quad + (1 - \lambda_{n,k}) \|U_{n,k-1}x - U_{k-1}x\| + |\lambda_k - \lambda_{n,k}| \|U_{k-1}x\| \\ &\leq \lambda_{n,k} \|U_{n,k-1}x - U_{k-1}x\| + (1 - \lambda_{n,k}) \|U_{n,k-1}x - U_{k-1}x\| + |\lambda_{n,k} - \lambda_k| (\|T_kU_{k-1}x\| + \|U_{k-1}x\|) \\ &= \|U_{n,k-1}x - U_{k-1}x\| + |\lambda_{n,k} - \lambda_k| (\|T_kU_{k-1}x\| + \|U_{k-1}x\|).\end{aligned}$$

It follows that

$$\begin{aligned}\|K_nx - Kx\| &= \|U_{n,N}x - U_Nx\| \leq \|U_{n,N-1}x - U_{N-1}x\| + |\lambda_{n,N} - \lambda_N| (\|T_NU_{N-1}x\| + \|U_{N-1}x\|) \\ &\leq \|U_{n,N-2}x - U_{N-2}x\| + |\lambda_{n,N-1} - \lambda_{N-1}| (\|T_{N-1}U_{N-2}x\| + \|U_{N-2}x\|) \\ &\quad + |\lambda_{n,N} - \lambda_N| (\|T_NU_{N-1}x\| + \|U_{N-1}x\|) \\ &= \|U_{n,N-2}x - U_{N-2}x\| + \sum_{j=N-1}^N |\lambda_{n,j} - \lambda_j| (\|T_jU_{j-1}x\| + \|U_{j-1}x\|) \\ &\quad \vdots \\ &\leq \|U_{n,1}x - U_1x\| + \sum_{j=2}^N |\lambda_{n,j} - \lambda_j| (\|T_jU_{j-1}x\| + \|U_{j-1}x\|) \\ &\leq |\lambda_{n,1} - \lambda_1| \|T_1x - x\| + \sum_{j=2}^N |\lambda_{n,j} - \lambda_j| (\|T_jU_{j-1}x\| + \|U_{j-1}x\|).\end{aligned}$$

Since  $\lambda_{n,i} \rightarrow \lambda_i$ , as  $n \rightarrow \infty$ , ( $i = 1, 2, \dots, N$ ) it follows that  $\lim_{n \rightarrow \infty} \|K_nx - Kx\| = 0$ .  $\square$

**Lemma 2.11.** Let  $H$  be a Hilbert space,  $C$  a closed convex nonempty subset of  $H$ ,  $\{T_i\}_{i=1}^N$  a finite family of nonexpansive mappings from  $H$  into itself with  $F = \bigcap_{i=1}^N F(T_i) \neq \emptyset$ , and let  $G : C \times C \rightarrow \mathbb{R}$  be a bifunction satisfying (A1)–(A4). For every  $n \in \mathbb{N}$ , let

Please cite this article in press as: A. Kangtunyakarn, S. Suantai, A new mapping for finding common solutions of equilibrium problems and fixed point problems of finite family of nonexpansive mappings, Nonlinear Analysis (2009), doi:10.1016/j.nla.2009.03.063

Let  $K$  be a  $K$ -mapping generated by  $T_1, \dots, T_N$  and  $\lambda_{n,1}, \dots, \lambda_{n,N}$  with  $\{\lambda_{n,i}\}_{i=1}^N \subset [a, b]$  where  $0 < a \leq b < 1$ . For a sequence  $\{r_n\}$  in  $(0, \infty)$ , let  $S_{r_n} : H \rightarrow C$  be defined by

$$S_{r_n}(x) = \left\{ z \in C : G(z, y) + \frac{1}{r_n} \langle y - z, z - x \rangle \geq 0, \forall y \in C \right\}.$$

If  $\liminf_{n \rightarrow \infty} r_n > 0$ ,  $\lim_{n \rightarrow \infty} \frac{r_n}{r_{n+1}} = 1$  and  $\lim_{n \rightarrow \infty} |\lambda_{n,i} - \lambda_{n-1,i}| = 0 \forall i \in \{1, 2, 3, \dots, N\}$ , then

- (1)  $\lim_{n \rightarrow \infty} \|K_{n+1}S_{r_{n+1}}w_n - K_{n+1}S_{r_n}w_n\| = 0$ ,
- (2)  $\lim_{n \rightarrow \infty} \|K_{n+1}w_n - K_nw_n\| = 0$

for every bounded sequence  $\{w_n\}$  in  $H$ .

**Proof.** By using the nonexpansivity of  $K_{n+1}$  and the proof of Step 2 in Theorem 3.1 of [17], it can be shown that (1) is satisfied.

Next, we show (2). For  $j \in \{2, \dots, N-2\}$ , we have

$$\begin{aligned} \|U_{n+1,N-j}w_n - U_{n,N-j}w_n\| &= \|\lambda_{n+1,N-j}T_{N-j}U_{n+1,N-j-1}w_n + (1 - \lambda_{n+1,N-j})U_{n+1,N-j-1}w_n \\ &\quad - \lambda_{n,N-j}T_{N-j}U_{n,N-j-1}w_n - (1 - \lambda_{n,N-j})U_{n,N-j-1}w_n\| \\ &= \|\lambda_{n+1,N-j}T_{N-j}U_{n+1,N-j-1}w_n - \lambda_{n+1,N-j}T_{N-j}U_{n,N-j-1}w_n \\ &\quad + \lambda_{n+1,N-j}T_{N-j}U_{n,N-j-1}w_n - \lambda_{n+1,N-j}U_{n,N-j-1}w_n \\ &\quad + \lambda_{n+1,N-j}U_{n,N-j-1}w_n + (1 - \lambda_{n+1,N-j})U_{n+1,N-j-1}w_n \\ &\quad - \lambda_{n,N-j}T_{N-j}U_{n,N-j-1}w_n - (1 - \lambda_{n,N-j})U_{n,N-j-1}w_n\| \\ &\leq \lambda_{n+1,N-j}\|T_{N-j}U_{n+1,N-j-1}w_n - T_{N-j}U_{n,N-j-1}w_n\| \\ &\quad + (1 - \lambda_{n+1,N-j})\|U_{n+1,N-j-1}w_n - U_{n,N-j-1}w_n\| \\ &\quad + |\lambda_{n+1,N-j} - \lambda_{n,N-j}|\|T_{N-j}U_{n,N-j-1}w_n\| + |\lambda_{n+1,N-j} - \lambda_{n,N-j}|\|U_{n,N-j-1}w_n\| \\ &\leq \|U_{n+1,N-j-1}w_n - U_{n,N-j-1}w_n\| + M|\lambda_{n+1,N-j} - \lambda_{n,N-j}| \end{aligned} \quad (2.3)$$

where  $M = \sup\{\sum_{j=2}^N (\|T_jU_{n,j-1}w_n\| + \|U_{n,j-1}w_n\|) + \|T_1w_n\| + \|w_n\|\} < \infty$ .

By (2.3), we have

$$\begin{aligned} \|K_{n+1}w_n - K_nw_n\| &= \|U_{n+1,N}w_n - U_{n,N}w_n\| \\ &\leq \|U_{n+1,N-1}w_n - U_{n,N-1}w_n\| + M|\lambda_{n+1,N} - \lambda_{n,N}| \\ &\leq \|U_{n+1,N-2}w_n - U_{n,N-2}w_n\| + M|\lambda_{n+1,N-1} - \lambda_{n,N-1}| + M|\lambda_{n+1,N} - \lambda_{n,N}| \\ &\vdots \\ &\leq M \sum_{j=2}^N |\lambda_{n+1,j} - \lambda_{n,j}| + \|U_{n+1,1}w_n - U_{n,1}w_n\|, \end{aligned} \quad (2.4)$$

and

$$\begin{aligned} \|U_{n+1,1}w_n - U_{n,1}w_n\| &= \|\lambda_{n+1,1}T_1w_n + (1 - \lambda_{n+1,1})w_n - \lambda_{n,1}T_1w_n - (1 - \lambda_{n,1})w_n\| \\ &\leq |\lambda_{n+1,1} - \lambda_{n,1}|\|T_1w_n\| + |\lambda_{n+1,1} - \lambda_{n,1}|\|w_n\| \\ &\leq |\lambda_{n+1,1} - \lambda_{n,1}|M. \end{aligned} \quad (2.5)$$

By (2.4), (2.5) and the condition  $\lim_{n \rightarrow \infty} |\lambda_{n+1,i} - \lambda_{n,i}| = 0$ , we can conclude that

$$\|K_{n+1}w_n - K_nw_n\| \leq M \sum_{j=1}^N |\lambda_{n+1,j} - \lambda_{n,j}| \rightarrow 0 \text{ as } n \rightarrow \infty.$$

Hence (2) is satisfied.  $\square$

## 1. Main result

In this section, we prove the strong convergence of the sequences  $\{u_n\}$  and  $\{x_n\}$  defined by the iteration scheme (1.10).

**Theorem 3.1.** Let  $H$  be a Hilbert space,  $C$  a closed convex nonempty subset of  $H$ ,  $\{T_i\}_{i=1}^N$  a finite family of nonexpansive mappings from  $H$  into itself with  $F = \bigcap_{i=1}^N F(T_i) \neq \emptyset$ ,  $G : C \times C \rightarrow \mathbb{R}$  a bifunction satisfying (A1)–(A4) with  $F \cap EP(G) \neq \emptyset$ ,  $A$  a strongly positive bounded linear operator on  $H$  with coefficient  $\bar{\gamma}$  and  $f$  an  $\alpha$ -contraction on  $H$  for some  $0 < \alpha < 1$ . Moreover, let  $\{\epsilon_n\}$

Please cite this article in press as: A. Kangtunyakarn, S. Suantai, A new mapping for finding common solutions of equilibrium problems and fixed point problems of finite family of nonexpansive mappings, Nonlinear Analysis (2009), doi:10.1016/j.na.2009.03.003

be a sequence in  $(0, 1)$ ,  $\{\lambda_{n,i}\}_{i=1}^N$  sequences in  $[a, b]$  with  $0 < a \leq b < 1$ ,  $\{r_n\}$  a sequence in  $(0, \infty)$  and let  $\gamma$  and  $\beta$  be two real numbers such that  $0 < \beta < 1$  and  $0 < \gamma < \frac{\bar{\gamma}}{\alpha}$ . Assume that

(i) the sequence  $\{r_n\}$  satisfies

$$(D1) \liminf_{n \rightarrow \infty} r_n > 0 \quad \text{and} \quad (D2) \lim_{n \rightarrow \infty} \frac{r_n}{r_{n+1}} = 1,$$

(ii) the finite family of sequences  $\{\lambda_{n,i}\}_{i=1}^N$  satisfies

$$(E1) \lim_{n \rightarrow \infty} |\lambda_{n,i} - \lambda_{n-1,i}| = 0, \quad \forall i = \{1, 2, 3, \dots, N\},$$

(iii) the sequence  $\{\epsilon_n\}$  satisfies

$$(C1) \lim_{n \rightarrow \infty} \epsilon_n = 0, \quad (C2) \sum_{n=1}^{\infty} \epsilon_n = \infty.$$

For every  $n \in N$ , let  $K_n$  be a  $K$ -mapping generated by  $T_1, \dots, T_N$  and  $\lambda_{n,1}, \dots, \lambda_{n,N}$  and let  $\{x_n\}$  and  $\{u_n\}$  be sequences generated by  $x_1 \in C$  and

$$\begin{cases} G(u_n, y) + \frac{1}{r_n} \langle y - u_n, u_n - x_n \rangle \geq 0, & \forall y \in C, \\ x_{n+1} = \epsilon_n \gamma f(x_n) + \beta x_n + ((1 - \beta)I - \epsilon_n A) K_n u_n, \end{cases} \quad (3.1)$$

where  $f : H \rightarrow H$  is an  $\alpha$ -contraction. Then both  $\{x_n\}$  and  $\{u_n\}$  converge strongly to  $x^* \in F = \bigcap_{i=1}^N F(T_i)$  where  $x^*$  is an equilibrium point for  $G$  and is the unique solution of the variational inequality (1.8), i.e.,

$$x^* = P_{F \cap EP(G)}(I - (A - \gamma f))x^*.$$

**Proof.** By Lemma 2.5, it follows that for every  $n \in N$ , there exists a nonexpansive mapping  $S_{r_n} : H \rightarrow H$  such that  $u_n = S_{r_n} x_n$  and  $EP(G) = F(S_{r_n})$ . Whenever needed, we shall write scheme (3.1) as

$$x_{n+1} = \epsilon_n \gamma f(x_n) + \beta x_n + ((1 - \beta)I - \epsilon_n A) K_n S_{r_n} x_n.$$

Moreover, we shall assume that  $\epsilon_n \leq (1 - \beta)\|A\|^{-1}$  and  $1 - \epsilon_n(\bar{\gamma} - \gamma\alpha) > 0$ .

Observe that, if  $\|u\| = 1$ , then

$$\langle ((1 - \beta)I - \epsilon_n A)u, u \rangle = (1 - \beta) - \epsilon_n \langle Au, u \rangle \geq (1 - \beta - \epsilon_n \|A\|) \geq 0.$$

By Lemma 2.4, we have

$$\|(1 - \beta)I - \epsilon_n A\| \leq 1 - \beta - \epsilon_n \bar{\gamma}.$$

We shall divide our proof into 7 steps.

**Step 1.** We shall show that the sequence  $\{x_n\}$  is bounded.

Let  $v \in EP(G) \cap F$ . Then

$$\begin{aligned} \|x_{n+1} - v\| &= \|\epsilon_n \gamma f(x_n) + \beta x_n + ((1 - \beta)I - \epsilon_n A) K_n u_n - v\| \\ &= \|((1 - \beta)I - \epsilon_n A)(K_n u_n - v) + \beta(x_n - v) + \epsilon_n(\gamma f(x_n) - Av)\| \\ &= \|((1 - \beta)I - \epsilon_n A)(K_n u_n - v) + \beta(x_n - v) + \epsilon_n(\gamma f(x_n) - \gamma f(v)) + \epsilon_n(\gamma f(v) - Av)\| \\ &\leq \|(1 - \beta)I - \epsilon_n A\| \|K_n S_{r_n} x_n - K_n S_{r_n} v\| + \beta \|x_n - v\| + \epsilon_n \gamma \alpha \|x_n - v\| + \epsilon_n \|\gamma f(v) - Av\| \\ &\leq (1 - \beta - \epsilon_n \bar{\gamma}) \|x_n - v\| + \beta \|x_n - v\| + \epsilon_n \gamma \alpha \|x_n - v\| + \epsilon_n \|\gamma f(v) - Av\| \\ &= (1 - \epsilon_n(\bar{\gamma} - \gamma\alpha)) \|x_n - v\| + \epsilon_n \|\gamma f(v) - Av\| \\ &\quad + (1 - \epsilon_n(\bar{\gamma} - \gamma\alpha)) \|x_n - v\| + \frac{\epsilon_n(\bar{\gamma} - \gamma\alpha)}{\bar{\gamma} - \gamma\alpha} \|\gamma f(v) - Av\| \\ &\leq \max \left\{ \|x_n - v\|, \frac{\|\gamma f(v) - Av\|}{\bar{\gamma} - \gamma\alpha} \right\}. \end{aligned}$$

By induction we can prove that  $\{x_n\}$  is bounded and also  $\{Ax_n\}$  and  $\{u_n\}$ .

**Step 2.** We will show that  $\lim_{n \rightarrow \infty} \|x_{n+1} - x_n\| = 0$ .

Define sequence  $\{z_n\}$  by  $z_n = \frac{1}{1-\beta}(x_{n+1} - \beta x_n)$ .

Then  $x_{n+1} = \beta x_n + (1 - \beta)z_n$ .

Since  $\{x_n\}$  is bounded, we have, for some big enough constant  $M > 0$ ,

$$\begin{aligned}
 \|z_{n+1} - z_n\| &= \frac{1}{1-\beta} \|x_{n+2} - \beta x_{n+1} - (x_{n+1} - \beta x_n)\| \\
 &= \frac{1}{1-\beta} \|\epsilon_{n+1} \gamma f(x_{n+1}) + ((1-\beta)I - \epsilon_{n+1}A)K_{n+1}u_{n+1} - (\epsilon_n \gamma f(x_n) + ((1-\beta)I - \epsilon_n A)K_n u_n)\| \\
 &= \frac{1}{1-\beta} \|\gamma(\epsilon_{n+1}f(x_{n+1}) - \epsilon_n f(x_n)) + ((1-\beta)I - \epsilon_{n+1}A)K_{n+1}u_{n+1} - ((1-\beta)I - \epsilon_n A)K_n u_n\| \\
 &= \frac{1}{1-\beta} \|\gamma(\epsilon_{n+1}f(x_{n+1}) - \epsilon_n f(x_n)) + (1-\beta)(K_{n+1}u_{n+1} - K_n u_n) - (\epsilon_{n+1}AK_{n+1}u_{n+1} - \epsilon_n AK_n u_n)\| \\
 &= \left\| \frac{\gamma}{1-\beta} (\epsilon_{n+1}f(x_{n+1}) - \epsilon_n f(x_n)) + (K_{n+1}u_{n+1} - K_n u_n) - \frac{1}{1-\beta} (\epsilon_{n+1}AK_{n+1}u_{n+1} - \epsilon_n AK_n u_n) \right\| \\
 &\leq \frac{\gamma}{1-\beta} (\epsilon_{n+1}\|f(x_{n+1})\| + \epsilon_n\|f(x_n)\|) + \|K_{n+1}u_{n+1} - K_n u_n\| + \frac{1}{1-\beta} (\epsilon_{n+1}\|AK_{n+1}u_{n+1}\| + \epsilon_n\|AK_n u_n\|) \\
 &\leq \|K_{n+1}S_{r_{n+1}}x_{n+1} - K_n S_{r_n}x_n\| + M(\epsilon_n + \epsilon_{n+1}) \\
 &\leq \|K_{n+1}S_{r_{n+1}}x_{n+1} - K_{n+1}S_{r_{n+1}}x_n\| + \|K_{n+1}S_{r_{n+1}}x_n - K_n S_{r_n}x_n\| + M(\epsilon_n + \epsilon_{n+1}) \\
 &\leq \|x_{n+1} - x_n\| + \|K_{n+1}S_{r_{n+1}}x_n - K_{n+1}S_{r_n}x_n\| + \|K_{n+1}S_{r_n}x_n - K_n S_{r_n}x_n\| + M(\epsilon_n + \epsilon_{n+1}).
 \end{aligned}$$

By condition on  $\{\epsilon_n\}$  and by Lemma 2.11, we can conclude that

$$\limsup_{n \rightarrow \infty} (\|z_{n+1} - z_n\| - \|x_{n+1} - x_n\|) \leq 0.$$

By Lemma 2.3, we obtain

$$\lim_{n \rightarrow \infty} \|x_n - z_n\| = 0.$$

This implies

$$\lim_{n \rightarrow \infty} \|x_{n+1} - x_n\| = (1-\beta) \lim_{n \rightarrow \infty} \|x_n - z_n\| = 0.$$

Step 3. We will show that  $\lim_{n \rightarrow \infty} \|x_n - K_n u_n\| = 0$  where  $u_n = S_{r_n}x_n$ .

Since

$$\begin{aligned}
 \|x_n - K_n u_n\| &\leq \|x_n - x_{n+1}\| + \|x_{n+1} - K_n u_n\| \\
 &= \|x_n - x_{n+1}\| + \|\epsilon_n \gamma f(x_n) + \beta x_n + (1-\beta)K_n u_n - \epsilon_n AK_n u_n - K_n u_n\| \\
 &\leq \|x_n - x_{n+1}\| + \epsilon_n \|\gamma f(x_n) - AK_n u_n\| + \beta \|x_n - K_n u_n\|,
 \end{aligned}$$

we have

$$\|x_n - K_n u_n\| \leq \frac{1}{(1-\beta)} (\|x_n - x_{n+1}\| + \epsilon_n \|\gamma f(x_n) - AK_n u_n\|).$$

By (3.1) and Step 2, we obtain  $\lim_{n \rightarrow \infty} \|x_n - K_n u_n\| = 0$ .

Step 4. We shall show that  $\lim_{n \rightarrow \infty} \|x_n - S_{r_n}x_n\| = 0$ .

Let  $v \in F \cap EP(G)$ . Since  $S_{r_n}$  is firmly nonexpansive, we have

$$\begin{aligned}
 \|v - S_{r_n}x_n\|^2 &= \|S_{r_n}v - S_{r_n}x_n\|^2 \\
 &\leq \langle S_{r_n}v - S_{r_n}x_n, v - x_n \rangle \\
 &= \frac{1}{2} (\|S_{r_n}x_n - v\|^2 + \|x_n - v\|^2 - \|S_{r_n}x_n - x_n\|^2).
 \end{aligned}$$

$$\|S_{r_n}x_n - v\|^2 \leq \|x_n - v\|^2 - \|S_{r_n}x_n - x_n\|^2. \quad (3.2)$$

Let  $\lambda_n = \gamma f(x_n) - AK_n u_n$  and  $\lambda > 0$  be a constant such that

$$\lambda > \sup_k \{\|y_k\|, \|x_k - v\|\}. \quad (3.3)$$

By (3.2) and (3.3), we have

$$\begin{aligned}
 \|x_{n+1} - v\|^2 &= \|\epsilon_n \gamma f(x_n) + \beta x_n + ((1 - \beta)I - \epsilon_n A)K_n u_n - v\|^2 \\
 &= \|[(1 - \beta)I - \epsilon_n A](K_n u_n - v) + \beta(x_n - v) + \epsilon_n(\gamma f(x_n) - Av)\|^2 \\
 &= \|(1 - \beta)(K_n u_n - v) - \epsilon_n A(K_n u_n - v) + \beta(x_n - v) + \epsilon_n(\gamma f(x_n) - Av)\|^2 \\
 &= \|(1 - \beta)(K_n u_n - v) + \beta(x_n - v) + \epsilon_n(\gamma f(x_n) - A(K_n u_n))\|^2 \\
 &\leq \|(1 - \beta)(K_n u_n - v) + \beta(x_n - v)\|^2 + 2\epsilon_n \langle \gamma f(x_n), x_{n+1} - v \rangle \\
 &\leq \|(1 - \beta)(K_n S_{r_n} x_n - v) + \beta(x_n - v)\|^2 + 2\epsilon_n \lambda^2 \\
 &\leq (1 - \beta) \|K_n S_{r_n} x_n - v\|^2 + \beta \|x_n - v\|^2 + 2\epsilon_n \lambda^2 \\
 &\leq (1 - \beta) \|S_{r_n} x_n - v\|^2 + \beta \|x_n - v\|^2 + 2\epsilon_n \lambda^2 \\
 &\leq \|x_n - v\|^2 - (1 - \beta) \|S_{r_n} x_n - x_n\|^2 + 2\epsilon_n \lambda^2.
 \end{aligned}$$

It follows that

$$\begin{aligned}
 \|S_{r_n} x_n - x_n\|^2 &\leq \frac{1}{1 - \beta} (\|x_n - v\|^2 - \|x_{n+1} - v\|^2 + 2\epsilon_n \lambda^2) \\
 &= \frac{1}{1 - \beta} ((\|x_n - v\| - \|x_{n+1} - v\|)(\|x_n - v\| + \|x_{n+1} - v\|) + 2\epsilon_n \lambda^2) \\
 &\leq \frac{1}{1 - \beta} (\|x_{n+1} - x_n\|(\|x_n - v\| + \|x_{n+1} - v\|) + 2\epsilon_n \lambda^2).
 \end{aligned}$$

By  $\|x_{n+1} - x_n\| \rightarrow 0$  and  $\epsilon_n \rightarrow 0$ , as  $n \rightarrow \infty$ , we obtain that

$$\lim_{n \rightarrow \infty} \|x_n - S_{r_n} x_n\| = 0.$$

**Step 5.** Let  $\omega(x_n)$  be the set of all weak  $\omega$ -limits of  $\{x_n\}$ . We shall show that  $\omega(x_n) \subset F \cap EP(G)$ . It is a consequence of Step 4 and [12, Lemma 2.13] that  $\omega(x_n) \subset EP(G)$ .

So, it remains to prove that  $z \in F$ . To see this, we observe that we may assume that

$$\lambda_{n_m, k} \rightarrow \lambda_k \in (0, 1) \text{ as } m \rightarrow \infty \text{ (} k = 1, 2, \dots, N \text{)}.$$

Let  $K$  be the  $K$ -mapping generated by  $T_1, T_2, \dots, T_N$  and  $\lambda_1, \dots, \lambda_N$ , then by Lemma 2.10, we have, for every  $x \in C$ ,

$$K_{n_m} x \rightarrow Kx \text{ as } m \rightarrow \infty. \quad (3.4)$$

We will show that  $z \in F = \bigcap_{i=1}^N F(T_i)$ . Assume that there exists  $j \in \{1, 2, \dots, N\}$  such that  $z \neq T_j z$ . By Lemma 2.9, we have  $z \neq Wz$ . Since  $z \in EP(G) = F(S_{r_n})$ , by Step 3, (3.4) and Opial's property of Hilbert space, we have

$$\begin{aligned}
 \liminf_{m \rightarrow \infty} \|x_{n_m} - z\| &< \liminf_{m \rightarrow \infty} \|x_{n_m} - Kz\| \\
 &\leq \liminf_{m \rightarrow \infty} (\|x_{n_m} - K_{n_m} S_{r_{n_m}} x_{n_m}\| + \|K_{n_m} S_{r_{n_m}} x_{n_m} - K_{n_m} S_{r_{n_m}} z\| + \|K_{n_m} S_{r_{n_m}} z - Kz\|) \\
 &\leq \liminf_{m \rightarrow \infty} \|x_{n_m} - z\|.
 \end{aligned}$$

This is a contradiction, then  $z \in F = \bigcap_{i=1}^N F(T_i)$ .

**Step 6.** Let  $x^*$  be the unique solution of the variational inequality,

$$\langle (A - \gamma f)x^*, x - x^* \rangle \geq 0, \quad \forall x \in F \cap EP(G). \quad (3.5)$$

We shall show that  $\limsup_{n \rightarrow \infty} \langle (\gamma f - A)x^*, x_n - x^* \rangle \leq 0$ .

Let  $\{x_{n_k}\}$  be a subsequence of  $\{x_n\}$  such that

$$\lim_{k \rightarrow \infty} \langle (\gamma f - A)x^*, x_{n_k} - x^* \rangle = \limsup_{n \rightarrow \infty} \langle (\gamma f - A)x^*, x_n - x^* \rangle. \quad (3.6)$$

Without loss of generality, we may assume that  $\{x_{n_k}\}$  weakly converges to some  $z$  in  $H$ . By Step 5,  $z \in F \cap EP(G)$ . Thus combining (3.5) and (3.6), we get

$$\begin{aligned}
 \limsup_{n \rightarrow \infty} \langle (\gamma f - A)x^*, x_n - x^* \rangle &= \lim_{k \rightarrow \infty} \langle (\gamma f - A)x^*, x_{n_k} - x^* \rangle \\
 &= \langle (\gamma f - A)x^*, z - x^* \rangle \leq 0
 \end{aligned} \quad (3.7)$$

as required.

**Step 7.** Finally, we will show that the sequences  $\{x_n\}$  and  $\{u_n\}$  converge strongly to  $x^* \in F \cap EP(G)$ . Let  $x^*$  be the unique fixed point of the mapping  $P_{F \cap EP(G)}(I - (A - \gamma f))$ , i.e. the unique solution of the variational inequality (1.8). By Lemmas 2.4 and 2.7, we have

$$\begin{aligned}
 \|x_{n+1} - x^*\|^2 &= \|\epsilon_n \gamma f(x_n) + \beta x_n + ((1 - \beta)I - \epsilon_n A)K_n u_n - x^*\|^2 \\
 &= \|((1 - \beta)I - \epsilon_n A)(K_n u_n - x^*) + \beta(x_n - x^*) + \epsilon_n(\gamma f(x_n) - Ax^*)\|^2 \\
 &\leq \|((1 - \beta)I - \epsilon_n A)(K_n u_n - x^*) + \beta(x_n - x^*)\|^2 + 2\epsilon_n \langle \gamma f(x_n) - Ax^*, x_{n+1} - x^* \rangle \\
 &= \left\| \frac{(1 - \beta)((1 - \beta)I - \epsilon_n A)(K_n u_n - x^*)}{(1 - \beta)} + \beta(x_n - x^*) \right\|^2 \\
 &\quad + 2\epsilon_n \gamma \langle f(x_n) - f(x^*), x_{n+1} - x^* \rangle + 2\epsilon_n \langle \gamma f(x^*) - Ax^*, x_{n+1} - x^* \rangle \\
 &\leq (1 - \beta) \left\| \frac{((1 - \beta)I - \epsilon_n A)(K_n u_n - x^*)}{(1 - \beta)} \right\|^2 + \beta \|x_n - x^*\|^2 \\
 &\quad + 2\epsilon_n \gamma \alpha \|x_n - x^*\| \|x_{n+1} - x^*\| + 2\epsilon_n \langle \gamma f(x^*) - Ax^*, x_{n+1} - x^* \rangle \\
 &\leq \frac{\|((1 - \beta)I - \epsilon_n A)(K_n u_n - x^*)\|^2}{(1 - \beta)} + \beta \|x_n - x^*\|^2 \\
 &\quad + 2\epsilon_n \gamma \alpha \|x_n - x^*\| \|x_{n+1} - x^*\| + 2\epsilon_n \langle \gamma f(x^*) - Ax^*, x_{n+1} - x^* \rangle \\
 &\leq \frac{\|(1 - \beta)I - \epsilon_n A\|^2}{(1 - \beta)} \|K_n u_n - x^*\|^2 + \beta \|x_n - x^*\|^2 \\
 &\quad + \epsilon_n \gamma \alpha [\|x_n - x^*\|^2 + \|x_{n+1} - x^*\|^2] + 2\epsilon_n \langle \gamma f(x^*) - Ax^*, x_{n+1} - x^* \rangle \\
 &\leq \frac{(1 - \beta - \epsilon_n \bar{\gamma})^2}{(1 - \beta)} \|x_n - x^*\|^2 + \beta \|x_n - x^*\|^2 + \epsilon_n \gamma \alpha (\|x_n - x^*\|^2 + \|x_{n+1} - x^*\|^2) \\
 &\quad + 2\epsilon_n \langle \gamma f(x^*) - Ax^*, x_{n+1} - x^* \rangle \\
 &= \left( \frac{(1 - \beta - \epsilon_n \bar{\gamma})^2}{(1 - \beta)} + \beta + \epsilon_n \gamma \alpha \right) \|x_n - x^*\|^2 + \epsilon_n \gamma \alpha \|x_{n+1} - x^*\|^2 \\
 &\quad + 2\epsilon_n \langle \gamma f(x^*) - Ax^*, x_{n+1} - x^* \rangle \\
 &= \left( \frac{(1 - \beta)^2 - 2(1 - \beta)\epsilon_n \bar{\gamma} + \epsilon_n^2 \bar{\gamma}^2}{(1 - \beta)} + \beta + \epsilon_n \gamma \alpha \right) \|x_n - x^*\|^2 + \epsilon_n \gamma \alpha \|x_{n+1} - x^*\|^2 \\
 &\quad + 2\epsilon_n \langle \gamma f(x^*) - Ax^*, x_{n+1} - x^* \rangle \\
 &= \left( (1 - \beta) - 2\epsilon_n \bar{\gamma} + \frac{\epsilon_n^2 \bar{\gamma}^2}{(1 - \beta)} + \beta + \epsilon_n \gamma \alpha \right) \|x_n - x^*\|^2 + \epsilon_n \gamma \alpha \|x_{n+1} - x^*\|^2 \\
 &\quad + 2\epsilon_n \langle \gamma f(x^*) - Ax^*, x_{n+1} - x^* \rangle \\
 &= \left( 1 - (2\bar{\gamma} - \alpha \gamma)\epsilon_n + \frac{\epsilon_n^2 \bar{\gamma}^2}{(1 - \beta)} \right) \|x_n - x^*\|^2 + \epsilon_n \gamma \alpha \|x_{n+1} - x^*\|^2 + 2\epsilon_n \langle \gamma f(x^*) - Ax^*, x_{n+1} - x^* \rangle,
 \end{aligned}$$

which implies

$$\begin{aligned}
 \|x_{n+1} - x^*\|^2 &\leq \frac{1}{1 - \epsilon_n \gamma \alpha} \left( 1 - (2\bar{\gamma} - \alpha \gamma)\epsilon_n + \frac{\epsilon_n^2 \bar{\gamma}^2}{(1 - \beta)} \right) \|x_n - x^*\|^2 + \frac{1}{1 - \epsilon_n \gamma \alpha} (2\epsilon_n \langle \gamma f(x^*) - Ax^*, x_{n+1} - x^* \rangle) \\
 &= \frac{1}{1 - \epsilon_n \gamma \alpha} ((1 - (2\bar{\gamma} - \alpha \gamma)\epsilon_n)) \|x_n - x^*\|^2 \\
 &\quad + \frac{1}{1 - \epsilon_n \gamma \alpha} \left( 2\epsilon_n \langle \gamma f(x^*) - Ax^*, x_{n+1} - x^* \rangle + \frac{\epsilon_n^2 \bar{\gamma}^2}{(1 - \beta)} \|x_n - x^*\|^2 \right) \\
 &= \frac{1}{1 - \epsilon_n \gamma \alpha} (1 - 2\epsilon_n \bar{\gamma} + \alpha \gamma \epsilon_n) \|x_n - x^*\|^2 \\
 &\quad + \frac{1}{1 - \epsilon_n \gamma \alpha} \left( 2\epsilon_n \langle \gamma f(x^*) - Ax^*, x_{n+1} - x^* \rangle + \frac{\epsilon_n^2 \bar{\gamma}^2}{(1 - \beta)} \|x_n - x^*\|^2 \right) \\
 &= \frac{1}{1 - \epsilon_n \gamma \alpha} (1 - 2\epsilon_n \bar{\gamma} + 2\alpha \gamma \epsilon_n - \alpha \gamma \epsilon_n) \|x_n - x^*\|^2 \\
 &\quad + \frac{1}{1 - \epsilon_n \gamma \alpha} \left( 2\epsilon_n \langle \gamma f(x^*) - Ax^*, x_{n+1} - x^* \rangle + \frac{\epsilon_n^2 \bar{\gamma}^2}{(1 - \beta)} \|x_n - x^*\|^2 \right)
 \end{aligned}$$

$$\begin{aligned}
&= \frac{1}{1 - \epsilon_n \gamma \alpha} (1 - \alpha \gamma \epsilon_n - 2\epsilon_n(\bar{\gamma} - \alpha \gamma)) \|x_n - x^*\|^2 \\
&\quad + \frac{\epsilon_n}{1 - \epsilon_n \gamma \alpha} \left( 2\langle \gamma f(x^*) - Ax^*, x_{n+1} - x^* \rangle + \frac{\epsilon_n \bar{\gamma}^2}{(1 - \beta)} \|x_n - x^*\|^2 \right) \\
&= \left( 1 - \frac{2\epsilon_n(\bar{\gamma} - \alpha \gamma)}{1 - \epsilon_n \gamma \alpha} \right) \|x_n - x^*\|^2 + \frac{\epsilon_n}{1 - \epsilon_n \gamma \alpha} \left( 2\langle \gamma f(x^*) - Ax^*, x_{n+1} - x^* \rangle + \frac{\epsilon_n \bar{\gamma}^2}{(1 - \beta)} \|x_n - x^*\|^2 \right) \\
&= \left( 1 - \frac{2\epsilon_n(\bar{\gamma} - \alpha \gamma)}{1 - \epsilon_n \gamma \alpha} \right) \|x_n - x^*\|^2 + \frac{2(\bar{\gamma} - \alpha \gamma)}{2(\bar{\gamma} - \alpha \gamma)} \frac{\epsilon_n}{1 - \epsilon_n \gamma \alpha} \\
&\quad \times \left( 2\langle \gamma f(x^*) - Ax^*, x_{n+1} - x^* \rangle + \frac{\epsilon_n \bar{\gamma}^2}{(1 - \beta)} \|x_n - x^*\|^2 \right) \\
&= \left( 1 - \frac{2\epsilon_n(\bar{\gamma} - \alpha \gamma)}{1 - \epsilon_n \gamma \alpha} \right) \|x_n - x^*\|^2 + \frac{2\epsilon_n(\bar{\gamma} - \alpha \gamma)}{1 - \epsilon_n \gamma \alpha} \\
&\quad \times \left( \frac{\langle \gamma f(x^*) - Ax^*, x_{n+1} - x^* \rangle}{(\bar{\gamma} - \alpha \gamma)} + \frac{\epsilon_n \bar{\gamma}^2}{2(1 - \beta)(\bar{\gamma} - \alpha \gamma)} \|x_n - x^*\|^2 \right). \tag{3.8}
\end{aligned}$$

We can rewrite (3.8) as

$$\|x_{n+1} - x^*\|^2 \leq (1 - \xi_n) \|x_n - x^*\|^2 + \xi_n \delta_n$$

where  $\xi_n = \frac{2\epsilon_n(\bar{\gamma} - \alpha \gamma)}{1 - \epsilon_n \gamma \alpha}$  and  $\delta_n = \left( \frac{\langle \gamma f(x^*) - Ax^*, x_{n+1} - x^* \rangle}{(\bar{\gamma} - \alpha \gamma)} + \frac{\epsilon_n \bar{\gamma}^2}{2(1 - \beta)(\bar{\gamma} - \alpha \gamma)} \|x_n - x^*\|^2 \right)$ .

By our hypotheses it is easily verified that  $\sum_{n=1}^{\infty} \xi_n = \infty$  and  $\limsup_{n \rightarrow \infty} \delta_n \leq 0$ .

Therefore, by Lemma 2.2, we can conclude that  $\|x_n - x^*\| \rightarrow 0$ .

Since  $\|u_n - x^*\| = \|S_{r_n} x_n - x^*\| \leq \|x_n - x^*\|$ , it follows that  $u_n \rightarrow x^*$  in norm. This completes the proof.  $\square$

**Remark.** (1) If we take  $N = 1$ ,  $T_1 = S$  and  $G(x, y) = 0$  for all  $x, y \in C$  and  $r_n = 1$  for all  $n \in \mathbb{N}$ , then the iterative scheme (3.1) reduces to the following scheme:

$$x_1 \in H, \quad x_{n+1} = \epsilon_n \gamma f(x_n) + \beta x_n + ((1 - \beta)I - \epsilon_n A) S x_n, \tag{3.9}$$

which is a modification of the iterative scheme (1.3) and by Theorem 3.1 we observe that the conditions (C1) and (C2) are sufficient for strong convergence of the sequence  $\{x_n\}$  generated by (3.9) to a fixed point of  $S$ .

(2) If we take  $N = 1$ ,  $T_1 = S$  and  $A = I$ , then the iterative scheme (3.1) reduces to the following scheme:

$$\begin{cases} x_1 \in C, \\ G(u_n, y) + \frac{1}{r_n} \langle y - u_n, u_n - x_n \rangle \geq 0, \quad \forall y \in C, \\ x_{n+1} = \epsilon_n f(x_n) + \beta x_n + (1 - \beta - \epsilon_n) S u_n, \end{cases} \tag{3.10}$$

which is a modification of the scheme in Theorem 1.2 defined by Takahashi and Takahashi [15], and by Theorem 3.1, we obtain strong convergence of the sequence  $\{x_n\}$  generated by (3.10) under the sufficient conditions of Theorem 1.2 but without the condition (C3).

(3) If we take  $N = 1$  and  $T_1 = S$  in Theorem 3.1, the iterative scheme (3.1) reduces to the following scheme:

$$\begin{cases} x_1 \in H, G(u_n, y) + \frac{1}{r_n} \langle y - u_n, u_n - x_n \rangle \geq 0, \quad \forall y \in H, \\ x_{n+1} = \epsilon_n \gamma f(x_n) + \beta x_n + ((1 - \beta)I - \epsilon_n A) S u_n, \end{cases} \tag{3.11}$$

which is a modification of the scheme in Theorem 1.3, and by Theorem 3.1, we obtain strong convergence of the sequence  $\{x_n\}$  generated by (3.11) under some sufficient conditions without the condition (C3).

## Acknowledgments

The authors would like to thank the Thailand Research Fund and Commission on Higher Education for their financial support during the preparation of this paper. The first author was also supported by the Graduate school Chiang Mai University.

## References

- [1] K. Goebel, W.A. Kirk, Topics in Metric Fixed Point Theory, in: Cambridge Stud. Adv. Math., vol. 28, Cambridge University Press, Cambridge, 1990.
- [2] H.H. Bauschke, The approximation of fixed points of compositions of nonexpansive mappings in Hilbert space, *J. Math. Anal. Appl.* 202 (1) (1996) 150–159.
- [3] H.H. Bauschke, J.M. Borwein, On projection algorithms for solving convex feasibility problems, *SIAM Rev.* 38 (3) (1996) 367–426.

Please cite this article in press as: A. Kangtunyakarn, S. Suantai, A new mapping for finding common solutions of equilibrium problems and fixed point problems of finite family of nonexpansive mappings, *Nonlinear Analysis* (2009), doi:10.1016/j.na.2009.03.003

# ARTICLE IN PRESS

A. Kangtunyakarn, S. Suantai / *Nonlinear Analysis* ■ (■■■■) ■■■–■■■

- [4] P.L. Combettes, The foundations of set theoretic estimation, *Proc. IEEE* 81 (2) (1993) 182–208.
- [5] P.L. Combettes, Constrained image recovery in a product space, in: *Proceedings of the IEEE International Conference on Image Processing*, Washington, DC, 1995, IEEE Computer Society Press, California, 1995, pp. 2025–2028.
- [6] F. Deutsch, H. Hundal, The rate of convergence of Dykstras cyclic projections algorithm: The polyhedral case, *Numer. Funct. Anal. Optim.* 15 (56) (1994) 537–565.
- [7] D.C. Youla, Mathematical theory of image restoration by the method of convex projections, in: H. Stark (Ed.), *Image Recovery: Theory and Applications*, Academic Press, Florida, 1987, pp. 29–77.
- [8] H.K. Xu, An iterative approach to quadratic optimization, *J. Optim. Theory Appl.* 116 (3) (2003) 659–678.
- [9] A. Moudafi, Viscosity approximation methods for fixed-points problems, *J. Math. Anal. Appl.* 24 (1) (2000) 46–55.
- [10] G. Marino, H.K. Xu, A general iterative method for nonexpansive mappings in Hilbert spaces, *J. Math. Anal. Appl.* 318 (1) (2006) 43–52.
- [11] E. Blum, W. Oettli, From optimization and variational inequalities to equilibrium problems, *Math. Student* 63 (14) (1994) 123–145.
- [12] P.L. Combettes, S.A. Hirstoaga, Equilibrium programming in Hilbert spaces, *J. Nonlinear Convex Anal.* 6 (1) (2005) 117–136.
- [13] S. Plubtieng, R. Punpaeng, A general iterative method for equilibrium problems and fixed point problems in Hilbert spaces, *J. Math. Anal. Appl.* 336 (1) (2007) 455–469.
- [14] W. Takahashi, K. Shimoji, Convergence theorems for nonexpansive mappings and feasibility problems, *Math. Comput. Modelling* 32 (2000) 1463–1471.
- [15] S. Takahashi, W. Takahashi, Viscosity approximation methods for equilibrium problems and fixed point problems in Hilbert spaces, *J. Math. Anal. Appl.* 331 (1) (2007) 506–515.
- [16] S. Atsushiba, W. Takahashi, Strong convergence theorems for a finite family of nonexpansive mappings and applications, in: B.N. Prasad Birth Centenary Commemoration Volume, *Indian J. Math.* 41 (3) (1999) 435–453.
- [17] V. Colao, G. Marino, H.K. Xu, An iterative method for finding common solutions of equilibrium and fixed point problems, *J. Math. Anal. Appl.* 344 (2008) 340–352.
- [18] F.E. Browder, Convergence of approximants to fixed points of nonexpansive nonlinear mappings in Banach space, *Arch. Ration. Mech. Anal.* 24 (1967) 82–89.
- [19] T. Suzuki, Strong convergence of Krasnoselskii and Manns type sequences for one-parameter nonexpansive semigroups without Bochner integrals, *J. Math. Anal. Appl.* 305 (1) (2005) 227–239.
- [20] W. Takahashi, *Nonlinear Functional Analysis: Fixed Point Theory and Its Applications*, Yokohama Publishers, Yokohama, 2000.

Please cite this article in press as: A. Kangtunyakarn, S. Suantai, A new mapping for finding common solutions of equilibrium problems and fixed point problems of finite family of nonexpansive mappings, *Nonlinear Analysis* (2009), doi:10.1016/j.na.2009.03.003



# Hybrid iterative scheme for generalized equilibrium problems and fixed point problems of finite family of nonexpansive mappings

Atid Kangtunyakarn, Suthep Suantai\*

Department of Mathematics, Faculty of Science, Chiang Mai University, Chiang Mai, 50200, Thailand

## ARTICLE INFO

### Article history:

Received 14 January 2009

Accepted 21 January 2009

### Keywords:

Strong convergence

Finite families of nonexpansive mapping

Fixed point

Generalized equilibrium problem

Inverse-strongly monotone

## ABSTRACT

In this paper, we introduce a new mapping and a Hybrid iterative scheme for finding a common element of the set of solutions of a generalized equilibrium problem and the set of common fixed points of a finite family of nonexpansive mappings in a Hilbert space. Then, we prove the strong convergence of the proposed iterative algorithm to a common fixed point of a finite family of nonexpansive mappings which is a solution of the generalized equilibrium problem. The results obtained in this paper extend the recent ones of Takahashi and Takahashi [S. Takahashi, W. Takahashi, Strong convergence theorem for a generalized equilibrium problem and a nonexpansive mapping in a Hilbert space, *Nonlinear Anal.* 69 (2008) 1025–1033].

© 2009 Elsevier Ltd. All rights reserved.

## 1. Introduction

Let  $H$  be a real Hilbert space and let  $C$  be a nonempty closed convex subset of  $H$  and  $A : C \rightarrow H$  be a nonlinear mapping and let  $P_C$  be the projection of  $H$  onto the convex subset  $C$ . A mapping  $T$  of  $H$  into itself is called *nonexpansive* if  $\|Tx - Ty\| \leq \|x - y\|$  for all  $x, y \in H$ . We denote by  $F(T)$  the set of fixed points of  $T$  (i.e.  $F(T) = \{x \in H : Tx = x\}$ ). Goebel and Kirk [1] showed that  $F(T)$  is always closed convex, and also nonempty provided  $T$  has a bounded trajectory. Let  $\{T_i\}_{i=1}^N$  be a finite family of nonexpansive mappings with  $\bigcap_{i=1}^N F(T_i) \neq \emptyset$ .

Let  $F : C \times C \rightarrow \mathbb{R}$  be a bifunction. The equilibrium problem for  $F$  is to determine its equilibrium points, i.e. the set

$$EP(F) = \{x \in C : F(x, y) \geq 0, \forall y \in C\}. \quad (1.1)$$

Many problems in physics, optimization, and economics require some elements of  $EP(F)$ , see [2–7]. Several iterative methods have been proposed to solve the equilibrium problem, see for instance [3,5–7]. In 2005, Combettes and Hirstoaga [3] introduced an iterative scheme for finding the best approximation to the initial data when  $EP(F)$  is nonempty and proved a strong convergence theorem.

The variational inequality problem is to find  $u \in C$  such that

$$\langle Au, v - u \rangle \geq 0 \quad (1.2)$$

for all  $v \in C$ . The set of solutions of the variational inequality is denoted by  $VI(C, A)$ .

For a bifunction  $F : C \times C \rightarrow \mathbb{R}$  and a nonlinear mapping  $A : C \rightarrow H$ , we consider the following equilibrium problem:

$$\text{Find } z \in C \text{ such that } F(z, y) + \langle Az, y - z \rangle \geq 0, \quad \forall y \in C. \quad (1.3)$$

The set of such  $z \in C$  is denoted by  $EP$ , i.e.,

$$EP = \{z \in C : F(z, y) + \langle Az, y - z \rangle \geq 0, \forall y \in C\}.$$

\* Corresponding author.

E-mail addresses: [beawrock@hotmail.com](mailto:beawrock@hotmail.com) (A. Kangtunyakarn), [scmti005@chiangmai.ac.th](mailto:scmti005@chiangmai.ac.th) (S. Suantai).

In the case of  $A \equiv 0$ ,  $EP$  is denoted by  $EP(F)$ . In the case of  $F \equiv 0$ ,  $EP$  is also denoted by  $VI(C, A)$ . Numerous problems in physics, optimization, variational inequalities, minimax problems, the Nash equilibrium problem in noncooperative games, economics reduce to finding a solution of (1.3) see, for instance, [2,4].

A mapping  $A$  of  $C$  into  $H$  is called  $\alpha$ -inverse strongly monotone, see [8], if there exists a positive real number  $\alpha$  such that

$$\langle x - y, Ax - Ay \rangle \geq \alpha \|Ax - Ay\|^2$$

for all  $x, y \in C$ .

For  $r > 0$ , let  $T_r : H \rightarrow C$  be defined by

$$T_r(x) = \left\{ z \in C : F(z, y) + \frac{1}{r}(y - z, z - x) \geq 0, \forall y \in C \right\}. \quad (1.4)$$

Combettes and Hirstoaga [9] showed that under some suitable conditions of  $F$ ,  $T_r$  is single-valued and firmly nonexpansive and satisfies  $F(T_r) = EP(F)$ .

In 2007, Takahashi and Takahashi [6] introduced a hybrid viscosity approximation method in the framework of a real Hilbert space  $H$ . They defined the iterative sequences  $\{x_n\}$  and  $\{u_n\}$  as follows:

$$\begin{cases} x_1 \in H, \text{ arbitrarily;} \\ F(u_n, y) + \frac{1}{r_n}(y - u_n, u_n - x_n) \geq 0, \quad \forall y \in C, \\ x_{n+1} = \alpha_n f(x_n) + (1 - \alpha_n)T_{r_n}u_n, \quad \forall n \in \mathbb{N}, \end{cases} \quad (1.5)$$

where  $f : H \rightarrow H$  is a contraction mapping with a constant  $\alpha \in (0, 1)$  and  $\{\alpha_n\} \subset [0, 1]$ ,  $\{r_n\} \subset (0, \infty)$ . They proved, under some suitable conditions on the sequence  $\{\alpha_n\}$ ,  $\{r_n\}$  and bifunction  $F$ , that  $\{x_n\}$  and  $\{u_n\}$  strongly converge to  $z \in F(T) \cap EP(F)$ , where  $z = P_{F(T) \cap EP(F)}f(z)$ .

Recently, in 2008, Takahashi and Takahashi [7] introduced a hybrid iterative method for finding a common element of  $EP$  and  $F(T)$ . They defined  $\{x_n\}$  in the following way:

$$\begin{cases} u, x_1 \in C, \text{ arbitrarily;} \\ F(z_n, y) + \langle Ax_n, y - z_n \rangle + \frac{1}{\lambda_n}(y - z_n, z_n - x_n) \geq 0, \quad \forall y \in C, \\ x_{n+1} = \beta_n x_n + (1 - \beta_n)T(a_n u + (1 - a_n)z_n), \quad \forall n \in \mathbb{N}, \end{cases} \quad (1.6)$$

where  $A$  be an  $\alpha$ -inverse strongly monotone mapping of  $C$  into  $H$  with positive real number  $\alpha$ , and  $\{a_n\} \in [0, 1]$ ,  $\{\beta_n\} \subset [0, 1]$ ,  $\{\lambda_n\} \subset [0, 2\alpha]$ , and proved strong convergence of the scheme (1.6) to  $z \in \bigcap_{i=1}^N F(T_i) \cap EP$ , where  $z = P_{\bigcap_{i=1}^N F(T_i) \cap EP}u$  in the framework of a Hilbert space, under some suitable conditions on  $\{a_n\}$ ,  $\{\beta_n\}$ ,  $\{\lambda_n\}$  and bifunction  $F$ .

In 1999, Atsushiba and Takahashi [10] defined the mapping  $W_n$  as follows:

$$\begin{aligned} U_{n,1} &= \lambda_{n,1}T_1 + (1 - \lambda_{n,1})I, \\ U_{n,2} &= \lambda_{n,2}T_2U_{n,1} + (1 - \lambda_{n,2})I, \\ U_{n,3} &= \lambda_{n,3}T_3U_{n,2} + (1 - \lambda_{n,3})I, \\ &\vdots \\ U_{n,N-1} &= \lambda_{n,N-1}T_{N-1}U_{n,N-2} + (1 - \lambda_{n,N-1})I, \\ W_n &= U_{n,N} = \lambda_{n,N}T_NU_{n,N-1} + (1 - \lambda_{n,N})I, \end{aligned} \quad (1.7)$$

where  $\{\lambda_{n,i}\}_i^N \subseteq [0, 1]$ . This mapping is called the  $W$ -mapping generated by  $T_1, T_2, \dots, T_N$  and  $\lambda_{n,1}, \lambda_{n,2}, \dots, \lambda_{n,N}$ . In 2000, Takahashi and Shimoji [11] proved that if  $X$  is a strictly convex Banach space, then  $F(W_n) = \bigcap_{i=1}^N F(T_i)$ , where  $0 < \lambda_{n,i} < 1$ ,  $i = 1, 2, \dots, N$ .

Let  $X$  be a real Hilbert space and  $C$  a nonempty closed convex subset of  $X$  and let  $\{T_i\}_{i=1}^N$  be a finite family of nonexpansive mappings of  $C$  into itself. For each  $n \in \mathbb{N}$ , and  $j = 1, 2, \dots, N$ , let  $\alpha_j^{(n)} = (\alpha_1^{n,j}, \alpha_2^{n,j}, \alpha_3^{n,j})$  be such that  $\alpha_1^{n,j}, \alpha_2^{n,j}, \alpha_3^{n,j} \in [0, 1]$  with  $\alpha_1^{n,j} + \alpha_2^{n,j} + \alpha_3^{n,j} = 1$ . We define mapping  $S_n : C \rightarrow C$  as follows:

$$\begin{aligned} U_{n,0} &= I \\ U_{n,1} &= \alpha_1^{n,1}T_1U_{n,0} + \alpha_2^{n,1}U_{n,0} + \alpha_3^{n,1}I \\ U_{n,2} &= \alpha_1^{n,2}T_2U_{n,1} + \alpha_2^{n,2}U_{n,1} + \alpha_3^{n,2}I \\ U_{n,3} &= \alpha_1^{n,3}T_3U_{n,2} + \alpha_2^{n,3}U_{n,2} + \alpha_3^{n,3}I \\ &\vdots \\ U_{n,N-1} &= \alpha_1^{n,N-1}T_{N-1}U_{n,N-2} + \alpha_2^{n,N-1}U_{n,N-2} + \alpha_3^{n,N-1}I \\ S_n &= U_{n,N} = \alpha_1^{n,N}T_NU_{n,N-1} + \alpha_2^{n,N}U_{n,N-1} + \alpha_3^{n,N}I. \end{aligned}$$

The mapping  $S_n$  is called the  $S$ -mapping generated by  $T_1, T_2, \dots, T_N$  and  $\alpha_1^{(n)}, \alpha_2^{(n)}, \dots, \alpha_N^{(n)}$ . For given  $u \in C$  and  $x_1 \in C$ , let  $\{z_n\} \subset C$  and  $\{x_n\} \subset C$  be sequences generated by

$$\begin{cases} F(z_n, y) + \langle Ax_n, y - z_n \rangle + \frac{1}{\lambda_n} \langle y - z_n, z_n - x_n \rangle \geq 0, & \forall y \in C, \\ x_{n+1} = \beta_n x_n + (1 - \beta_n) S_n(a_n u + (1 - a_n) z_n). & \forall n \in \mathbb{N}. \end{cases} \quad (1.8)$$

In this paper, we show that if  $X$  is strictly convex, then  $F(S_n) = \bigcap_{i=1}^N F(T_i)$  if  $\alpha_1^{n,j} \in (0, 1)$  for all  $j = 1, 2, \dots, N-1$ ,  $\alpha_1^{n,N} \in (0, 1)$  and  $\alpha_2^{n,j}, \alpha_3^{n,j} \in [0, 1)$  for all  $j = 1, 2, \dots, N$ , and we prove that under some suitable conditions, the sequence  $\{x_n\}$  converges strongly to a point  $z = P_{\bigcap_{i=1}^N F(T_i) \cap EP} u$ .

## 2. Preliminaries

In this section, we collect and give some useful lemmas that will be used for our main result in the next section.

Let  $C$  be the closed convex subset of a real Hilbert space  $H$ , let  $P_C$  be the metric projection of  $H$  onto  $C$  i.e., for  $x \in H$ ,  $P_C x$  satisfies the property

$$\|x - P_C x\| = \min_{y \in C} \|x - y\|.$$

The following characterizes the projection  $P_C$ .

**Lemma 2.1** (See [12]). Given  $x \in H$  and  $y \in C$ . Then  $P_C x = y$  if and only if there holds the inequality  $\langle x - y, y - z \rangle \geq 0 \quad \forall z \in C$ .

**Lemma 2.2** (See [11]). In a strictly convex Banach space  $E$ , if

$$\|x\| = \|y\| = \|\lambda x + (1 - \lambda)y\|$$

for all  $x, y \in E$  and  $\lambda \in (0, 1)$ , then  $x = y$ .

**Lemma 2.3** (See [13]). Let  $\{s_n\}$  be a sequence of nonnegative real numbers satisfying

$$s_{n+1} = (1 - \alpha_n)s_n + \alpha_n \beta_n, \quad \forall n \geq 0 \text{ where } \{\alpha_n\}, \{\beta_n\} \text{ satisfy the conditions}$$

$$(1) \{\alpha_n\} \subset [0, 1], \quad \sum_{n=1}^{\infty} \alpha_n = \infty, \quad (2) \limsup_{n \rightarrow \infty} \beta_n \leq 0.$$

Then  $\lim_{n \rightarrow \infty} s_n = 0$ .

**Lemma 2.4** (See [14]). Let  $\{x_n\}$  and  $\{z_n\}$  be bounded sequences in a Banach space  $X$  and let  $\{\beta_n\}$  be a sequence in  $[0, 1]$  with  $0 < \liminf_{n \rightarrow \infty} \beta_n \leq \limsup_{n \rightarrow \infty} \beta_n < 1$ . Suppose

$$x_{n+1} = \beta_n x_n + (1 - \beta_n) z_n \text{ for all integer } n \geq 0 \text{ and } \limsup_{n \rightarrow \infty} (\|z_{n+1} - z_n\| - \|x_{n+1} - x_n\|) \leq 0.$$

Then  $\lim_{n \rightarrow \infty} \|x_n - z_n\| = 0$ .

For solving the equilibrium problem for a bifunction  $F : C \times C \rightarrow \mathbb{R}$ , let us assume that  $F$  satisfies the following conditions:

(A1)  $F(x, x) = 0 \quad \forall x \in C$ ;

(A2)  $F$  is monotone, i.e.  $F(x, y) + F(y, x) \leq 0, \quad \forall x, y \in C$ ;

(A3)  $\forall x, y, z \in C$ ,

$$\lim_{t \rightarrow 0^+} F(tz + (1 - t)x, y) \leq F(x, y);$$

(A4)  $\forall x \in C, y \mapsto F(x, y)$  is convex and lower semicontinuous.

The following lemma appears implicitly in [2].

**Lemma 2.5** (See [2]). Let  $C$  be a nonempty closed convex subset of  $H$  and let  $F$  be a bifunction of  $C \times C$  into  $\mathbb{R}$  satisfying (A1)–(A4). Let  $r > 0$  and  $x \in H$ . Then, there exists  $z \in C$  such that

$$F(z, y) + \frac{1}{r} \langle y - z, z - x \rangle \geq 0 \quad (2.1)$$

for all  $y \in C$ .

**Lemma 2.6** (See [9]). Assume that  $F : C \times C \rightarrow \mathbb{R}$  satisfies (A1)–(A4). For  $r > 0$  and  $x \in H$ , define a mapping  $T_r : H \rightarrow C$  as follows:

$$T_r(x) = \left\{ z \in C : F(z, y) + \frac{1}{r} \langle y - z, z - x \rangle \geq 0, \quad \forall y \in C \right\} \quad (2.2)$$

for all  $z \in H$ . Then, the following hold:

- (1)  $T_r$  is single-valued;  
 (2)  $T_r$  is firmly nonexpansive i.e.

$$\|T_r(x) - T_r(y)\|^2 \leq \langle T_r(x) - T_r(y), x - y \rangle \quad \forall x, y \in H;$$

- (3)  $F(T_r) = EP(F)$ ;  
 (4)  $EP(F)$  is closed and convex.

**Definition 2.7.** Let  $C$  be a nonempty convex subset of real Banach space. Let  $\{T_i\}_{i=1}^N$  be a finite family of nonexpansive mappings of  $C$  into itself. For each  $j = 1, 2, \dots, N$ , let  $\alpha_j = (\alpha_1^j, \alpha_2^j, \alpha_3^j)$  where  $\alpha_1^j, \alpha_2^j, \alpha_3^j \in [0, 1]$  and  $\alpha_1^j + \alpha_2^j + \alpha_3^j = 1$ . We define the mapping  $S : C \rightarrow C$  as follows:

$$\begin{aligned} U_0 &= I \\ U_1 &= \alpha_1^1 T_1 U_0 + \alpha_2^1 U_0 + \alpha_3^1 I \\ U_2 &= \alpha_1^2 T_2 U_1 + \alpha_2^2 U_1 + \alpha_3^2 I \\ U_3 &= \alpha_1^3 T_3 U_2 + \alpha_2^3 U_2 + \alpha_3^3 I \\ &\vdots \\ U_{N-1} &= \alpha_1^{N-1} T_{N-1} U_{N-2} + \alpha_2^{N-1} U_{N-2} + \alpha_3^{N-1} I \\ S &= U_N = \alpha_1^N T_N U_{N-1} + \alpha_2^N U_{N-1} + \alpha_3^N I. \end{aligned} \quad (2.3)$$

This mapping is called  $S$ -mapping generated by  $T_1, \dots, T_N$  and  $\alpha_1, \alpha_2, \dots, \alpha_N$ .

Next, we prove a lemma which is very useful for our consideration.

**Lemma 2.8.** Let  $C$  be a nonempty closed convex subset of a strictly convex Banach space. Let  $\{T_i\}_{i=1}^N$  be a finite family of nonexpansive mappings of  $C$  into itself with  $\bigcap_{i=1}^N F(T_i) \neq \emptyset$  and let  $\alpha_j = (\alpha_1^j, \alpha_2^j, \alpha_3^j)$ ,  $j = 1, 2, 3, \dots, N$ , where  $\alpha_1^j, \alpha_2^j, \alpha_3^j \in [0, 1]$ ,  $\alpha_1^j + \alpha_2^j + \alpha_3^j = 1$ ,  $\alpha_1^j \in (0, 1)$  for all  $j = 1, 2, \dots, N-1$ ,  $\alpha_1^N \in (0, 1]$ ,  $\alpha_2^j, \alpha_3^j \in [0, 1)$  for all  $j = 1, 2, \dots, N$ . Let  $S$  be the mapping generated by  $T_1, \dots, T_N$  and  $\alpha_1, \alpha_2, \dots, \alpha_N$ . Then  $F(S) = \bigcap_{i=1}^N F(T_i)$ .

**Proof.** It is clear that  $\bigcap_{i=1}^N F(T_i) \subseteq F(S)$ . Next, we show that  $F(S) \subseteq \bigcap_{i=1}^N F(T_i)$ . To show this, let  $x_0 \in F(S)$  and  $x^* \in \bigcap_{i=1}^N F(T_i)$ . Then we have

$$\begin{aligned} \|x_0 - x^*\| &= \|Sx_0 - x^*\| = \|\alpha_1^N (T_N U_{N-1} x_0 - x^*) + \alpha_2^N (U_{N-1} x_0 - x^*) + \alpha_3^N (x_0 - x^*)\| \\ &\leq \alpha_1^N \|T_N U_{N-1} x_0 - x^*\| + \alpha_2^N \|U_{N-1} x_0 - x^*\| + \alpha_3^N \|x_0 - x^*\| \\ &\leq (1 - \alpha_3^N) \|U_{N-1} x_0 - x^*\| + (1 - (1 - \alpha_3^N)) \|x_0 - x^*\| \\ &= (1 - \alpha_3^N) \|\alpha_1^{N-1} (T_{N-1} U_{N-2} x_0 - x^*) + \alpha_2^{N-1} (U_{N-2} x_0 - x^*) + \alpha_3^{N-1} (x_0 - x^*)\| \\ &\quad + (1 - (1 - \alpha_3^N)) \|x_0 - x^*\| \\ &\leq (1 - \alpha_3^N) (\alpha_1^{N-1} \|T_{N-1} U_{N-2} x_0 - x^*\| + \alpha_2^{N-1} \|U_{N-2} x_0 - x^*\| + \alpha_3^{N-1} \|x_0 - x^*\|) \\ &\quad + (1 - (1 - \alpha_3^N)) \|x_0 - x^*\| \end{aligned} \quad (2.4)$$

$$\leq \prod_{j=N-1}^N (1 - \alpha_3^j) \|U_{N-2} x_0 - x^*\| + \left(1 - \prod_{j=N-1}^N (1 - \alpha_3^j)\right) \|x_0 - x^*\| \quad (2.5)$$

$$\begin{aligned} &= \prod_{j=N-1}^N (1 - \alpha_3^j) \|\alpha_1^{N-2} (T_{N-2} U_{N-3} x_0 - x^*) + \alpha_2^{N-2} (U_{N-3} x_0 - x^*) + \alpha_3^{N-2} (x_0 - x^*)\| \\ &\quad + \left(1 - \prod_{j=N-1}^N (1 - \alpha_3^j)\right) \|x_0 - x^*\| \end{aligned} \quad (2.6)$$

$$\begin{aligned} &\leq \prod_{j=N-1}^N (1 - \alpha_3^j) (\alpha_1^{N-2} \|T_{N-2} U_{N-3} x_0 - x^*\| + \alpha_2^{N-2} \|U_{N-3} x_0 - x^*\| + \alpha_3^{N-2} \|x_0 - x^*\|) \\ &\quad + \left(1 - \prod_{j=N-1}^N (1 - \alpha_3^j)\right) \|x_0 - x^*\| \end{aligned}$$

$$\begin{aligned}
&\leq \prod_{j=N-2}^N (1 - \alpha_3^j) \|U_{N-3}x_0 - x^*\| + \left(1 - \prod_{j=N-2}^N (1 - \alpha_3^j)\right) \|x_0 - x^*\| \\
&\leq \dots \\
&\leq \prod_{j=3}^N (1 - \alpha_3^j) \|\alpha_1^2 (T_2 U_1 x_0 - x^*) + \alpha_2^2 (U_1 x_0 - x^*) + \alpha_3^2 (x_0 - x^*)\| \\
&\quad + \left(1 - \prod_{j=3}^N (1 - \alpha_3^j)\right) \|x_0 - x^*\| \tag{2.7}
\end{aligned}$$

$$\begin{aligned}
&\leq \prod_{j=3}^N (1 - \alpha_3^j) (\alpha_1^2 \|T_2 U_1 x_0 - x^*\| + \alpha_2^2 \|U_1 x_0 - x^*\| + \alpha_3^2 \|x_0 - x^*\|) \\
&\quad + \left(1 - \prod_{j=3}^N (1 - \alpha_3^j)\right) \|x_0 - x^*\| \\
&\leq \prod_{j=2}^N (1 - \alpha_3^j) \|U_1 x_0 - x^*\| + \left(1 - \prod_{j=2}^N (1 - \alpha_3^j)\right) \|x_0 - x^*\| \tag{2.8}
\end{aligned}$$

$$= \prod_{j=2}^N (1 - \alpha_3^j) \|\alpha_1^1 (T_1 x_0 - x^*) + (1 - \alpha_1^1) (x_0 - x^*)\| + \left(1 - \prod_{j=2}^N (1 - \alpha_3^j)\right) \|x_0 - x^*\| \tag{2.9}$$

$$\begin{aligned}
&\leq \prod_{j=2}^N (1 - \alpha_3^j) (\alpha_1^1 \|T_1 x_0 - x^*\| + (1 - \alpha_1^1) \|x_0 - x^*\|) \\
&\quad + \left(1 - \prod_{j=2}^N (1 - \alpha_3^j)\right) \|x_0 - x^*\| \tag{2.10} \\
&\leq \prod_{j=2}^N (1 - \alpha_3^j) \|x_0 - x^*\| + \left(1 - \prod_{j=2}^N (1 - \alpha_3^j)\right) \|x_0 - x^*\| \\
&= \|x_0 - x^*\|.
\end{aligned}$$

This implies by (2.9) that

$$\|x_0 - x^*\| = \prod_{j=2}^N (1 - \alpha_3^j) \|\alpha_1^1 (T_1 x_0 - x^*) + (1 - \alpha_1^1) (x_0 - x^*)\| + \left(1 - \prod_{j=2}^N (1 - \alpha_3^j)\right) \|x_0 - x^*\|,$$

hence

$$\|x_0 - x^*\| = \|\alpha_1^1 (T_1 x_0 - x^*) + (1 - \alpha_1^1) (x_0 - x^*)\|. \tag{2.11}$$

By (2.10), we obtain

$$\|x_0 - x^*\| = \prod_{j=2}^N (1 - \alpha_3^j) [\alpha_1^1 \|T_1 x_0 - x^*\| + (1 - \alpha_1^1) \|x_0 - x^*\|] + \left(1 - \prod_{j=2}^N (1 - \alpha_3^j)\right) \|x_0 - x^*\|,$$

which implies

$$\|x_0 - x^*\| = \alpha_1^1 \|T_1 x_0 - x^*\| + (1 - \alpha_1^1) \|x_0 - x^*\|.$$

It follows that

$$\|x_0 - x^*\| = \|T_1 x_0 - x^*\|. \tag{2.12}$$

From (2.11) and (2.12), we have by Lemma 2.2 that  $T_1 x_0 = x_0$ , that is  $x_0 \in F(T_1)$ .

It implies that

$$U_1 x_0 = \lambda_1 T_1 x_0 + (1 - \lambda_1) x_0 = x_0.$$

By (2.7), we have

$$\|x_0 - x^*\| = \prod_{j=3}^N (1 - \alpha_3^j) \|\alpha_1^2 (T_2 U_1 x_0 - x^*) + \alpha_2^2 (U_1 x_0 - x^*) + \alpha_3^2 (x_0 - x^*)\| + \left[1 - \prod_{j=3}^N (1 - \alpha_3^j)\right] \|x_0 - x^*\|.$$

It follows that

$$\begin{aligned} \|x_0 - x^*\| &= \|\alpha_1^2 (T_2 U_1 x_0 - x^*) + \alpha_2^2 (U_1 x_0 - x^*) + \alpha_3^2 (x_0 - x^*)\| \\ &= \|\alpha_1^2 (T_2 x_0 - x^*) + (1 - \alpha_1^2) (x_0 - x^*)\|. \end{aligned} \quad (2.13)$$

By (2.8), we have

$$\|x_0 - x^*\| = \prod_{j=3}^N (1 - \alpha_3^j) (\alpha_1^2 \|T_2 U_1 x_0 - x^*\| + \alpha_2^2 \|U_1 x_0 - x^*\| + \alpha_3^2 \|x_0 - x^*\|) + \left(1 - \prod_{j=3}^N (1 - \alpha_3^j)\right) \|x_0 - x^*\|,$$

which implies

$$\begin{aligned} \|x_0 - x^*\| &= \alpha_1^2 \|T_2 U_1 x_0 - x^*\| + \alpha_2^2 \|U_1 x_0 - x^*\| + \alpha_3^2 \|x_0 - x^*\| \\ &= \alpha_1^2 \|T_2 x_0 - x^*\| + (1 - \alpha_1^2) \|x_0 - x^*\|. \end{aligned}$$

Hence, we obtain

$$\|x_0 - x^*\| = \|T_2 x_0 - x^*\|. \quad (2.14)$$

From (2.13) and (2.14), we have by Lemma 2.2 that  $T_2 x_0 = x_0$ , that is  $x_0 \in F(T_2)$ .

This implies that  $U_2 x_0 = \alpha_1^2 T_2 U_1 x_0 + \alpha_2^2 U_1 x_0 + \alpha_3^2 x_0 = x_0$ .

By continuing in this way, we can show that  $x_0 \in F(T_i)$  and  $x_0 \in F(U_i)$  for all  $i = 1, 2, \dots, N-1$ .

Finally, we shall show that  $x_0 \in F(T_N)$ .

Since

$$\begin{aligned} 0 &= Sx_0 - x_0 = \alpha_1^N T_N U_{N-1} x_0 + \alpha_2^N U_{N-1} x_0 + \alpha_3^N x_0 - x_0 \\ &= \alpha_1^N (T_N x_0 - x_0), \end{aligned}$$

and  $\alpha_1^N \in (0, 1]$ , we obtain  $T_N x_0 = x_0$  so that  $x_0 \in F(T_N)$ . Hence  $F(S) \subseteq \bigcap_{i=1}^N F(T_i)$ .  $\square$

**Lemma 2.9.** Let  $C$  be a nonempty closed convex subset of Banach space. Let  $\{T_i\}_{i=1}^N$  be a finite family of nonexpansive mappings of  $C$  into itself and for each  $n \in \mathbb{N}$  and  $j \in \{1, 2, \dots, N\}$ , let  $\alpha_j^{(n)} = (\alpha_1^{n,j}, \alpha_2^{n,j}, \alpha_3^{n,j})$ ,  $\alpha_j = (\alpha_1^j, \alpha_2^j, \alpha_3^j)$  where  $\alpha_1^{n,j}, \alpha_2^{n,j}, \alpha_3^{n,j} \in [0, 1]$ ,  $\alpha_1^j, \alpha_2^j, \alpha_3^j \in [0, 1]$ ,  $\alpha_1^{n,j} + \alpha_2^{n,j} + \alpha_3^{n,j} = 1$  and  $\alpha_1^j + \alpha_2^j + \alpha_3^j = 1$ . Suppose  $\alpha_i^{n,j} \rightarrow \alpha_i^j$  as  $n \rightarrow \infty$  for  $i = 1, 2, 3$  and  $j = 1, 2, 3, \dots, N$ . Let  $S$  and  $S_n$  be the  $S$ -mappings generated by  $T_1, T_2, \dots, T_N$  and  $\alpha_1, \alpha_2, \dots, \alpha_N$  and  $T_1, T_2, \dots, T_N$  and  $\alpha_1^{(n)}, \alpha_2^{(n)}, \dots, \alpha_N^{(n)}$ , respectively. Then  $\lim_{n \rightarrow \infty} \|S_n x - Sx\| = 0$  for every  $x \in C$ .

**Proof.** Let  $x \in C$ ,  $U_k$  and  $U_{n,k}$  be generated by  $T_1, T_2, \dots, T_N$  and  $\alpha_1, \alpha_2, \dots, \alpha_N$  and  $T_1, T_2, \dots, T_N$  and  $\alpha_1^{(n)}, \alpha_2^{(n)}, \dots, \alpha_N^{(n)}$ , respectively. For each  $n \in \mathbb{N}$  and for  $k \in \{2, 3, \dots, N\}$ , we have

$$\begin{aligned} \|U_{n,1}x - U_1x\| &= \|\alpha_1^{n,1} T_1 x + (1 - \alpha_1^{n,1})x - \alpha_1^1 T_1 x - (1 - \alpha_1^1)x\| \\ &= |\alpha_1^{n,1} - \alpha_1^1| \|T_1 x - x\|, \end{aligned} \quad (2.15)$$

and

$$\begin{aligned} \|U_{n,k}x - U_kx\| &= \|\alpha_1^{n,k} T_k U_{n,k-1}x + \alpha_2^{n,k} U_{n,k-1}x + \alpha_3^{n,k} x - \alpha_1^k T_k U_{k-1}x - \alpha_2^k U_{k-1}x - \alpha_3^k x\| \\ &= \|\alpha_1^{n,k} (T_k U_{n,k-1}x - T_k U_{k-1}x) + (\alpha_1^{n,k} - \alpha_1^k) T_k U_{k-1}x \\ &\quad + (\alpha_3^{n,k} - \alpha_3^k)x + \alpha_2^{n,k} (U_{n,k-1}x - U_{k-1}x) + (\alpha_2^{n,k} - \alpha_2^k) U_{k-1}x\| \\ &\leq \alpha_1^{n,k} \|T_k U_{n,k-1}x - T_k U_{k-1}x\| + |\alpha_1^{n,k} - \alpha_1^k| \|T_k U_{k-1}x\| \\ &\quad + |\alpha_3^{n,k} - \alpha_3^k| \|x\| + \alpha_2^{n,k} \|U_{n,k-1}x - U_{k-1}x\| + |\alpha_2^{n,k} - \alpha_2^k| \|U_{k-1}x\| \\ &\leq \alpha_1^{n,k} \|U_{n,k-1}x - U_{k-1}x\| + |\alpha_1^{n,k} - \alpha_1^k| \|T_k U_{k-1}x\| \\ &\quad + \alpha_2^{n,k} \|U_{n,k-1}x - U_{k-1}x\| + (|\alpha_1^k - \alpha_1^{n,k}| + |\alpha_3^{n,k} - \alpha_3^k|) \|U_{k-1}x\| + |\alpha_3^{n,k} - \alpha_3^k| \|x\| \\ &\leq \|U_{n,k-1}x - U_{k-1}x\| + |\alpha_1^{n,k} - \alpha_1^k| (\|T_k U_{k-1}x\| + \|U_{k-1}x\|) \\ &\quad + |\alpha_3^{n,k} - \alpha_3^k| (\|U_{k-1}x\| + \|x\|). \end{aligned} \quad (2.16)$$