

บทคัดย่อ

ออกซิโดสควาลีนไซเคเลส เป็นเอนไซม์สำคัญในชีวสังเคราะห์ของไตรเทอร์พีน ทำหน้าที่เปลี่ยนออกซิโดสควาลีน ไปเป็นไตรเทอร์พีนวงแหวน ได้หลากหลายชนิด โดยเอนไซม์นี้ถูกสร้างโดย ยีนออกซิโดสควาลีนไซเคเลส โครงการนี้มีวัตถุประสงค์เพื่อโคลนยีนออกซิโดสควาลีนไซเคเลส จากใบบัวบกและใบเปล้าน้อย ด้วยการเทคนิคพีซีอาร์ โดยออกแบบไพรเมอร์ของ ออกซิโดสควาลีน ไซเคเลส จากข้อมูลยีนที่มีรายงานมาก่อนนี้ ร่วมกับการใช้เทคนิคการหาลำดับนิวคลีโอไทด์ส่วปลาย 3' และ 5' ของยีน ในการโคลนยีนในส่วนกลางและปลายของยีน สามารถโคลนยีนจากบัวบกได้ 2 ชนิด คือ CAL และ CAM ซึ่งมีความยาว 1572 คู่เบส และ 1269 คู่เบส ตามลำดับ และโคลนยีนจากเปล้าน้อยได้ 2 ชนิดคือ CSC และ CSD ซึ่งมีความยาว 1502 คู่เบส และ 1497 คู่เบส ตามลำดับ แต่ไม่สามารถโคลนยีนในส่วนต้น (ปลาย 5') ได้สำเร็จ อย่างไรก็ตามในการศึกษานี้ได้โคลนยีนออกซิโดสควาลีนไซเคเลส ชนิด CabAS จากบัวบก ตามข้อมูลที่มีรายงานไว้ก่อนหน้านี้ (Kim et al., 2005) เรียกยีนที่โคลนได้จากโครงการนี้ ว่า CAK โดยยีน CAK มีความแตกต่างของลำดับนิวคลีโอไทด์ 3 ตำแหน่ง ส่งผลให้มีลำดับกรดอะมิโนต่างจากยีนCabAS 3 ตำแหน่งคือ V631G, V645A และ F692L ซึ่งผลการศึกษาบทบาทหน้าที่ของ CAK พบว่าเป็น multifunctional triterpene synthase ซึ่งสร้าง α -และ β -amyrin เป็นหลัก ต่างจาก CabAS ที่รายงานว่าเป็น dammarenediol synthase (Kim et al., 2009)

Abstract

Oxidosqualene cyclases (OSC) are significant enzymes in the triterpene biosynthesis. OSC convert 2,3-oxidosqualene to several cyclic triterpenes. The OSC was controlled by oxidosqualene cyclase gene (*osc*). In this study, oxidosqualene cyclases from *Centella asiatica* and *Croton stellatopilosus* leaves were cloned using homology-base PCR and RACE technics. Two partial cDNAs, 1572-bp CAL and 1269-bp CAM, were obtained from *C. asiatica*. Other two, 1502-bp CSC and 1497-bp CSD were obtained from *C. stellatopilosus*. These partial cDNA are the fragment from core to 3'-end of each gene. The 5'-end cDNA fragments have not been successfully cloned in this study. However another known *osc* gene from *C. asiatica*, namely CabAS (Kim et al., 2005), was cloned using local plant. The obtained cDNA was named as CAK. Three nucleotides of CAK was different from CabAS resulting in 3 amino acids different between both clones. Three amino acids include V631G, V645A and F692L. The functional analysis of CAK in mutant yeast was revealed that CAK codes for multifunctional triterpene synthase, which produced α -and β -amyrin as major product, while the CabAS was reported as dammarenediol synthase (Kim et al., 2009)