



รายงานวิจัยฉบับสมบูรณ์

โครงการ: การยับยั้งการส่งสัญญาณภายในเซลล์เนื่องจาก
การกระตุ้น STAT3 โดย IL-6 ในเซลล์มะเร็งท่อน้ำดีของมนุษย์
เป็นเป้าหมายการออกฤทธิ์ของเคมีต้านมะเร็ง

โดย ลัดดาวัลย์ เส็งกันไพร และคณะ

มิถุนายน 2556

รายงานวิจัยฉบับสมบูรณ์

โครงการ: การยับยั้งการส่งสัญญาณภายในเซลล์เนื่องจาก
การกระตุ้น STAT3 โดย IL-6 ในเซลล์มะเร็งท่อน้ำดีของมนุษย์
เป็นเป้าหมายการออกฤทธิ์ของเคมีต้านมะเร็ง

ผู้วิจัย

- | | |
|------------------------------|----------------------------------|
| 1. น.ส.ลัดดาวัลย์ เส็งกันไพร | คณะแพทยศาสตร์ มหาวิทยาลัยขอนแก่น |
| 2. นายวีรพล คู่คงวิริยพันธุ์ | คณะแพทยศาสตร์ มหาวิทยาลัยขอนแก่น |
| 3. น.ส.เอื้อมเดือน ประวาฬ | คณะแพทยศาสตร์ มหาวิทยาลัยขอนแก่น |
| 4. นางยุพา คู่คงวิริยพันธุ์ | คณะแพทยศาสตร์ มหาวิทยาลัยขอนแก่น |

สนับสนุนโดยสำนักงานคณะกรรมการการอุดมศึกษา
สำนักงานกองทุนสนับสนุนการวิจัย และ มหาวิทยาลัยขอนแก่น

(ความเห็นในรายงานนี้เป็นของผู้วิจัย สกอ. และ สกว. ไม่จำเป็นต้องเห็นด้วยเสมอไป)

บทคัดย่อ

รหัสโครงการ: MRG5480011

ชื่อโครงการ: การยับยั้งการส่งสัญญาณภายในเซลล์เนื่องจากการกระตุ้น STAT3 โดย IL-6 ในเซลล์มะเร็งท่อน้ำดีของมนุษย์เป็นเป้าหมายการออกฤทธิ์ของเคมีต้านมะเร็ง

ชื่อนักวิจัย และสถาบัน

หัวหน้าโครงการ : ผศ.ดร.ลัดดาวัลย์ เส็งกันไพร คณะแพทยศาสตร์ มหาวิทยาลัยขอนแก่น

ผู้ร่วมวิจัย : 1. รศ.ดร.วีรพล คู่คงวิริยพันธุ์ คณะแพทยศาสตร์ มหาวิทยาลัยขอนแก่น

2. ผศ.ดร.เอื้อมเดือน ประวาฬ คณะแพทยศาสตร์ มหาวิทยาลัยขอนแก่น

3. รศ.ดร.ยุพา คู่คงวิริยพันธุ์ คณะแพทยศาสตร์ มหาวิทยาลัยขอนแก่น

อีเมล: laddas@kku.ac.th

ระยะเวลาโครงการ: 2 ปี (15 มิถุนายน 2554 ถึง 14 มิถุนายน 2556)

บทคัดย่อ:

Quercetin และ EGCG เป็นสารพฤกษเคมีที่ได้จากอาหารธรรมชาติ ซึ่งมีฤทธิ์ต้านการอักเสบและต้านมะเร็ง ในการศึกษาครั้งนี้ ผู้วิจัยทำการตรวจสอบฤทธิ์ของสารสองชนิดนี้ต่อวิถี JAK/ STAT ในเซลล์มะเร็งท่อน้ำดี ซึ่งเป็นมะเร็งที่ร้ายแรงและมีการพยากรณ์โรคแย่ และวิถี JAK/STAT มีบทบาทที่สำคัญต่อกระบวนการอักเสบและการเกิดมะเร็ง ผลการศึกษาพบว่า สาร quercetin และ EGCG มีผลยับยั้งการกระตุ้นวิถี JAK/STAT โดยสาร cytokine ซึ่งแสดงผลโดยมีการลดลงของระดับโปรตีน phosphorylated-STAT1 และ phosphorylated-STAT3 โดยผลการยับยั้งเป็นไปตาม dose สาร cytokine สามารถเหนี่ยวนำให้มีการแสดงออกของยีน inducible nitric oxide synthase (iNOS) และ intercellular adhesion molecule-1 (ICAM-1) เพิ่มขึ้น ซึ่งผลการเหนี่ยวนำนี้ถูกยับยั้งได้ด้วยสาร quercetin และ EGCG นอกจากนี้สาร quercetin และ EGCG ยังมีผลยับยั้งการเจริญเติบโตและยับยั้งการเคลื่อนที่ของเซลล์มะเร็งท่อน้ำดีที่เหนี่ยวนำโดยสาร cytokine ด้วย การศึกษาผลของสารยับยั้งวิถี JAK/STAT แบบจำเพาะได้แก่ AG490 และ piceatannol พบว่าสารทั้งสองนี้มีฤทธิ์ยับยั้งการแสดงออกของยีน iNOS และ ICAM-1 ที่เหนี่ยวนำโดย cytokine ได้ จากผลการศึกษาทั้งหมดแสดงให้เห็นถึงประโยชน์ของสาร quercetin และ EGCG ในการออกฤทธิ์ยับยั้งวิถี JAK/STAT ในเซลล์มะเร็งท่อน้ำดี ดังนั้นสารสองชนิดนี้จึงเป็นสารที่มีประสิทธิภาพในการนำมาใช้เป็นเคมีป้องกันการเกิดมะเร็งท่อน้ำดี

คำหลัก : JAK/STAT, quercetin, EGCG, มะเร็งท่อน้ำดี, เคมีป้องกัน

Abstract

Project Code : MRG5480011

Project Title : Suppression of IL-6-mediated STAT3 signaling cascade in human cholangiocarcinomas is a target of anti-cancer drugs

Investigator :

Principal investigator: Assist.Prof. Laddawan Senggunprai,
Faculty of Medicine, Khon Kaen University

Co-investigators: 1. Assoc.Prof. Veerapol Kukongviriyapan,
Faculty of Medicine, Khon Kaen University

2. Assist.Prof. Auemduan Prawan,
Faculty of Medicine, Khon Kaen University

3. Assoc.Prof. Upa Kukongviriyapan,
Faculty of Medicine, Khon Kaen University

E-mail Address : laddas@kku.ac.th

Project Period : 2 years (June 15, 2011 – June 14, 2013)

Abstract:

Quercetin and epigallocatechin-3-gallate (EGCG) are dietary phytochemicals with anti-inflammatory and antitumor effects. In the present study we examined effects of these two compounds on JAK/STAT pathway of cholangiocarcinoma (CCA) cells, because CCA is one of the aggressive cancers with very poor prognosis and JAK/STAT pathway is critically important in inflammation and carcinogenesis. The results showed that the JAK/STAT pathway activation by proinflammatory cytokine interleukin-6 and interferon- γ in CCA cells was suppressed by pretreatment with quercetin and EGCG, evidently by decrease of the elevated phosphorylated-STAT1 and -STAT3 proteins in a dose-dependent manner. The cytokine-mediated up-regulation of inducible nitric oxide synthase (iNOS) and intercellular adhesion molecule-1 (ICAM-1) via JAK/STAT cascade was abolished by both quercetin and EGCG pretreatment. Moreover these flavonoids also could inhibit growth and cytokine-induced migration of CCA cells. Pretreatment with specific JAK inhibitors, AG490 and piceatannol, abolished cytokine-induced iNOS and ICAM-1 expression. These results demonstrate beneficial effects of quercetin and EGCG in suppression of JAK/STAT cascade of CCA cells. Quercetin and EGCG would be potentially useful as cancer chemopreventive agents against CCA.

Keywords : JAK/STAT, quercetin, EGCG, cholangiocarcinoma, chemopreventive

เนื้อหางานวิจัย

1. บทคัดย่อ

Quercetin และ EGCG เป็นสารพฤกษเคมีที่ได้จากอาหารธรรมชาติ ซึ่งมีฤทธิ์ต้านการอักเสบและต้านมะเร็ง ในการศึกษาครั้งนี้ ผู้วิจัยทำการตรวจสอบฤทธิ์ของสารสองชนิดนี้ต่อวิถี JAK/ STAT ในเซลล์มะเร็งท่อน้ำดี ซึ่งเป็นมะเร็งที่ร้ายแรงและมีการพยากรณ์โรคแย่ และวิถี JAK/STAT มีบทบาทที่สำคัญต่อกระบวนการอักเสบและการเกิดมะเร็ง ผลการศึกษาพบว่า สาร quercetin และ EGCG มีผลยับยั้งการกระตุ้นวิถี JAK/STAT โดยสาร cytokine ซึ่งแสดงผลโดยมีการลดลงของระดับโปรตีน phosphorylated-STAT1 และ phosphorylated-STAT3 โดยผลการยับยั้งเป็นไปตาม dose สาร cytokine สามารถเหนี่ยวนำให้มีการแสดงออกของยีน inducible nitric oxide synthase (iNOS) และ intercellular adhesion molecule-1 (ICAM-1) เพิ่มขึ้น ซึ่งผลการเหนี่ยวนำนี้ถูกยับยั้งได้ด้วยสาร quercetin และ EGCG นอกจากนี้สาร quercetin และ EGCG ยังมีผลยับยั้งการเจริญเติบโตและยับยั้งการเคลื่อนที่ของเซลล์มะเร็งท่อน้ำดีที่เหนี่ยวนำโดยสาร cytokine ด้วย การศึกษาผลของสารยับยั้งวิถี JAK/STAT แบบจำเพาะได้แก่ AG490 และ piceatannol พบว่าสารทั้งสองนี้มีฤทธิ์ยับยั้งการแสดงออกของยีน iNOS และ ICAM-1 ที่เหนี่ยวนำโดย cytokine ได้ จากผลการศึกษาทั้งหมดแสดงให้เห็นถึงประโยชน์ของสาร quercetin และ EGCG ในการออกฤทธิ์ยับยั้งวิถี JAK/STAT ในเซลล์มะเร็งท่อน้ำดี ดังนั้นสารสองชนิดนี้จึงเป็นสารที่มีประสิทธิภาพในการนำมาใช้เป็นเคมีป้องกันการเกิดมะเร็งท่อน้ำดี

Quercetin and epigallocatechin-3-gallate (EGCG) are dietary phytochemicals with anti-inflammatory and antitumor effects. In the present study we examined effects of these two compounds on JAK/STAT pathway of cholangiocarcinoma (CCA) cells, because CCA is one of the aggressive cancers with very poor prognosis and JAK/STAT pathway is critically important in inflammation and carcinogenesis. The results showed that the JAK/STAT pathway activation by proinflammatory cytokine interleukin-6 and interferon- γ in CCA cells was suppressed by pretreatment with quercetin and EGCG, evidently by decrease of the elevated phosphorylated-STAT1 and -STAT3 proteins in a dose-dependent manner. The cytokine-mediated up-regulation of inducible nitric oxide synthase (iNOS) and intercellular adhesion molecule-1 (ICAM-1) via JAK/STAT cascade was abolished by both quercetin and EGCG pretreatment. Moreover these flavonoids also could inhibit growth and cytokine-induced migration of CCA cells. Pretreatment with specific JAK inhibitors, AG490 and piceatannol, abolished cytokine-induced iNOS and ICAM-1 expression. These results demonstrate beneficial effects of quercetin and EGCG in suppression of JAK/STAT cascade of CCA cells. Quercetin and EGCG would be potentially useful as cancer chemopreventive agents against CCA.

2. EXECUTIVE SUMMARY

Epidemiological studies have shown that the amount of consumption of vegetables and fruits rich in flavonoids inversely associated with the risk of human cancers (Riboli and Norat, 2003). The anti-inflammatory effect of flavonoids is recognized as an important action for their cancer chemoprevention property (Garcia-Lafuente *et al.*, 2009). Quercetin, the most abundant flavonoid in the human diet, and epigallocatechin-3-gallate (EGCG), the biologically most active constituent in green tea, are proposed to have cancer-preventive activities (Hemalswarya and Doble, 2006). The potential chemopreventive effects of these two compounds have been attributed to various mechanisms including anti-oxidative activity as well as capability to modulate cellular signal transduction pathways involving anti-inflammatory, anti-proliferative and induction of apoptosis of tumor cells (Shanmugam *et al.*, 2011). Quercetin and EGCG can interact with non-receptor protein kinases and receptor tyrosine kinases such as epidermal growth factor receptor and vascular endothelial growth factor receptor (Shanmugam *et al.*, 2011). At present, quercetin and EGCG have been used in clinical trials for prevention of several types of cancers such as breast, lung and prostate cancers (<http://www.clinicaltrials.gov/>).

Cholangiocarcinoma (CCA) is a malignant epithelial neoplasm of the biliary tree with very poor prognosis. It is a rare type of cancer worldwide, however populations residing in the Southeast Asian region are at very high risk. The high incidence of CCA in this region is associated with background conditions particularly liver fluke infection (i.e. *Opisthorchis viverrini* and *Chlonorchis sinensis*) that causes long-standing inflammation, cell injury, and reparative biliary epithelial cell proliferation (Sripa and Pairojkul, 2008). Persistent inflammation can create a local environment enriched with cytokines and other growth factors that primes for cells to develop autonomous proliferative signaling and enhanced production of mitogenic factors (Wise *et al.*, 2008). Excretory/secretory products of liver flukes can stimulate epithelial bile duct cells to secrete proinflammatory cytokine interleukin-6 (IL-6) (Ninlawan *et al.*, 2010). In addition, serum interferon-gamma (IFN- γ) and its expression level in the liver of mice were increased by *Opisthorchis viverrini* infection (Nair *et al.*, 2011). These suggest that various inflammatory mediators, particularly proinflammatory cytokines, are common and important contributors for cholangiocarcinogenesis.

Proinflammatory cytokines activate the signal transducers and activation of transcription (STAT) proteins, which are considered to be oncogenic transcription factors (Hodge *et al.*, 2005). Stimuli to cell surface cytokine receptors activates the Janus-like kinase (JAK) family of protein kinases, which, in turn phosphorylates and activates latent cytoplasmic STAT proteins to an active dimer, leading to nuclear translocation, DNA binding and subsequently modulating gene transcription (Darnell, 1997). Several other kinases including members of the Src and Abl family have also been implicated in the phosphorylation of STATs (Darnell, 1997). Among seven known mammalian STAT proteins, STAT1 and STAT3 are the most widely studied isoforms due to their constitutively activated state in many tumors but not in normal cells (Hodge *et al.*, 2005). STAT1 is relatively specific to IFNs and a pivotal transcription factor in IFN- γ -induced expression of inflammatory genes, whereas STAT3 is mainly activated by IL-6 and other gp130-related cytokines (Yoshimura, 2006).

The crucial roles of STATs, particularly STAT1 and STAT3, in inflammation and tumorigenesis have been demonstrated in several studies such as inflammation-associated gastric

tumorigenesis (Ernst *et al.*, 2008) and colitis-associated tumorigenesis (Grivennikov *et al.*, 2009). The activation of JAK/STAT cascade triggered by inflammatory cytokines produced by tumor infiltrating immune cells can lead to the expression of STAT-regulated genes including proinflammatory enzymes and proteins in which then enhance inflammatory condition (Dalwadi *et al.*, 2005). In addition, the activation of JAK/STAT pathway also results in induction of genes that mediate cell proliferation, suppression of apoptosis and promotion of angiogenesis (Barton *et al.*, 2004).

Presently there is neither effective treatment for patients with the advanced stage of CCA nor any effective agents for chemoprevention of CCA. Similar to several cancers, the JAK/STAT signaling pathway is involved in the development of CCA (Isomoto *et al.*, 2007). Thus, STAT signaling pathway inhibitors may be useful for prevention and treatment of CCA. Currently, there is only few studies addressed the role of JAK/STAT signaling pathway in carcinogenesis of CCA.

In the present study, we have investigated effects of quercetin and EGCG on JAK/STAT pathway using CCA cells treated with proinflammatory cytokine mixture, IL-6 and IFN- γ , for mimicking inflammatory condition (Kaur *et al.*, 2003). The results show that both quercetin and EGCG could inhibit JAK/STAT signaling cascade. They also could suppress cytokine-induced the expression of inducible nitric oxide synthase (iNOS) and intercellular adhesion molecule-1 (ICAM-1), the key molecules involving in inflammatory and tumorigenesis processes. Moreover, they inhibited growth and migration of CCA cells. Our findings provide new insights into the intervention strategy for prevention of CCA.

3. OBJECTIVES

- 3.1 To investigate the role of JAK/STAT pathway in cholangiocarcinoma cells
- 3.2 To examine the effects of quercetin and EGCG on JAK/STAT pathway in cholangiocarcinoma cells
- 3.3 To determine the effects of quercetin and EGCG on STAT-related genes
- 3.4 To determine the effects of quercetin and EGCG on CCA cell growth

4. MATERIALS AND METHODS

4.1 Reagents

Quercetin, EGCG and piceatannol were purchased from Sigma Chemical (St.Louis, MO, USA). AG490 was obtained from Calbiochem (San Diego, CA, USA). Interleukin-6 (IL-6), interferon- γ (IFN- γ) and the primary antibodies against STAT1, STAT3, phospho-STAT1 (Tyr⁷⁰¹) and phospho-STAT3 (Tyr⁷⁰⁵) were purchased from Cell Signaling Technology (Danvers, MA, USA). RIPA buffer was from Amresco (Solon, OH, USA). The pGL4 basic and pGL4[*hRluc*] vector, LipofectamineTM 2000 and Dual-Luciferase[®] Reporter Assay Kit were obtained from Promega (Madison, WI, USA). Reagents for cell culture were from Gibco BRL Life Technologies (Grand Island, NY, USA).

4.2 Cell lines and cell cultures

The human CCA cell lines; KKU100, KKU-M139 and KKU-M213 used in this study were kindly provided by Dr. Banchob Sripa of Department of Pathology, Faculty of Medicine, Khon Kaen University. These cell lines were cultured in complete media consisting of Ham's F12 media,

supplemented with 10% fetal calf serum, 12.5 mM HEPES, pH 7.3, 100 U/mL penicillin G and 100 µg/mL streptomycin and maintained under an atmosphere of 5% CO₂ at 37°C. The cells were subcultured every 3 days using 0.25% trypsin-EDTA.

4.3 Cells treatment

At about 70% confluency, cultures were starved in serum-free medium for 16 h and then treated with cytokine mixture, combination of IL-6 and IFN-γ (Kaur *et al.*, 2003), in serum-free medium for further 1 h to 24 h. The concentration used for both IL-6 and IFN-γ were 100 ng/mL. In experiments involving flavonoids or JAK inhibitors, the cells were pretreated with various concentrations of quercetin (1, 10, 50 and 100 µM) or EGCG (1, 10, 25 and 50 µM) or piceatannol (JAK1 inhibitor) (Barton *et al.*, 2004) (1, 10 and 100 µM) or AG490 (JAK2 inhibitor) (Barton *et al.*, 2004) (1, 10 and 100 µM) in DMSO for 2 h and then further treated with cytokine mixture. An equal amount of DMSO (vehicle) was present in each treatment, including control. DMSO concentration did not exceed 0.1% (v/v) in any treatment.

4.4 Protein extraction and Western blot analysis

After treatments for designated period of times, medium was aspirated, cells were washed twice with cold PBS and whole cell lysates were prepared using RIPA cell lysis buffer (Amresco, OH, USA) according to the manufacturer's instructions. The proteins were resolved by SDS-PAGE using 10% polyacrylamide gel and the separated proteins on the gel were electrophoretically transferred to a PVDF membrane. The membranes were blocked with 5% (w/v) bovine serum albumin in Tris buffered saline containing 0.1% Tween-20 for 2 h at room temperature, followed by incubation with primary antibody against STAT1 (1:1000), phospho-STAT1 (Tyr⁷⁰¹) (1:1000), STAT3 (1:1000), phospho-STAT3 (Tyr⁷⁰⁵) (1:1000) and β-actin (1:5000) at 4°C overnight. After washing, the membranes were incubated with the appropriate horseradish peroxidase conjugated secondary antibody to detect bands by AmershamTM ECLTM Prime (Amersham Biosciences Corp, NJ, USA). The densities of the specific protein bands were visualized and captured by ImageQuantTM 400.

4.5 Real-time polymerase chain reaction

Total RNA was extracted from cells using Trizol reagent (Invitrogen, USA) according to the manufacturer's instructions. One µg of total RNA was then reverse transcribed to single-stranded cDNA by the ImProm-IITM reverse transcription system (Promega, WI, USA). PCR was carried out using specific primers as follows: iNOS forward 5'-GTTCTCACGGCACAGGTCTC-3' and iNOS reverse 5'-GCAGGTCACTTATGTCAC TTATC-3' (Wu *et al.*, 2011), ICAM-1 forward 5'-CAAGGCCTCAGTCAGTGTGA-3' and ICAM-1 reverse 5'-CCTCTGGCTTCGTCAGAATC-3' (Kim *et al.*, 2010), GAPDH forward 5'-GTGGTGGACCTGACCTGC-3' and GAPDH reverse 5'-TGAGCTTGACA AAGTGGTCG-3' (Kim *et al.*, 2010). The relative expression of iNOS and ICAM-1 was analyzed using quantitative RT-PCR with GAPDH as an internal control. The PCR was performed with SsoFastTM EvaGreen Supermix with low Rox (Bio-Rad, CA, USA) using an ABI Prism 7500 Sequence Detection System (Applied Biosystems, Foster City, CA, USA) under the following conditions: denaturation at 95°C for 3 min and amplification by cycling 40 times at 95°C for 15 sec and 60°C for 31 sec.

4.6 Cell growth assay

Sulforhodamine B (SRB) was used to measure the effect of quercetin and EGCG on the proliferation of CCA cell lines as described previously (Prawan *et al.*, 2009). Briefly, KCU100 cells were plated in a 96-well plate for 24 h. After exposure of cultured cells to quercetin or EGCG at various concentrations for 48 h, the culture medium was removed, 10% cold trichloroacetic acid was added for 1 h at 4 °C, and subsequently washed 5 times with deionized H₂O. The plate was air dried, and 0.4% SRB in 1% acetic acid was added for 30 min. Unbound dye was washed out 5 times with 1% acetic acid. After air drying, SRB dye within cells was solubilized with 200 µl of 10 mM unbuffered Tris base solution. The plate was shaken for 10 min, and the absorbance was measured at 540 nm by using a microplate reader. Cell growth inhibition was expressed in terms of percentage of untreated control absorbance following subtraction of mean background absorbance. The IC₅₀ concentration (50% inhibition of cell growth values) was calculated from the dose-response curves.

4.7 Wound healing assay

KCU100 cells (1.5×10^5) were seeded into a 24-well plate and incubated overnight in Ham's F12 medium, supplemented with 10% fetal calf serum. A scratch wound was made with a sterile 200 µL pipette tip. After washing with PBS to remove any detached cell, a series of images of the scratched wound were taken from 0 to 24 h. The cells were pretreated with quercetin (10 µM) or EGCG (5 µM) or vehicle for 2 h, then with cytokine mixture afterward. The closing of scratched wound, represented the migration process, was determined by capture of the denuded area along the scratch using Image-Pro Plus software (Media Cybernetics, LP, USA). The wound distance was calculated by dividing the area by the length of scratch.

4.8 STAT3 luciferase reporter assay

To construct the STAT3 luciferase vector, the STAT3 transcriptional regulatory element 5'-GTCGACATTTCCCGTAAATCGTCGA-3' (Bergad *et al.*, 2000) containing NheI/HindIII restriction sites was subcloned into the multiple cloning site of the pGL4 luciferase reporter plasmid (Promega, WI, USA). The sequences of the constructs were verified by the MegaBACE DNA Analysis System (Amersham Biosciences Corp, NJ, USA). For reporter assay, the cells (1×10^4) were seeded into each well of 96-well plates, and cultured for 24 h. The cells were transfected with 0.2 µg of STAT3-pGL4 reporter vector according to the manufacturer's protocols using XtremeGene HP (Roche) for KCU100 cells or LipofectamineTM 2000 (Promega, WI, USA) for KCU-M139 and KCU-M213 cells. The cells were also cotransfected with 0.01 µg of pGL4[*hRluc*/TK] vector containing the *Renilla luciferase* gene (Promega) to control transfection efficiency. After 6 h of transfection, cells were pretreated with flavonoids for 2 h and then stimulated with the cytokine mixture for additional 18 h before being washed and lysed in luciferase lysis buffer (Promega). Luciferase activity was measured with a luminometer by using a Dual-Luciferase[®] Reporter Assay Kit (Promega).

5. RESULTS

5.1 Cytokine mixture induced STAT1 and STAT3 activation in K KU100 cells

To clarify whether JAK/STAT signaling is intact in CCA cells, the effect of cytokine mixture (100 ng/mL IL-6 and 100 ng/mL IFN- γ) on STAT levels in K KU100 cells was determined at various time points (1 h to 24 h). In separate experiments, we confirmed that the dose range of cytokine mixture used in this study did not affect the cell viability (data not shown). After treatment with the cytokine mixture, phosphorylation of STAT1 (Tyr⁷⁰¹) in K KU100 cells was dramatically increased as early as within the first hour, and then gradually decreased afterwards, although it was still detectable at 24 h (Fig. 1A). For STAT3, the cytokine mixture induced transient STAT3 (Tyr⁷⁰⁵) phosphorylation in K KU100 cells within the first hour, but the phospho-STAT3 was diminished rapidly within 2 h (Fig. 1A). From these results, functional JAK/STAT signaling cascade is present in CCA cells.




Figure 1: Effects of cytokine mixture on the activation of STAT and effects of quercetin and EGCG on cytokine-induced STAT activation. (A) K KU100 cells were incubated with cytokine combination (100 ng/mL IL-6 and 100 ng/mL IFN- γ) for various time points (1 to 24 h). (B-C) K KU100 cells were pretreated with the indicated concentrations of (B) quercetin or (C) EGCG for 2 h and then treated with cytokine mixture for 1 h. Whole cell extracts were prepared and 20 μ g aliquots of those extracts were resolved on 10% SDS-PAGE, electrotransferred onto PVDF membranes and probed using the indicated antibodies as described under Materials and Methods. Anti- β -actin antibody was used as a loading control. Results shown are representative of at least two reproducible independent experiments.

5.2 Quercetin and EGCG inhibited cytokine-induced STAT1 and STAT3 phosphorylation

To determine whether quercetin and EGCG can suppress the proinflammatory cytokine-induced JAK/STAT phosphorylation, KKU100 cells were pre-incubated with various concentrations of quercetin or EGCG for 2 h and then stimulated with the cytokine mixture for 1 h. Under these conditions, the cell viability was not affected (data not shown). As shown in Fig. 1, when KKU100 cells were pretreated with quercetin (Fig. 1B) or EGCG (Fig. 1C), cytokine-induced STAT1 and STAT3 phosphorylation was markedly suppressed in a dose-dependent manner, although STAT1 and STAT3 protein levels were not affected.

5.3 Quercetin and EGCG suppressed cytokine mixture-induced iNOS and ICAM-1 expression

It is known that iNOS and ICAM-1 play a critical role in inflammatory and tumorigenesis processes (Yamada *et al.*, 2006; Kostourou *et al.*, 2011). Therefore, we first examined the effect of cytokine treatment on the expression of iNOS and ICAM-1 in CCA cells using real time-PCR. The results showed that cytokine treatment markedly up-regulated the mRNA expression of iNOS and ICAM-1 (Fig. 2). Pretreatment with quercetin and EGCG significantly inhibited cytokine-mediated up-regulation of iNOS and ICAM-1 expression in a dose-dependent manner (Fig. 2A and 2B, respectively). The inhibitory effect of EGCG was more potent than that of quercetin.




Figure 2: Effects of quercetin and EGCG on cytokine-induced expression of iNOS and ICAM-1. KKU100 cells were pretreated with various concentrations of quercetin (1 - 100 μM) or EGCG (1 - 50 μM) for 2 h and then treated with cytokine combination (100 ng/mL IL-6 and 100 ng/mL IFN-γ) for 3 h. At the end of treatments, total RNA of cells were collected and real time RT-PCR was carried out for (A) iNOS and (B) ICAM-1 expression. The bars represent relative expression of indicated genes normalized with GAPDH. The data are expressed as mean ± SD, each from three independent experiments. *, $p < 0.05$ vs cytokine treatment alone.

5.4 JAK1 and JAK2 inhibition attenuated cytokine-induced expression of iNOS and ICAM-1

To validate whether JAK/STAT signaling contributes to the up-regulation of iNOS and ICAM-1, effects of piceatannol (JAK1 inhibitor) and AG490 (JAK2 inhibitor) on the cytokine-induced STAT1 and STAT3 phosphorylation in KKKU100 cells were examined. As illustrated in Fig. 3A, pretreatment with these inhibitors could abrogate STAT1 and STAT3 phosphorylation in a dose-dependent manner. Subsequently, KKKU100 cells were pre-incubated with various concentration of piceatannol or AG490 for 2 h and then treated with the cytokine mixture for 3 h. After that the mRNA expression of iNOS and ICAM-1 were evaluated. The results showed that cytokine-induced iNOS expression in KKKU100 cells were suppressed dose-dependently by piceatannol and AG490 pretreatment (Fig. 3B). Similarly, piceatannol and AG490 could suppress the up-regulation of ICAM-1 expression, although the higher concentration was required for the inhibition compared to the effects on iNOS expression (Fig. 3C). Under these conditions, the cell viability was only slightly reduced (data not shown).




Figure 3: Effects of JAK inhibitors on cytokine-mediated phosphorylation of STAT and -induced iNOS and ICAM-1 expression. (A) KKKU100 cells were pretreated with the indicated concentrations of AG490 or piceatannol for 2 h and then treated with cytokine mixture for 1 h. Whole cell extracts were prepared and 20 μ g aliquots of those extracts were resolved on 10% SDS-PAGE, electrotransferred onto PVDF membranes and probed using the indicated antibodies as described under Materials and Methods. Anti- β -actin antibody was used as a loading control. Results shown are representative of at least two reproducible independent experiments. (B-C) KKKU100 cells were pretreated with indicated concentrations (1 - 100 μ M) of AG490 or piceatannol for 2 h and then treated with cytokine combination (100 ng/mL IL-6 and 100 ng/mL IFN- γ) for 3 h. At the end of treatments, total RNA of cells were collected and analyzed for (B) iNOS and (C) ICAM-1 expression by real time RT-PCR. The bars represent relative expression of indicated genes normalized with GAPDH. The data are expressed as mean $\pm$ SD, each from three independent experiments. *, $p < 0.05$ vs cytokine treatment alone.

5.5 Quercetin and EGCG inhibited growth and migration of CCA cell lines

The anticancer effect of quercetin and EGCG was further evaluated by examining some downstream effects of JAK/STAT signaling i.e. anti-proliferation and anti-migration. The anti-proliferative activity was assayed on KKU100 cells using SRB assay. Quercetin and EGCG had similar potency to inhibit KKU100 cell growth with the IC_{50} values of 31.1 ± 7.1 and 24.3 ± 3.1 μ M, respectively (Fig. 4A).

Since quercetin and EGCG showed strong inhibitory effect on cytokine-induced expression of ICAM-1, an adhesion molecule involved in cell migration and invasion, the effect of these two compounds on CCA cell migration was investigated by wound healing assay using the monolayer culture of KKU100 cells. The results are shown in Fig. 4B and 4C. Closure of the scratched wound was stimulated by cytokine mixture, determined at 24 h. The cytokine-mediated stimulation of cell migration was suppressed by quercetin and EGCG, as the migration distance of cultured cells in those flavonoid treated groups was shorter than that of cytokine treatment only.




Figure 4: Effects of quercetin and EGCG on the growth and cytokine-induced migration of CCA cells. (A) KKU100 cells were treated with indicated concentration of quercetin or EGCG for 48 h and cell numbers were determined by SRB assay. The results are presented as percentage of control. The data are the mean $\pm$ SD averaged from three independent experiments. (B) Scratched wounds of monolayer KKU100 cells were pretreated with vehicle or 10 μ M of quercetin or 5 μ M of EGCG for 2 h and then treated with cytokine mixture. Cell migration was monitored under phase-contrast microscopy (x4 magnification). Representative images of wound healing were obtained at the time of the scratch and 24 h later. (C) The graph shows the level of cell migration into the wound scratch quantified as the percentage of wound closure at 24 h. The data are mean $\pm$ SD averaged from quadruplets of one experiment. CK: cytokine, QC: quercetin.

5.6 Quercetin and EGCG inhibited STAT3-dependent luciferase activity in CCA cells

The suppressive effects of quercetin and EGCG on JAK/STAT signaling were confirmed further by STAT3-luciferase reporter assay. Firstly, we demonstrated that quercetin and EGCG suppressed cytokine-induced STAT3 luciferase activity in KKU100 cells (Fig. 5A). To explore whether the suppressive effects of flavonoids on STAT3 signaling is common in various CCA cells, in addition to KKU100 cells, KKU-M139 and KKU-M213 cells were employed in this experiment. These three human CCA cell lines were established from primary tumors of liver fluke-associated CCA patients with different histological types i.e. poorly differentiated, adeno-squamous, and mixed papillary and non-papillary adenocarcinoma, respectively. As was expected, cytokine-induced STAT3 activity in KKU-M139 and KKU-M213 cells were strongly suppressed by quercetin and EGCG as shown in Fig. 5B and 5C, respectively, suggesting the effect of the flavonoids are not restricted to a special cell type.




Figure 5: Effects of quercetin and EGCG on cytokine-induced transcriptional activity of STAT3. The CCA cells: (A) KKU100 (B) KKU-M139 and (C) KKU-M213 were transfected with STAT3-pGL4 reporter vector together with renilla luciferase plasmid. After 6 h of transfection period, cells were pretreated with indicated concentrations of quercetin or EGCG for 2 h before exposure to cytokine combination (100 ng/mL IL-6 and 100 ng/mL IFN- γ) for additional 18 h. After that the luciferase activity was measured and presented as fold induction to the control cells without cytokine treatment. Each bar indicates the mean $\pm$ SD of five to six determinations. *, $p < 0.05$ vs cytokine treatment alone.

6. DISCUSSION

Chemoprevention and chemotherapy of CCA are so far quite disappointing. Several efforts have been made to identify and develop new agents for both treatment and prevention of this deadly disease. These include targeted therapy on some critical molecules of cancer cells, such as NQO1 and HO-1, where inhibition of the enzymes make CCA cells sensitive to chemotherapeutic agents (Buranrat *et al.*, 2010; Kongpetch *et al.*, 2012). In the present study, we have demonstrated that JAK/STAT signaling pathway may be an important target for inhibition of CCA. JAK/STAT signaling plays an important role in the expression of iNOS and ICAM-1, both of which are known critical molecules involving in inflammation and carcinogenesis processes. Quercetin and EGCG inhibited JAK/STAT signaling cascade in CCA cells leading to the down-regulation of iNOS and ICAM-1 and suppression of cell migration. In addition, these two phytochemical compounds could also inhibit the growth of CCA cells. The results provide evidence supporting the use of these two flavonoids as chemopreventive agents for CCA.

Quercetin and EGCG are potent inhibitors of tumor growth and inflammation and have traditionally been used to treat many inflammatory disorders (Garcia-Lafuente *et al.*, 2009). Quercetin and EGCG have been shown to exert their effects through inhibiting several signaling pathway such as NF- κ B (Priyadarsini and Nagini, 2012), AP-1 (Shanmugam *et al.*, 2011) and JAK/STAT signaling pathway (Tedeschi *et al.*, 2002; Michaud-Levesque *et al.*, 2012). We also found that quercetin and EGCG are potent inhibitors of cytokine-induced STAT1 as well as STAT3 activation in CCA cells.

In the present study, we demonstrated that the activation of JAK/STAT pathway is associated with the increase in expression of iNOS and ICAM-1. The iNOS has critical functions in inflammation-related diseases. Sustained and excess nitric oxide generation, most of which is attributable to iNOS expression, often is pathogenic to cause inflammation, angiogenesis and neoplasia (Kostourou *et al.*, 2011). Quercetin as well as EGCG are known to exhibit suppression of iNOS expression in various cancer cells (Gerhauser *et al.*, 2003). It has been reported that iNOS gene promoter has binding sites for STAT, AP-1, NF- κ B and HIF-1 α (Chittezhath *et al.*, 2008). In this study, we demonstrated that cytokine-induced STAT activation is an important determinant of iNOS expression in CCA cells, as JAK inhibitors markedly inhibited the expression. The inhibition of STAT activation by quercetin correlated with the decreases in iNOS mRNA levels. Similar inhibitory patterns were observed after EGCG treatment, and the inhibitory effect of EGCG was stronger than quercetin. These results indicate that the inhibitory effects of quercetin and EGCG on cytokine-induced iNOS expression are mediated partly via suppression of JAK/STAT pathway. Because no other pathways were tested in the present study, further studies are required to elucidate the action of quercetin and EGCG on the other signaling pathways involving in iNOS expression.

ICAM-1, of which expression is regulated by JAK/STAT pathway, plays a critical role in inflammation-related diseases by promoting trafficking of leukocytes across endothelia and epithelial barriers. It also implicated in carcinogenic processes by contributing in tumor metastasis (Yamada *et al.*, 2006). The modulation of ICAM-1 expression is, therefore, an important target for chemoprevention and chemotherapy, as shown by the beneficial effect of ICAM-1 inhibitors on the tumorigenesis and cancer progression (Lee *et al.*, 2012). Similar to iNOS, transcription factors

important to the activation of ICAM-1 expression include AP-1, NF- κ B and STAT (van de Stolpe and van der Saag, 1996).

In this study, we found that although AG490 at 100 μ M profoundly suppressed the phosphorylation of STAT1 and STAT3, this treatment caused a significant but small suppression on ICAM-1 expression. These results suggest that ICAM-1 expression may be modulated by multiple signaling pathways. The suppressive effect of piceatannol on ICAM-1 expression was also incomplete. It should be noted that piceatannol may inhibit NF- κ B which is also involved in the regulation of ICAM-1 expression (Ashikawa *et al.*, 2002). In this study we found that quercetin and EGCG both dramatically suppress cytokine-induced ICAM-1 expression, where EGCG have higher inhibitory effect than quercetin. Quercetin and EGCG exert their suppressive effects on cytokine-induced ICAM-1 up-regulation, at least in part, through modulating the JAK/STAT cascade in CCA cells.

Enhanced cell migration and invasion are important aspects of cancer phenotypes, where ICAM-1 plays an important role (Yamada *et al.*, 2006). In this study, we found that cytokine-induced migration of CCA cells was associated with ICAM-1 expression. Quercetin and EGCG could significantly abolish the cytokine-dependent CCA cell migration. It should be noted that the anti-migratory effect of flavonoids was detected at low concentrations which minimally inhibited cell growth. In fact, the modulation of cell migration ability is a target for chemopreventive compound. Thus, our findings provide a supportive evidence that quercetin and EGCG are potentially useful as cancer chemopreventive agents for CCA.

To ascertain that the inhibition of JAK/STAT signaling by quercetin and EGCG is not confined to only one cell type, we performed the reporter assay with other two CCA cell lines. It is clearly demonstrated that both flavonoids inhibited cytokine-induced STAT3 luciferase activity in all three CCA cell lines examined. These results imply that quercetin and EGCG may efficiently inhibit JAK/STAT signaling pathway in most CCA cells and this confers a high potential therapeutic opportunity.

In summary, the present study showed that JAK/STAT signaling cascade is functioning in CCA cells and upon activation it induces the expression of iNOS and ICAM-1. Quercetin and EGCG can suppress STAT1- and STAT3-phosphorylation and also suppress iNOS and ICAM-1 expression. Moreover, these two compounds can inhibit cytokine-induced CCA cell migration as well. Given that JAK/STAT cascade plays a critical role in the inflammation and carcinogenesis processes, inhibition of JAK/STAT signaling pathway could be a valuable chemopreventive target of natural compounds to protect against inflammation-associated cancer.

REFERENCES

- Ashikawa K, Majumdar S, Banerjee S, Bharti AC, Shishodia S, Aggarwal BB. 2002. Piceatannol inhibits TNF-induced NF- κ B activation and NF- κ B-mediated gene expression through suppression of IkappaB α kinase and p65 phosphorylation. *J Immunol* **169**(11): 6490-6497.
- Barton BE, Karras JG, Murphy TF, Barton A, Huang HF. 2004. Signal transducer and activator of transcription 3 (STAT3) activation in prostate cancer: Direct STAT3 inhibition induces apoptosis in prostate cancer lines. *Mol Cancer Ther* **3**(1):11-20.

- Bergad PL, Schwarzenberg SJ, Humbert JT, Morrison M, Amarasinghe S, Towle HC, *et al.* 2000. Inhibition of growth hormone action in models of inflammation. *Am J Physiol Cell Physiol* **279**(6): C1906-1917.
- Buranrat B, Prawan A, Kukongviriyapan U, Kongpetch S, Kukongviriyapan V. 2010. Dicoumarol enhances gemcitabine-induced cytotoxicity in high NQO1-expressing cholangiocarcinoma cells. *World J Gastroenterol* **16**(19): 2362-2370.
- Chittezhath M, Deep G, Singh RP, Agarwal C, Agarwal R. 2008. Silibinin inhibits cytokine-induced signaling cascades and down-regulates inducible nitric oxide synthase in human lung carcinoma A549 cells. *Mol Cancer Ther* **7**(7): 1817-1826.
- Dalwadi H, Krysan K, Heuze-Vourc'h N, Dohadwala M, Elashoff D, Sharma S, *et al.* 2005. Cyclooxygenase-2-dependent activation of signal transducer and activator of transcription 3 by interleukin-6 in non-small cell lung cancer. *Clin Cancer Res* **11**(21): 7674-7682.
- Darnell JE, Jr. 1997. STATs and gene regulation. *Science* **277**(5332): 1630-1635.
- Ernst M, Najdovska M, Grail D, Lundgren-May T, Buchert M, Tye H, *et al.* 2008. STAT3 and STAT1 mediate IL-11-dependent and inflammation-associated gastric tumorigenesis in gp130 receptor mutant mice. *J Clin Invest* **118**(5): 1727-1738.
- Garcia-Lafuente A, Guillaumon E, Villares A, Rostagno MA, Martinez JA. 2009. Flavonoids as anti-inflammatory agents: implications in cancer and cardiovascular disease. *Inflamm Res* **58**(9): 537-552.
- Gerhauser C, Klimo K, Heiss E, Neumann I, Gamal-Eldeen A, Knauff J, *et al.* 2003. Mechanism-based in vitro screening of potential cancer chemopreventive agents. *Mutat Res* **523-524**: 163-172.
- Grivennikov S, Karin E, Terzic J, Mucida D, Yu GY, Vallabhapurapu S, *et al.* 2009. IL-6 and Stat3 are required for survival of intestinal epithelial cells and development of colitis-associated cancer. *Cancer Cell* **15**(2): 103-113.
- Hemalswarya S, Doble M. 2006. Potential synergism of natural products in the treatment of cancer. *Phytother Res* **20**(4): 239-249.
- Hodge DR, Hurt EM, Farrar WL. 2005. The role of IL-6 and STAT3 in inflammation and cancer. *Eur J Cancer* **41**(16): 2502-2512.
- Isomoto H, Mott JL, Kobayashi S, Werneburg NW, Bronk SF, Haan S, *et al.* 2007. Sustained IL-6/STAT-3 signaling in cholangiocarcinoma cells due to SOCS-3 epigenetic silencing. *Gastroenterology* **132**(1): 384-396.
- Kaur N, Kim IJ, Higgins D, Halvorsen SW. 2003. Induction of an interferon-gamma Stat3 response in nerve cells by pre-treatment with gp130 cytokines. *J Neurochem* **87**(2): 437-447.
- Kim YS, Im J, Choi JN, Kang SS, Lee YJ, Lee CH, *et al.* 2010. Induction of ICAM-1 by *Armillariella mellea* is mediated through generation of reactive oxygen species and JNK activation. *J Ethnopharmacol* **128**(1): 198-205.
- Kongpetch S, Kukongviriyapan V, Prawan A, Senggunprai L, Kukongviriyapan U, Buranrat B. 2012. Crucial role of heme oxygenase-1 on the sensitivity of cholangiocarcinoma cells to chemotherapeutic agents. *PLoS One* **7**(4): e34994.
- Kostourou V, Cartwright JE, Johnstone AP, Boulton JK, Cullis ER, Whitley G, *et al.* 2011. The role of tumour-derived iNOS in tumour progression and angiogenesis. *Br J Cancer* **104**(1): 83-90.
- Lee IT, Lin CC, Lee CY, Hsieh PW, Yang CM. 2012. Protective effects of (-)-epigallocatechin-3-gallate against TNF- α -induced lung inflammation via ROS-dependent ICAM-1 inhibition. *J Nutr Biochem* **68**(3): 946-955.

- Michaud-Levesque J, Bousquet-Gagnon N, Beliveau R. 2012. Quercetin abrogates IL-6/STAT3 signaling and inhibits glioblastoma cell line growth and migration. *Exp Cell Res* **318**(8): 925-935.
- Nair SS, Bommana A, Pakala SB, Ohshiro K, Lyon AJ, Suttiprapa S, *et al.* 2011. Inflammatory response to liver fluke *Opisthorchis viverrini* in mice depends on host master coregulator MTA1, a marker for parasite-induced cholangiocarcinoma in humans. *Hepatology* **54**(4): 1388-1397.
- Ninlawan K, O'Hara SP, Splinter PL, Yongvanit P, Kaewkes S, Surapaitoon A, *et al.* 2010. *Opisthorchis viverrini* excretory/secretory products induce toll-like receptor 4 upregulation and production of interleukin 6 and 8 in cholangiocyte. *Parasitol Int* **59**(4): 616-621.
- Prawan A, Buranrat B, Kukongviriyapan U, Sripa B, Kukongviriyapan V. 2009. Inflammatory cytokines suppress NAD(P)H:quinone oxidoreductase-1 and induce oxidative stress in cholangiocarcinoma cells. *J Cancer Res Clin Oncol* **135**(4): 515-522.
- Priyadarsini RV, Nagini S. 2012. Quercetin suppresses cytochrome P450 mediated ROS generation and NFkappaB activation to inhibit the development of 7,12-dimethylbenz[a]anthracene (DMBA) induced hamster buccal pouch carcinomas. *Free Radic Res* **46**(1): 41-49.
- Riboli E, Norat T. 2003. Epidemiologic evidence of the protective effect of fruit and vegetables on cancer risk. *Am J Clin Nutr* **78**(3 Suppl): 559S-569S.
- Shanmugam MK, Kannaiyan R, Sethi G. 2011. Targeting cell signaling and apoptotic pathways by dietary agents: role in the prevention and treatment of cancer. *Nutr Cancer* **63**(2): 161-173.
- Sripa B, Pairojkul C. 2008. Cholangiocarcinoma: lessons from Thailand. *Curr Opin Gastroenterol* **24**(3): 349-356.
- Tedeschi E, Suzuki H, Menegazzi M. 2002. Antiinflammatory action of EGCG, the main component of green tea, through STAT-1 inhibition. *Ann N Y Acad Sci* **973**: 435-437.
- van de Stolpe A, van der Saag PT. 1996. Intercellular adhesion molecule-1. *J Mol Med (Berl)* **74**(1): 13-33.
- Wise C, Pилanthanonond M, Perry BF, Alpini G, McNeal M, Glaser SS. 2008. Mechanisms of biliary carcinogenesis and growth. *World J Gastroenterol* **14**(19): 2986-2999.
- Wu SQ, Otero M, Unger FM, Goldring MB, Phrutivorapongkul A, Chiari C, *et al.* 2011. Anti-inflammatory activity of an ethanolic *Caesalpinia sappan* extract in human chondrocytes and macrophages. *J Ethnopharmacol* **138**(2): 364-372.
- Yamada M, Yanaba K, Hasegawa M, Matsushita Y, Horikawa M, Komura K, *et al.* 2006. Regulation of local and metastatic host-mediated anti-tumour mechanisms by L-selectin and intercellular adhesion molecule-1. *Clin Exp Immunol* **143**(2): 216-227.
- Yoshimura A. 2006. Signal transduction of inflammatory cytokines and tumor development. *Cancer Sci* **97**(6): 439-447.

7. Appendix

- The revised manuscript submitted to Editor of Phytotherapy Research Journal
- The poster presented a part of the study at 13th International Congress of the Society for Ethnopharmacology, Austria on September 2-6, 2012.

Appendix

Quercetin and EGCG exhibit chemopreventive effects in cholangiocarcinoma cells via suppression of JAK/STAT signaling pathway

Journal:	<i>Phytotherapy Research</i>
Manuscript ID:	PTR-12-1189.R1
Wiley - Manuscript type:	Full Paper
Date Submitted by the Author:	03-Jun-2013
Complete List of Authors:	Senggunprai, Laddawan; Khon Kaen University, Department of Pharmacology, Faculty of Medicine, and Liver Fluke and Cholangiocarcinoma Research Center Kukongviriyapan, Veerapol; Khon Kaen University, Department of Pharmacology, Faculty of Medicine, and Liver Fluke and Cholangiocarcinoma Research Center Prawan, Auemduan; Khon Kaen University, Department of Pharmacology, Faculty of Medicine, and Liver Fluke and Cholangiocarcinoma Research Center Kukongviriyapan, Upa; Khon Kaen University, Department of Physiology, Faculty of Medicine
Keyword:	JAK/STAT, quercetin, EGCG, cholangiocarcinoma, phytochemical, chemopreventive

SCHOLARONE™
Manuscripts

Title page

Quercetin and EGCG exhibit chemopreventive effects in cholangiocarcinoma cells via suppression of JAK/STAT signaling pathway

Laddawan Senggunprai¹, Veerapol Kukongviriyapan¹, Auemduan Prawan¹ and
Upa Kukongviriyapan²

¹Department of Pharmacology, Faculty of Medicine, and Liver Fluke and
Cholangiocarcinoma Research Center, Khon Kaen University, Thailand 40002

²Department of Physiology, Faculty of Medicine, Khon Kaen University, Thailand 40002

Corresponding author: Laddawan Senggunprai, Department of Pharmacology, Faculty of
Medicine, Khon Kaen University, Khon Kaen, Thailand 40002. Tel: +66-43-348397; Fax:
+66-43-348397; e-mail: laddas@kku.ac.th, laddanok@yahoo.com

Short title: Quercetin and EGCG inhibit JAK/STAT in cholangiocarcinoma

ABSTRACT

Quercetin and epigallocatechin-3-gallate (EGCG) are dietary phytochemicals with anti-inflammatory and antitumor effects. In the present study we examined effects of these two compounds on JAK/STAT pathway of cholangiocarcinoma (CCA) cells, because CCA is one of the aggressive cancers with very poor prognosis and JAK/STAT pathway is critically important in inflammation and carcinogenesis. The results showed that the JAK/STAT pathway activation by proinflammatory cytokine interleukin-6 and interferon- γ in CCA cells was suppressed by pretreatment with quercetin and EGCG, evidently by decrease of the elevated phosphorylated-STAT1 and -STAT3 proteins in a dose-dependent manner. The cytokine-mediated up-regulation of inducible nitric oxide synthase (iNOS) and intercellular adhesion molecule-1 (ICAM-1) via JAK/STAT cascade was abolished by both quercetin and EGCG pretreatment. Moreover these flavonoids also could inhibit growth and cytokine-induced migration of CCA cells. Pretreatment with specific JAK inhibitors, AG490 and piceatannol, abolished cytokine-induced iNOS and ICAM-1 expression. These results demonstrate beneficial effects of quercetin and EGCG in suppression of JAK/STAT cascade of CCA cells. Quercetin and EGCG would be potentially useful as cancer chemopreventive agents against CCA.

Key words: JAK/STAT, quercetin, EGCG, cholangiocarcinoma, phytochemical, chemopreventive

INTRODUCTION

Epidemiological studies have shown that the amount of consumption of vegetables and fruits rich in flavonoids inversely associated with the risk of human cancers (Riboli and Norat, 2003). The anti-inflammatory effect of flavonoids is recognized as an important action for their cancer chemoprevention property (Garcia-Lafuente *et al.*, 2009). Quercetin, the most abundant flavonoid in the human diet, and epigallocatechin-3-gallate (EGCG), the biologically most active constituent in green tea, are proposed to have cancer-preventive activities (Hemalswarya and Doble, 2006). The potential chemopreventive effects of these two compounds have been attributed to various mechanisms including anti-oxidative activity as well as capability to modulate cellular signal transduction pathways involving anti-inflammatory, anti-proliferative and induction of apoptosis of tumor cells (Shanmugam *et al.*, 2011). Quercetin and EGCG can interact with non-receptor protein kinases and receptor tyrosine kinases such as epidermal growth factor receptor and vascular endothelial growth factor receptor (Shanmugam *et al.*, 2011). At present, quercetin and EGCG have been used in clinical trials for prevention of several types of cancers such as breast, lung and prostate cancers (<http://www.clinicaltrials.gov/>).

Cholangiocarcinoma (CCA) is a malignant epithelial neoplasm of the biliary tree with very poor prognosis. It is a rare type of cancer worldwide, however populations residing in the Southeast Asian region are at very high risk. The high incidence of CCA in this region is associated with background conditions particularly liver fluke infection (i.e. *Opisthorchis viverrini* and *Chlonorchis sinensis*) that causes long-standing inflammation, cell injury, and reparative biliary epithelial cell proliferation (Sripa and Pairojkul, 2008).

Persistent inflammation can create a local environment enriched with cytokines and other growth factors that primes for cells to develop autonomous proliferative signaling and enhanced production of mitogenic factors (Wise *et al.*, 2008). Excretory/secretory products of liver flukes can stimulate epithelial bile duct cells to secrete proinflammatory cytokine interleukin-6 (IL-6) (Ninlawan *et al.*, 2010). In addition, serum interferon-gamma (IFN- γ) and its expression level in the liver of mice were increased by *Opisthorchis viverrini* infection (Nair *et al.*, 2011). These suggest that various inflammatory mediators, particularly proinflammatory cytokines, are common and important contributors for cholangiocarcinogenesis.

Proinflammatory cytokines activate the signal transducers and activation of transcription (STAT) proteins, which are considered to be oncogenic transcription factors (Hodge *et al.*, 2005). Stimuli to cell surface cytokine receptors activates the Janus-like kinase (JAK) family of protein kinases, which, in turn phosphorylates and activates latent cytoplasmic STAT proteins to an active dimer, leading to nuclear translocation, DNA binding and subsequently modulating gene transcription (Darnell, 1997). Several other kinases including members of the Src and Abl family have also been implicated in the phosphorylation of STATs (Darnell, 1997). Among seven known mammalian STAT proteins, STAT1 and STAT3 are the most widely studied isoforms due to their constitutively activated state in many tumors but not in normal cells (Hodge *et al.*, 2005). STAT1 is relatively specific to IFNs and a pivotal transcription factor in IFN- γ -induced expression of inflammatory genes, whereas STAT3 is mainly activated by IL-6 and other gp130-related cytokines (Yoshimura, 2006).

The crucial roles of STATs, particularly STAT1 and STAT3, in inflammation and tumorigenesis have been demonstrated in several studies such as inflammation-associated gastric tumorigenesis (Ernst *et al.*, 2008) and colitis-associated tumorigenesis (Grivennikov *et al.*, 2009). The activation of JAK/STAT cascade triggered by inflammatory cytokines produced by tumor infiltrating immune cells can lead to the expression of STAT-regulated genes including proinflammatory enzymes and proteins in which then enhance inflammatory condition (Dalwadi *et al.*, 2005). In addition, the activation of JAK/STAT pathway also results in induction of genes that mediate cell proliferation, suppression of apoptosis and promotion of angiogenesis (Barton *et al.*, 2004).

Presently there is neither effective treatment for patients with the advanced stage of CCA nor any effective agents for chemoprevention of CCA. Similar to several cancers, the JAK/STAT signaling pathway is involved in the development of CCA (Isomoto *et al.*, 2007). Thus, STAT signaling pathway inhibitors may be useful for prevention and treatment of CCA. Currently, there is only few studies addressed the role of JAK/STAT signaling pathway in carcinogenesis of CCA.

In the present study, we have investigated effects of quercetin and EGCG on JAK/STAT pathway using CCA cells treated with proinflammatory cytokine mixture, IL-6 and IFN- γ , for mimicking inflammatory condition (Kaur *et al.*, 2003). The results show that both quercetin and EGCG could inhibit JAK/STAT signaling cascade. They also could suppress cytokine-induced the expression of inducible nitric oxide synthase (iNOS) and intercellular adhesion molecule-1 (ICAM-1), the key molecules involving in inflammatory

and tumorigenesis processes. Moreover, they inhibited growth and migration of CCA cells. Our findings provide new insights into the intervention strategy for prevention of CCA.

MATERIALS AND METHODS

Reagents

Quercetin, EGCG and piceatannol were purchased from Sigma Chemical (St.Louis, MO, USA). AG490 was obtained from Calbiochem (San Diego, CA, USA). Interleukin-6 (IL-6), interferon- γ (IFN- γ) and the primary antibodies against STAT1, STAT3, phospho-STAT1 (Tyr⁷⁰¹) and phospho-STAT3 (Tyr⁷⁰⁵) were purchased from Cell Signaling Technology (Danvers, MA, USA). RIPA buffer was from Amresco (Solon, OH, USA). The pGL4 basic and pGL4[*hRluc*] vector, LipofectamineTM 2000 and Dual-Luciferase[®] Reporter Assay Kit were obtained from Promega (Madison, WI, USA). Reagents for cell culture were from Gibco BRL Life Technologies (Grand Island, NY, USA).

Cell lines and cell cultures

The human CCA cell lines; KKU100, KKU-M139 and KKU-M213 used in this study were kindly provided by Dr. Banchob Sripa of Department of Pathology, Faculty of Medicine, Khon Kaen University. These cell lines were cultured in complete media consisting of Ham's F12 media, supplemented with 10% fetal calf serum, 12.5 mM HEPES, pH 7.3, 100 U/mL penicillin G and 100 μ g/mL streptomycin and maintained under an atmosphere of 5% CO₂ at 37°C. The cells were subcultured every 3 days using 0.25% trypsin-EDTA.

Cells treatment

At about 70% confluency, cultures were starved in serum-free medium for 16 h and then treated with cytokine mixture, combination of IL-6 and IFN- γ (Kaur *et al.*, 2003), in serum-free medium for further 1 h to 24 h. The concentration used for both IL-6 and IFN- γ were 100 ng/mL. In experiments involving flavonoids or JAK inhibitors, the cells were pretreated with various concentrations of quercetin (1, 10, 50 and 100 μ M) or EGCG (1, 10, 25 and 50 μ M) or piceatannol (JAK1 inhibitor) (Barton *et al.*, 2004) (1, 10 and 100 μ M) or AG490 (JAK2 inhibitor) (Barton *et al.*, 2004) (1, 10 and 100 μ M) in DMSO for 2 h and then further treated with cytokine mixture. An equal amount of DMSO (vehicle) was present in each treatment, including control. DMSO concentration did not exceed 0.1% (v/v) in any treatment.

Protein extraction and Western blot analysis

After treatments for designated period of times, medium was aspirated, cells were washed twice with cold PBS and whole cell lysates were prepared using RIPA cell lysis buffer (Amresco, OH, USA) according to the manufacturer's instructions. The proteins were resolved by SDS-PAGE using 10% polyacrylamide gel and the separated proteins on the gel were electrophoretically transferred to a PVDF membrane. The membranes were blocked with 5% (w/v) bovine serum albumin in Tris buffered saline containing 0.1% Tween-20 for 2 h at room temperature, followed by incubation with primary antibody against STAT1 (1:1000), phospho-STAT1 (Tyr⁷⁰¹) (1:1000), STAT3 (1:1000), phospho-STAT3 (Tyr⁷⁰⁵) (1:1000) and β -actin (1:5000) at 4°C overnight. After washing, the

membranes were incubated with the appropriate horseradish peroxidase conjugated secondary antibody to detect bands by AmershamTM ECLTM Prime (Amersham Biosciences Corp, NJ, USA). The densities of the specific protein bands were visualized and captured by ImageQuantTM 400.

Real-time polymerase chain reaction

Total RNA was extracted from cells using Trizol reagent (Invitrogen, USA) according to the manufacturer's instructions. One µg of total RNA was then reverse transcribed to single-stranded cDNA by the ImProm-IITM reverse transcription system (Promega, WI, USA). PCR was carried out using specific primers as follows: iNOS forward 5'-GTTCTCACGGCACAGGTCTC-3' and iNOS reverse 5'-GCAGGTCACCTTATGTCAC TTATC-3' (Wu *et al.*, 2011), ICAM-1 forward 5'-CAAGGCCTCAGTCAGTGTGA-3' and ICAM-1 reverse 5'-CCTCTGGCTTCGTCAGAATC-3' (Kim *et al.*, 2010), GAPDH forward 5'-GTGGTGGACCTGACCTGC-3' and GAPDH reverse 5'-TGAGCTTGACA AAGTGGTCG-3' (Kim *et al.*, 2010). The relative expression of iNOS and ICAM-1 was analyzed using quantitative RT-PCR with GAPDH as an internal control. The PCR was performed with SsoFastTM EvaGreen Supermix with low Rox (Bio-Rad, CA, USA) using an ABI Prism 7500 Sequence Detection System (Applied Biosystems, Foster City, CA, USA) under the following conditions: denaturation at 95°C for 3 min and amplification by cycling 40 times at 95°C for 15 sec and 60°C for 31 sec.

Cell growth assay

Sulforhodamine B (SRB) was used to measure the effect of quercetin and EGCG on the proliferation of CCA cell lines as described previously (Prawan *et al.*, 2009). Briefly, K KU100 cells were plated in a 96-well plate for 24 h. After exposure of cultured cells to quercetin or EGCG at various concentrations for 48 h, the culture medium was removed, 10% cold trichloroacetic acid was added for 1 h at 4 °C, and subsequently washed 5 times with deionized H₂O. The plate was air dried, and 0.4% SRB in 1% acetic acid was added for 30 min. Unbound dye was washed out 5 times with 1% acetic acid. After air drying, SRB dye within cells was solubilized with 200 µl of 10 mM unbuffered Tris base solution. The plate was shaken for 10 min, and the absorbance was measured at 540 nm by using a microplate reader. Cell growth inhibition was expressed in terms of percentage of untreated control absorbance following subtraction of mean background absorbance. The IC₅₀ concentration (50% inhibition of cell growth values) was calculated from the dose-response curves.

Wound healing assay

K KU100 cells (1.5 x 10⁵) were seeded into a 24-well plate and incubated overnight in Ham's F12 medium, supplemented with 10% fetal calf serum. A scratch wound was made with a sterile 200 µL pipette tip. After washing with PBS to remove any detached cell, a series of images of the scratched wound were taken from 0 to 24 h. The cells were pretreated with quercetin (10 µM) or EGCG (5 µM) or vehicle for 2 h, then with cytokine mixture afterward. The closing of scratched wound, represented the migration process, was

determined by capture of the denuded area along the scratch using Image-Pro Plus software (Media Cybernetics, LP, USA). The wound distance was calculated by dividing the area by the length of scratch.

STAT3 luciferase reporter assay

To construct the STAT3 luciferase vector, the STAT3 transcriptional regulatory element 5'-GTCGACATTTCCCGTAAATCGTCGA-3' (Bergad *et al.*, 2000) containing NheI/HindIII restriction sites was subcloned into the multiple cloning site of the pGL4 luciferase reporter plasmid (Promega, WI, USA). The sequences of the constructs were verified by the MegaBACE DNA Analysis System (Amersham Biosciences Corp, NJ, USA). For reporter assay, the cells (1×10^4) were seeded into each well of 96-well plates, and cultured for 24 h. The cells were transfected with 0.2 μ g of STAT3-pGL4 reporter vector according to the manufacturer's protocols using X-tremeGene HP (Roche) for KKKU100 cells or LipofectamineTM 2000 (Promega, WI, USA) for KKKU-M139 and KKKU-M213 cells. The cells were also cotransfected with 0.01 μ g of pGL4[*hRluc*/TK] vector containing the *Renilla luciferase* gene (Promega) to control transfection efficiency. After 6 h of transfection, cells were pretreated with flavonoids for 2 h and then stimulated with the cytokine mixture for additional 18 h before being washed and lysed in luciferase lysis buffer (Promega). Luciferase activity was measured with a luminometer by using a Dual-Luciferase[®] Reporter Assay Kit (Promega).

RESULTS

Cytokine mixture induced STAT1 and STAT3 activation in KKU100 cells

To clarify whether JAK/STAT signaling is intact in CCA cells, the effect of cytokine mixture (100 ng/mL IL-6 and 100 ng/mL IFN- γ) on STAT levels in KKU100 cells was determined at various time points (1 h to 24 h). In separate experiments, we confirmed that the dose range of cytokine mixture used in this study did not affect the cell viability (data not shown). After treatment with the cytokine mixture, phosphorylation of STAT1 (Tyr⁷⁰¹) in KKU100 cells was dramatically increased as early as within the first hour, and then gradually decreased afterwards, although it was still detectable at 24 h (Fig. 1A). For STAT3, the cytokine mixture induced transient STAT3 (Tyr⁷⁰⁵) phosphorylation in KKU100 cells within the first hour, but the phospho-STAT3 was diminished rapidly within 2 h (Fig. 1A). From these results, functional JAK/STAT signaling cascade is present in CCA cells.

Quercetin and EGCG inhibited cytokine-induced STAT1 and STAT3 phosphorylation

To determine whether quercetin and EGCG can suppress the proinflammatory cytokine-induced JAK/STAT phosphorylation, KKU100 cells were pre-incubated with various concentrations of quercetin or EGCG for 2 h and then stimulated with the cytokine mixture for 1 h. Under these conditions, the cell viability was not affected (data not shown). As shown in Fig. 1, when KKU100 cells were pretreated with quercetin (Fig. 1B) or EGCG (Fig. 1C), cytokine-induced STAT1 and STAT3 phosphorylation was markedly

suppressed in a dose-dependent manner, although STAT1 and STAT3 protein levels were not affected.

Quercetin and EGCG suppressed cytokine mixture-induced iNOS and ICAM-1 expression

It is known that iNOS and ICAM-1 play a critical role in inflammatory and tumorigenesis processes (Yamada *et al.*, 2006; Kostourou *et al.*, 2011). Therefore, we first examined the effect of cytokine treatment on the expression of iNOS and ICAM-1 in CCA cells using real time-PCR. The results showed that cytokine treatment markedly up-regulated the mRNA expression of iNOS and ICAM-1 (Fig. 2). Pretreatment with quercetin and EGCG significantly inhibited cytokine-mediated up-regulation of iNOS and ICAM-1 expression in a dose-dependent manner (Fig. 2A and 2B, respectively). The inhibitory effect of EGCG was more potent than that of quercetin.

JAK1 and JAK2 inhibition attenuated cytokine-induced expression of iNOS and ICAM-1

To validate whether JAK/STAT signaling contributes to the up-regulation of iNOS and ICAM-1, effects of piceatannol (JAK1 inhibitor) and AG490 (JAK2 inhibitor) on the cytokine-induced STAT1 and STAT3 phosphorylation in KKU100 cells were examined. As illustrated in Fig. 3A, pretreatment with these inhibitors could abrogate STAT1 and STAT3 phosphorylation in a dose-dependent manner. Subsequently, KKU100 cells were pre-incubated with various concentration of piceatannol or AG490 for 2 h and then treated

with the cytokine mixture for 3 h. After that the mRNA expression of iNOS and ICAM-1 were evaluated. The results showed that cytokine-induced iNOS expression in K KU100 cells were suppressed dose-dependently by piceatannol and AG490 pretreatment (Fig. 3B). Similarly, piceatannol and AG490 could suppress the up-regulation of ICAM-1 expression, although the higher concentration was required for the inhibition compared to the effects on iNOS expression (Fig. 3C). Under these conditions, the cell viability was only slightly reduced (data not shown).

Quercetin and EGCG inhibited growth and migration of CCA cell lines

The anticancer effect of quercetin and EGCG was further evaluated by examining some downstream effects of JAK/STAT signaling i.e. anti-proliferation and anti-migration. The anti-proliferative activity was assayed on K KU100 cells using SRB assay. Quercetin and EGCG had similar potency to inhibit K KU100 cell growth with the IC₅₀ values of 31.1 ± 7.1 and 24.3 ± 3.1 μM, respectively (Fig. 4A).

Since quercetin and EGCG showed strong inhibitory effect on cytokine-induced expression of ICAM-1, an adhesion molecule involved in cell migration and invasion, the effect of these two compounds on CCA cell migration was investigated by wound healing assay using the monolayer culture of K KU100 cells. The results are shown in Fig. 4B and 4C. Closure of the scratched wound was stimulated by cytokine mixture, determined at 24 h. The cytokine-mediated stimulation of cell migration was suppressed by quercetin and EGCG, as the migration distance of cultured cells in those flavonoid treated groups was shorter than that of cytokine treatment only.

Quercetin and EGCG inhibited STAT3-dependent luciferase activity in CCA cells

The suppressive effects of quercetin and EGCG on JAK/STAT signaling were confirmed further by STAT3-luciferase reporter assay. Firstly, we demonstrated that quercetin and EGCG suppressed cytokine-induced STAT3 luciferase activity in KKU100 cells (Fig. 5A). To explore whether the suppressive effects of flavonoids on STAT3 signaling is common in various CCA cells, in addition to KKU100 cells, KKU-M139 and KKU-M213 cells were employed in this experiment. These three human CCA cell lines were established from primary tumors of liver fluke-associated CCA patients with different histological types i.e. poorly differentiated, adeno-squamous, and mixed papillary and non-papillary adenocarcinoma, respectively. As was expected, cytokine-induced STAT3 activity in KKU-M139 and KKU-M213 cells were strongly suppressed by quercetin and EGCG as shown in Fig. 5B and 5C, respectively, suggesting the effect of the flavonoids are not restricted to a special cell type.

DISCUSSION

Chemoprevention and chemotherapy of CCA are so far quite disappointing. Several efforts have been made to identify and develop new agents for both treatment and prevention of this deadly disease. These include targeted therapy on some critical molecules of cancer cells, such as NQO1 and HO-1, where inhibition of the enzymes make CCA cells sensitive to chemotherapeutic agents (Buranrat *et al.*, 2010; Kongpetch *et al.*, 2012). In the present study, we have demonstrated that JAK/STAT signaling pathway may be an important target for inhibition of CCA. JAK/STAT signaling plays an important role

in the expression of iNOS and ICAM-1, both of which are known critical molecules involving in inflammation and carcinogenesis processes. Quercetin and EGCG inhibited JAK/STAT signaling cascade in CCA cells leading to the down-regulation of iNOS and ICAM-1 and suppression of cell migration. In addition, these two phytochemical compounds could also inhibit the growth of CCA cells. The results provide evidence supporting the use of these two flavonoids as chemopreventive agents for CCA.

Quercetin and EGCG are potent inhibitors of tumor growth and inflammation and have traditionally been used to treat many inflammatory disorders (Garcia-Lafuente *et al.*, 2009). Quercetin and EGCG have been shown to exert their effects through inhibiting several signaling pathway such as NF- κ B (Priyadarsini and Nagini, 2012), AP-1 (Shanmugam *et al.*, 2011) and JAK/STAT signaling pathway (Tedeschi *et al.*, 2002; Michaud-Levesque *et al.*, 2012). We also found that quercetin and EGCG are potent inhibitors of cytokine-induced STAT1 as well as STAT3 activation in CCA cells.

In the present study, we demonstrated that the activation of JAK/STAT pathway is associated with the increase in expression of iNOS and ICAM-1. The iNOS has critical functions in inflammation-related diseases. Sustained and excess nitric oxide generation, most of which is attributable to iNOS expression, often is pathogenic to cause inflammation, angiogenesis and neoplasia (Kostourou *et al.*, 2011). Quercetin as well as EGCG are known to exhibit suppression of iNOS expression in various cancer cells (Gerhauser *et al.*, 2003). It has been reported that iNOS gene promoter has binding sites for STAT, AP-1, NF- κ B and HIF-1 α (Chittezhath *et al.*, 2008). In this study, we demonstrated that cytokine-induced STAT activation is an important determinant of iNOS

expression in CCA cells, as JAK inhibitors markedly inhibited the expression. The inhibition of STAT activation by quercetin correlated with the decreases in iNOS mRNA levels. Similar inhibitory patterns were observed after EGCG treatment, and the inhibitory effect of EGCG was stronger than quercetin. These results indicate that the inhibitory effects of quercetin and EGCG on cytokine-induced iNOS expression are mediated partly via suppression of JAK/STAT pathway. Because no other pathways were tested in the present study, further studies are required to elucidate the action of quercetin and EGCG on the other signaling pathways involving in iNOS expression.

ICAM-1, of which expression is regulated by JAK/STAT pathway, plays a critical role in inflammation-related diseases by promoting trafficking of leukocytes across endothelia and epithelial barriers. It also implicated in carcinogenic processes by contributing in tumor metastasis (Yamada *et al.*, 2006). The modulation of ICAM-1 expression is, therefore, an important target for chemoprevention and chemotherapy, as shown by the beneficial effect of ICAM-1 inhibitors on the tumorigenesis and cancer progression (Lee *et al.*, 2012). Similar to iNOS, transcription factors important to the activation of ICAM-1 expression include AP-1, NF- κ B and STAT (van de Stolpe and van der Saag, 1996).

In this study, we found that although AG490 at 100 μ M profoundly suppressed the phosphorylation of STAT1 and STAT3, this treatment caused a significant but small suppression on ICAM-1 expression. These results suggest that ICAM-1 expression may be modulated by multiple signaling pathways. The suppressive effect of piceatannol on ICAM-1 expression was also incomplete. It should be noted that piceatannol may inhibit

NF- κ B which is also involved in the regulation of ICAM-1 expression (Ashikawa *et al.*, 2002). In this study we found that quercetin and EGCG both dramatically suppress cytokine-induced ICAM-1 expression, where EGCG have higher inhibitory effect than quercetin. Quercetin and EGCG exert their suppressive effects on cytokine-induced ICAM-1 up-regulation, at least in part, through modulating the JAK/STAT cascade in CCA cells.

Enhanced cell migration and invasion are important aspects of cancer phenotypes, where ICAM-1 plays an important role (Yamada *et al.*, 2006). In this study, we found that cytokine-induced migration of CCA cells was associated with ICAM-1 expression. Quercetin and EGCG could significantly abolish the cytokine-dependent CCA cell migration. It should be noted that the anti-migratory effect of flavonoids was detected at low concentrations which minimally inhibited cell growth. In fact, the modulation of cell migration ability is a target for chemopreventive compound. Thus, our findings provide a supportive evidence that quercetin and EGCG are potentially useful as cancer chemopreventive agents for CCA.

To ascertain that the inhibition of JAK/STAT signaling by quercetin and EGCG is not confined to only one cell type, we performed the reporter assay with other two CCA cell lines. It is clearly demonstrated that both flavonoids inhibited cytokine-induced STAT3 luciferase activity in all three CCA cell lines examined. These results imply that quercetin and EGCG may efficiently inhibit JAK/STAT signaling pathway in most CCA cells and this confers a high potential therapeutic opportunity.

1
2
3
4
5 In summary, the present study showed that JAK/STAT signaling cascade is
6
7 functioning in CCA cells and upon activation it induces the expression of iNOS and
8
9 ICAM-1. Quercetin and EGCG can suppress STAT1- and STAT3-phosphorylation and
10
11 also suppress iNOS and ICAM-1 expression. Moreover, these two compounds can inhibit
12
13 cytokine-induced CCA cell migration as well. Given that JAK/STAT cascade plays a
14
15 critical role in the inflammation and carcinogenesis processes, inhibition of JAK/STAT
16
17 signaling pathway could be a valuable chemopreventive target of natural compounds to
18
19 protect against inflammation-associated cancer.
20
21
22
23
24
25

26 **Acknowledgements**

27
28 This work was supported by the Thailand Research Fund (Grant No.
29
30 MRG5480011), Office of the Higher Education Commission, Khon Kaen University, and
31
32 the Higher Education Research Promotion and National Research University Project of
33
34 Thailand, Office of the Higher Education Commission, through the Health cluster (SHeP-
35
36 GMS) of Khon Kaen University. The authors would like to thank Prof. Dr. Yukifumi
37
38 Nawa, Invited Professor/Consultant, Khon Kaen University, for editing the manuscript.
39
40
41
42
43
44

45 **Conflict of Interest**

46
47 The authors have declared that there is no conflict of interest.
48
49
50
51
52
53
54
55
56
57
58
59
60

REFERENCES

Ashikawa K, Majumdar S, Banerjee S, Bharti AC, Shishodia S, Aggarwal BB. 2002. Piceatannol inhibits TNF-induced NF-kappaB activation and NF-kappaB-mediated gene expression through suppression of IkappaBalpha kinase and p65 phosphorylation. *J Immunol* **169**(11): 6490-6497.

Barton BE, Karras JG, Murphy TF, Barton A, Huang HF. 2004. Signal transducer and activator of transcription 3 (STAT3) activation in prostate cancer: Direct STAT3 inhibition induces apoptosis in prostate cancer lines. *Mol Cancer Ther* **3**(1):11-20.

Bergad PL, Schwarzenberg SJ, Humbert JT, Morrison M, Amarasinghe S, Towle HC, *et al.* 2000. Inhibition of growth hormone action in models of inflammation. *Am J Physiol Cell Physiol* **279**(6): C1906-1917.

Buranrat B, Prawan A, Kukongviriyapan U, Kongpetch S, Kukongviriyapan V. 2010. Dicoumarol enhances gemcitabine-induced cytotoxicity in high NQO1-expressing cholangiocarcinoma cells. *World J Gastroenterol* **16**(19): 2362-2370.

Chittezhath M, Deep G, Singh RP, Agarwal C, Agarwal R. 2008. Silibinin inhibits cytokine-induced signaling cascades and down-regulates inducible nitric oxide synthase in human lung carcinoma A549 cells. *Mol Cancer Ther* **7**(7): 1817-1826.

Dalwadi H, Krysan K, Heuze-Vourc'h N, Dohadwala M, Elashoff D, Sharma S, *et al.* 2005. Cyclooxygenase-2-dependent activation of signal transducer and activator of transcription 3 by interleukin-6 in non-small cell lung cancer. *Clin Cancer Res* **11**(21): 7674-7682.

Darnell JE, Jr. 1997. STATs and gene regulation. *Science* **277**(5332): 1630-1635.

- Ernst M, Najdovska M, Grail D, Lundgren-May T, Buchert M, Tye H, *et al.* 2008. STAT3 and STAT1 mediate IL-11-dependent and inflammation-associated gastric tumorigenesis in gp130 receptor mutant mice. *J Clin Invest* **118**(5): 1727-1738.
- Garcia-Lafuente A, Guillamon E, Villares A, Rostagno MA, Martinez JA. 2009. Flavonoids as anti-inflammatory agents: implications in cancer and cardiovascular disease. *Inflamm Res* **58**(9): 537-552.
- Gerhauser C, Klimo K, Heiss E, Neumann I, Gamal-Eldeen A, Knauf J, *et al.* 2003. Mechanism-based in vitro screening of potential cancer chemopreventive agents. *Mutat Res* **523-524**: 163-172.
- Grivennikov S, Karin E, Terzic J, Mucida D, Yu GY, Vallabhapurapu S, *et al.* 2009. IL-6 and Stat3 are required for survival of intestinal epithelial cells and development of colitis-associated cancer. *Cancer Cell* **15**(2): 103-113.
- HemaIswarya S, Doble M. 2006. Potential synergism of natural products in the treatment of cancer. *Phytother Res* **20**(4): 239-249.
- Hodge DR, Hurt EM, Farrar WL. 2005. The role of IL-6 and STAT3 in inflammation and cancer. *Eur J Cancer* **41**(16): 2502-2512.
- Isomoto H, Mott JL, Kobayashi S, Werneburg NW, Bronk SF, Haan S, *et al.* 2007. Sustained IL-6/STAT-3 signaling in cholangiocarcinoma cells due to SOCS-3 epigenetic silencing. *Gastroenterology* **132**(1): 384-396.
- Kaur N, Kim IJ, Higgins D, Halvorsen SW. 2003. Induction of an interferon-gamma Stat3 response in nerve cells by pre-treatment with gp130 cytokines. *J Neurochem* **87**(2): 437-447.

- Kim YS, Im J, Choi JN, Kang SS, Lee YJ, Lee CH, *et al.* 2010. Induction of ICAM-1 by *Armillariella mellea* is mediated through generation of reactive oxygen species and JNK activation. *J Ethnopharmacol* **128**(1): 198-205.
- Kongpetch S, Kukongviriyapan V, Prawan A, Senggunprai L, Kukongviriyapan U, Buranrat B. 2012. Crucial role of heme oxygenase-1 on the sensitivity of cholangiocarcinoma cells to chemotherapeutic agents. *PLoS One* **7**(4): e34994.
- Kostourou V, Cartwright JE, Johnstone AP, Boulton JK, Cullis ER, Whitley G, *et al.* 2011. The role of tumour-derived iNOS in tumour progression and angiogenesis. *Br J Cancer* **104**(1): 83-90.
- Lee IT, Lin CC, Lee CY, Hsieh PW, Yang CM. 2012. Protective effects of (-)-epigallocatechin-3-gallate against TNF- α -induced lung inflammation via ROS-dependent ICAM-1 inhibition. *J Nutr Biochem* **68**(3): 946-955.
- Michaud-Levesque J, Bousquet-Gagnon N, Beliveau R. 2012. Quercetin abrogates IL-6/STAT3 signaling and inhibits glioblastoma cell line growth and migration. *Exp Cell Res* **318**(8): 925-935.
- Nair SS, Bommana A, Pakala SB, Ohshiro K, Lyon AJ, Suttiprapa S, *et al.* 2011. Inflammatory response to liver fluke *Opisthorchis viverrini* in mice depends on host master coregulator MTA1, a marker for parasite-induced cholangiocarcinoma in humans. *Hepatology* **54**(4): 1388-1397.
- Ninlawan K, O'Hara SP, Splinter PL, Yongvanit P, Kaewkes S, Surapaitoon A, *et al.* 2010. *Opisthorchis viverrini* excretory/secretory products induce toll-like receptor 4

- upregulation and production of interleukin 6 and 8 in cholangiocyte. *Parasitol Int* **59**(4): 616-621.
- Prawan A, Buranrat B, Kukongviriyapan U, Sripa B, Kukongviriyapan V. 2009. Inflammatory cytokines suppress NAD(P)H:quinone oxidoreductase-1 and induce oxidative stress in cholangiocarcinoma cells. *J Cancer Res Clin Oncol* **135**(4): 515-522.
- Priyadarsini RV, Nagini S. 2012. Quercetin suppresses cytochrome P450 mediated ROS generation and NFkappaB activation to inhibit the development of 7,12-dimethylbenz[a]anthracene (DMBA) induced hamster buccal pouch carcinomas. *Free Radic Res* **46**(1): 41-49.
- Riboli E, Norat T. 2003. Epidemiologic evidence of the protective effect of fruit and vegetables on cancer risk. *Am J Clin Nutr* **78**(3 Suppl): 559S-569S.
- Shanmugam MK, Kannaiyan R, Sethi G. 2011. Targeting cell signaling and apoptotic pathways by dietary agents: role in the prevention and treatment of cancer. *Nutr Cancer* **63**(2): 161-173.
- Sripa B, Pairojkul C. 2008. Cholangiocarcinoma: lessons from Thailand. *Curr Opin Gastroenterol* **24**(3): 349-356.
- Tedeschi E, Suzuki H, Menegazzi M. 2002. Antiinflammatory action of EGCG, the main component of green tea, through STAT-1 inhibition. *Ann N Y Acad Sci* **973**: 435-437.
- van de Stolpe A, van der Saag PT. 1996. Intercellular adhesion molecule-1. *J Mol Med (Berl)* **74**(1): 13-33.

Wise C, Pilanthananond M, Perry BF, Alpini G, McNeal M, Glaser SS. 2008. Mechanisms of biliary carcinogenesis and growth. *World J Gastroenterol* **14**(19): 2986-2999.

Wu SQ, Otero M, Unger FM, Goldring MB, Phrutivorapongkul A, Chiari C, *et al.* 2011. Anti-inflammatory activity of an ethanolic *Caesalpinia sappan* extract in human chondrocytes and macrophages. *J Ethnopharmacol* **138**(2): 364-372.

Yamada M, Yanaba K, Hasegawa M, Matsushita Y, Horikawa M, Komura K, *et al.* 2006. Regulation of local and metastatic host-mediated anti-tumour mechanisms by L-selectin and intercellular adhesion molecule-1. *Clin Exp Immunol* **143**(2): 216-227.

Yoshimura A. 2006. Signal transduction of inflammatory cytokines and tumor development. *Cancer Sci* **97**(6): 439-447.

FIGURE LEGENDS

Figure 1: Effects of cytokine mixture on the activation of STAT and effects of quercetin and EGCG on cytokine-induced STAT activation. (A) K KU100 cells were incubated with cytokine combination (100 ng/mL IL-6 and 100 ng/mL IFN- γ) for various time points (1 to 24 h). (B-C) K KU100 cells were pretreated with the indicated concentrations of (B) quercetin or (C) EGCG for 2 h and then treated with cytokine mixture for 1 h. Whole cell extracts were prepared and 20 μ g aliquots of those extracts were resolved on 10% SDS-PAGE, electrotransferred onto PVDF membranes and probed using the indicated antibodies as described under Materials and Methods. Anti- β -actin antibody was used as a loading control. Results shown are representative of at least two reproducible independent experiments.

Figure 2: Effects of quercetin and EGCG on cytokine-induced expression of iNOS and ICAM-1. KKKU100 cells were pretreated with various concentrations of quercetin (1 - 100 μ M) or EGCG (1 - 50 μ M) for 2 h and then treated with cytokine combination (100 ng/mL IL-6 and 100 ng/mL IFN- γ) for 3 h. At the end of treatments, total RNA of cells were collected and real time RT-PCR was carried out for (A) iNOS and (B) ICAM-1 expression. The bars represent relative expression of indicated genes normalized with GAPDH. The data are expressed as mean $\pm$ SD, each from three independent experiments. *, $p < 0.05$ vs cytokine treatment alone.

Figure 3: Effects of JAK inhibitors on cytokine-mediated phosphorylation of STAT and -induced iNOS and ICAM-1 expression. (A) KKKU100 cells were pretreated with the indicated concentrations of AG490 or piceatannol for 2 h and then treated with cytokine mixture for 1 h. Whole cell extracts were prepared and 20 μ g aliquots of those extracts were resolved on 10% SDS-PAGE, electrotransferred onto PVDF membranes and probed using the indicated antibodies as described under Materials and Methods. Anti- β -actin antibody was used as a loading control. Results shown are representative of at least two reproducible independent experiments. (B-C) KKKU100 cells were pretreated with indicated concentrations (1 - 100 μ M) of AG490 or piceatannol for 2 h and then treated with cytokine combination (100 ng/mL IL-6 and 100 ng/mL IFN- γ) for 3 h. At the end of treatments, total RNA of cells were collected and analyzed for (B) iNOS and (C) ICAM-1 expression by real time RT-PCR. The bars represent relative expression of indicated genes normalized with GAPDH. The data are expressed as mean $\pm$ SD, each from three independent experiments. *, $p < 0.05$ vs cytokine treatment alone.

Figure 4: Effects of quercetin and EGCG on the growth and cytokine-induced migration of CCA cells. (A) K KU100 cells were treated with indicated concentration of quercetin or EGCG for 48 h and cell numbers were determined by SRB assay. The results are presented as percentage of control. The data are the mean $\pm$ SD averaged from three independent experiments. (B) Scratched wounds of monolayer K KU100 cells were pretreated with vehicle or 10 μ M of quercetin or 5 μ M of EGCG for 2 h and then treated with cytokine mixture. Cell migration was monitored under phase-contrast microscopy (x4 magnification). Representative images of wound healing were obtained at the time of the scratch and 24 h later. (C) The graph shows the level of cell migration into the wound scratch quantified as the percentage of wound closure at 24 h. The data are mean $\pm$ SD averaged from quadruplets of one experiment. CK: cytokine, QC: quercetin.

Figure 5: Effects of quercetin and EGCG on cytokine-induced transcriptional activity of STAT3. The CCA cells: (A) K KU100 (B) K KU-M139 and (C) K KU-M213 were transfected with STAT3-pGL4 reporter vector together with renilla luciferase plasmid. After 6 h of transfection period, cells were pretreated with indicated concentrations of quercetin or EGCG for 2 h before exposure to cytokine combination (100 ng/mL IL-6 and 100 ng/mL IFN- γ) for additional 18 h. After that the luciferase activity was measured and presented as fold induction to the control cells without cytokine treatment. Each bar indicates the mean $\pm$ SD of five to six determinations. *, $p < 0.05$ vs cytokine treatment alone.




Fig.1: Effects of cytokine mixture on the activation of STAT and effects of quercetin and EGCG on cytokine-induced STAT activation
190x275mm (300 x 300 DPI)




Fig.2: Effects of quercetin and EGCG on cytokine-induced expression of iNOS and ICAM-1
190x275mm (300 x 300 DPI)




Fig.3: Effects of JAK inhibitors on cytokine-mediated phosphorylation of STAT and -induced iNOS and ICAM-1 expression
190x275mm (300 x 300 DPI)




Fig.4: Effects of quercetin and EGCG on the growth and cytokine-induced migration of CCA cells
127x95mm (300 x 300 DPI)

FIG. 5




Fig.5: Effects of quercetin and EGCG on cytokine-induced transcriptional activity of STAT3
476x900mm (96 x 96 DPI)



International Society for
Ethnopharmacology

BOOK OF ABSTRACTS

13th Congress of the International Society for Ethnopharmacology

in collaboration with the

**Society for Medicinal Plant and
Natural Product Research**

and

Eurasia-Pacific Uninet

**Graz, Austria
September 2 - 6, 2012**



Das Land
Steiermark

<http://ise13.uni-graz.at>

Hypoglycemic and hypolipidemic effect of morin (2',3,4',5,7-pentahydroxyflavone) in streptozotocin-induced diabetic rats

Muhammed A¹, Gül N¹

¹ Department of Biology, Faculty of Science, Ankara University, 06100 Tandoğan, Ankara, Turkey.

Diabetes mellitus (DM) is one of the major public health burden worldwide. Recently, the treatment of diabetes with natural products (flavonoids specially) witnessed a growing interest because side effects of insulin and oral antidiabetic drugs. The present work is carried out to study the beneficial effects of a flavonoid extracted from plants which called morin (2',3,4',5,7-pentahydroxyflavone) on blood glucose level and lipid profile level in normal and streptozotocin induced diabetic rats (STZ 45 mg/kg, i.p). Animals were divided into seven groups (n: 6 in each group): Group 1: Normal control, group 2: Normal + morin 25 mg/kg, group 3: Normal + morin 50 mg/kg, group 4: Diabetic control, group 5: Diabetic + morin 25 mg/kg, group 6: Diabetic + morin 50 mg/kg and group 7: Diabetic + Tolbutamide 100 mg/kg. Morin could not prevent loss of body weight, but significantly decrease water and food (25 mg/kg) intake. It showed significant prevention in elevation of glucose and prevent increase in the cholesterol, triglyceride, LDL-cholesterol, VLDL-cholesterol levels in serum of diabetic rats. Morin also significantly prevented decrease in the HDL-cholesterol level and increase in AST and ALT levels in diabetic rats compared to that of diabetic control. These results indicated that morin could prevent abnormal changes in blood glucose and lipid profile in streptozotocin induced diabetic rats.

Chemopreventive effects of quercetin and EGCG acting by suppression on JAK/STAT3 signaling pathway

Senggunprai L, Prawan A, Kukongviriyapan V, Zeekpudsa P, Boonlor K

Department of Pharmacology, Faculty of Medicine, and Liver Fluke and Cholangiocarcinoma Research Center, Khon Kaen University, Khon Kaen 40002, Thailand

Quercetin and epigallocatechin-3-gallate (EGCG) are dietary phytochemicals with antiinflammatory and antitumor effects [1]. The mechanisms underlying their effects may involve induction of antioxidant enzymes and suppression of pro-inflammatory cytokine activity. As JAK/STAT3 pathway is critical in inflammation and carcinogenesis and constitutively active in most tumor but not in normal cells [2], in the present study we examined the chemopreventive effect of quercetin and EGCG on this pathway using cholangiocarcinoma (CCA) cells treated with pro-inflammatory cytokine mixture (IL-6 + IFN- γ) for mimicking inflammatory condition. The results showed that cytokine mixture treatment could activate the STAT3 signaling cascade in CCA cells demonstrated by increased levels of phosphorylated-STAT3 protein (p-STAT3) in both whole cell and nuclear lysates. Pretreatment of cells with quercetin and EGCG dose-dependently suppressed the STAT3 activation. We further examined some down-stream genes of JAK/STAT3 signaling pathways. The expression of inducible nitric oxide synthase (iNOS) was strongly induced by treatment with the cytokine combination in relation to the activation of STAT3. Both quercetin and EGCG were able to inhibit the cytokine-mediated up-regulation of iNOS expression. Detailed molecular analyses revealed that pretreatment of cells with specific JAK inhibitor, AG490, diminished cytokine-induced iNOS expression in a dose-dependent manner, suggesting a role of JAK in cytokine mediated iNOS expression via STAT3 activation in CCA cells. Taken together, the results suggested that quercetin and EGCG could downregulate iNOS expression through inhibiting JAK/STAT3 signaling cascade. Inhibition of STAT3 pathway could be valuable chemopreventive targets of natural chemopreventive compounds to protect against inflammatory-associated cancer.

Acknowledgements: This work was supported by the Thailand Research Fund, Office of the Higher Education Commission, the National Research University Project and Office of the Higher Education Commission through SHEP-GMS of Khon Kaen University, Thailand.

References: 1. Garcia-Lafuente, A. et al. (2009). *Inflamm Res* 58(9): 537-52. 2. Hodge, R. et al. (2005). *Eur J Cancer* 41(16): 2502-12.

Chemopreventive effects of quercetin and EGCG acting by suppression on JAK/STAT3 signaling pathway

Laddawan Senggunprai, Veerapol Kukongviriyapan, Auemduan Prawan, Ponsilp Zeekpudsa, Kampeeporn Boonlor

Department of Pharmacology, Faculty of Medicine, and Liver Fluke and Cholangiocarcinoma Research Center, Khon Kaen University, Khon Kaen, Thailand

Quercetin, the most abundant flavonoid in the human diet, and the biologically most active constituent of green tea, epigallocatechin-3-gallate (EGCG), are dietary phytochemicals with anti-inflammatory and antitumor effects [1]. The mechanisms underlying their effects may involve induction of antioxidant enzymes and suppression of pro-inflammatory cytokine activity. As JAK/STAT3 pathway is critical in inflammation and carcinogenesis and constitutively active in most tumor but not in normal cells [2], in the present study we examined the chemopreventive effects of quercetin and EGCG on this pathway using cholangiocarcinoma (CCA) cells treated with pro-inflammatory cytokine mixture (100 ng/mL IL-6 + 100 ng/mL IFN- γ) for mimicking inflammatory condition.



Methods

The effect of quercetin and EGCG on inducible JAK/STAT3 signaling pathway in CCA cells, KKU-100, was assessed by determination of p-STAT3 protein using western blot analysis. The mRNA expression of iNOS was determined by real-time PCR with specific primers.

RESULTS




Fig. 1 Effects of cytokine mixture on JAK/STAT3 pathway
cytokine mixture treatment could activate the STAT3 signaling cascade in CCA cells demonstrated by increased levels of phosphorylated-STAT3 protein (p-STAT3) in whole cell lysates




Fig. 2 Effects of quercetin and EGCG on cytokine-induced STAT3 phosphorylation

Cells were pre-incubated for 2 h with quercetin or EGCG, and then were treated with cytokine combination for 1 h. After that whole cell lysates were prepared for western blotting.
Pre-treatment of cells with quercetin and EGCG dose-dependently suppressed the STAT3 activation.

CONCLUSION

Quercetin and EGCG suppresses inducible JAK/STAT3 signaling cascade in CCA cells leading to the down-regulation of iNOS which is a known critical molecule involving in inflammation and carcinogenesis processes. The results provide evidence supporting the use of these two flavonoids as chemopreventive agents for cholangiocarcinoma.

RESULTS




Fig. 3 Effects of cytokine mixture in the presence or absence

of quercetin or EGCG on iNOS mRNA expression

Cells were pre-incubated for 2 h with quercetin or EGCG, and then were treated with cytokine combination for 3 h. After that mRNA was extracted for real-time PCR. * $p < 0.05$ compared to cytokine treatment alone.

Quercetin and EGCG could significantly inhibit the cytokine-mediated up-regulation of iNOS expression in a dose response manner.




Fig. 4 Effects of specific JAK inhibitors on cytokine-induced iNOS expression

Cells were pre-incubated for 2 h with AG490 (JAK2 inhibitor) or piceatannol (JAK1 inhibitor), and then were treated with cytokine combination for 3 h. After that mRNA was extracted for real-time PCR. * $p < 0.05$ compared to cytokine treatment alone.

Piceatannol and AG490 treatment dose-dependently suppressed cytokine-dependent induction of iNOS expression.

Acknowledgements

This work was supported by the Thailand Research Fund, Office of the Higher Education Commission, the National Research University Project and Office of the Higher Education Commission through SHeP-GMS of Khon Kaen University, Thailand.

References

- Garcia-Lafuente, A. et al. (2009). *Inflamm Res* 58(9): 537-52.
- Hodge, R. et al. (2005). *Eur J Cancer* 41(16): 2502-12.