

บทคัดย่อ

รหัสโครงการ: RSA4780018

ชื่อโครงการ: บทบาทในการต้านอนุมูลอิสระของเมลานินและผลของเมลานินในการป้องกันการเสื่อมสลายของเซลล์ประสาท

ชื่อนักวิจัยและสถานที่ทำงาน: รองศาสตราจารย์บัณฑิต เจตน์สว่าง

โครงการวิจัยชีววิทยาระบบประสาทและพฤติกรรม สถาบันวิจัยและพัฒนาวิทยาศาสตร์และเทคโนโลยี มหาวิทยาลัยมหิดล ศาลายา จ. นครปฐม 73170

E-mail Address: grbcs@mahidol.ac.th

ระยะเวลาโครงการ: 3 ปี (31 สิงหาคม 2547 - 30 สิงหาคม 2550)

การศึกษานี้ต้องการที่จะศึกษาหาสาเหตุและกลไกในระดับเซลล์และโมเลกุลที่อนุมูลอิสระก่อให้เกิดการเสื่อมสลายของเซลล์ประสาทและเส้นเลือดสมอง รวมทั้งผลในการเป็นสารต้านอนุมูลอิสระของเมลานินในการป้องกันการเสื่อมสลายนี้นี้ ผลจากการทดลองพบว่า hydrogen peroxide สามารถเหนี่ยวนำให้เกิดการเสื่อมสลายของเซลล์ประสาทโดปามีนชนิด SH-SY5Y มีการเพิ่มขึ้นของปริมาณ NF- κ B phosphorylation (pNF- κ B) อย่างมีนัยสำคัญและเหนี่ยวนำให้มีการ translocation ของ pNF- κ B จาก cytoplasm เข้าไปที่บริเวณรอบๆนิวเคลียสและในนิวเคลียสของเซลล์เพิ่มมากขึ้นอีกด้วยโดยผลดังกล่าวสามารถยับยั้งได้ด้วยเมลานิน นอกจากนี้ hydrogen peroxide ยังมีผลในการเพิ่มปริมาณของโปรตีน Bax และ Bcl-2 โดยเซลล์ที่ได้รับ hydrogen peroxide อย่างเดียวมีอัตราการเสื่อมสลายของเซลล์สูงการเพิ่มขึ้นของ Bax จะมากกว่า Bcl-2 แต่ในทางตรงกันข้ามเซลล์ที่ได้รับ hydrogen peroxide และเมลานินด้วยเซลล์มีอัตราการเสื่อมสลายของเซลล์ลดลงมีการลดลงของปริมาณของ Bax แต่ยังคงมีการเพิ่มขึ้นของปริมาณ Bcl-2 และในการศึกษานี้ก็ได้ทำการทดสอบและพบว่าสารที่ยับยั้งการทำงานของ caspase enzyme สามารถป้องกันการเสื่อมสลายของเซลล์ประสาทโดย hydrogen peroxide ได้ เมื่อทำการศึกษาผลของสารพิษ MPP⁺ พบว่า MPP⁺ สามารถเหนี่ยวนำให้เกิดการเสื่อมสลายของเซลล์ประสาทโดปามีนชนิด SK-N-SH ได้และยังสามารถเหนี่ยวนำให้มีการเพิ่มขึ้นของปริมาณ c-Jun phosphorylation เพิ่มการทำงานของ caspase enzyme และมีการย่อยสลาย DNA fragmentation 45 เพิ่มมากขึ้นอีกด้วย ซึ่งผลดังกล่าวข้างต้นของ MPP⁺ สามารถยับยั้งได้ด้วยเมลานิน เมื่อทำการศึกษาผลของเมลานินในการป้องกันการเสื่อมสลายของเส้นเลือดสมองจากสภาวะ oxidative stress พบว่าเมลานินสามารถลดการเพิ่มขึ้นของเอนไซม์ endothelial nitric oxide synthase และลดการเพิ่มขึ้นของ calcium/calmodulin-dependent protein kinase II phosphorylation ในเส้นเลือดสมองว้ที่ถูกเหนี่ยวนำให้เกิดสภาวะ oxidative stress จาก hydrogen peroxide ผลจากการศึกษาวิจัยแสดงให้เห็นถึงคุณสมบัติในการเป็นสารป้องกันการเสื่อมสลายของเซลล์ประสาทและเส้นเลือดสมองจากสภาวะ oxidative stress ของเมลานินซึ่งสภาวะดังกล่าวสามารถเหนี่ยวนำให้มีการกระตุ้นการทำงานของ NF- κ B, c-Jun-N-terminal kinases, CaMKII และ caspase-

dependent signaling ข้อมูลและความรู้ที่ได้จากการศึกษาวิจัยในครั้งนี้สามารถนำไปประยุกต์ใช้เพื่อการศึกษาวิจัยที่เกี่ยวข้องกับเมลานินกับสภาวะ oxidative stress และการตายของเซลล์ประสาทได้ต่อไปในอนาคต

คำหลัก: เมลานิน, hydrogen peroxide, MPP^+ , เซลล์ประสาท, เส้นเลือดสมอง

Abstract

Project Code: RSA4780018

Project Title: The antioxidative role of melatonin: The protective effects of melatonin in the process of neurodegeneration

Investigator: Associate Professor Banthit Chetsawang
Neuro-Behavioural Biology Center, Institute of Science and Technology for Research and Development, Mahidol University, Salaya, Nakhonpathom 73170

E-mail Address: grbcs@mahidol.ac.th

Project Period: 3 years (31 August 2004 - 30 August 2007)

Neurodegenerative diseases are illnesses associated with high morbidity and mortality with few, or no effective, options available for their treatment. The direct cause of neuronal cell loss has not been clearly understood. In addition, the neuroprotective effect of melatonin has been observed both in vivo and in vitro. The objective of this research, therefore, was to better understand the cellular mechanisms of neuronal cell degeneration induced via oxidative stress and neurotoxin. Taken together, the protective roles of melatonin on this cell death have also been studied. In the present study, the effect of melatonin on hydrogen peroxide (H_2O_2) and 1-methyl, 4-phenyl, pyridinium ion (MPP^+) induced neuronal cell degeneration in human dopaminergic neuroblastoma cultured cells were investigated. The results showed that H_2O_2 significantly decreased cell viability and melatonin reversed the toxic effects of H_2O_2 . An inhibition of caspase enzyme activity by Ac-DEVD-CHO, a caspase-3 inhibitor, significantly increased cell viability in H_2O_2 -treated cells. The phosphorylation of transcription factors, nuclear factor kappa B (NF- κ B) was increased in H_2O_2 -treated cells and this effect was abolished by melatonin. Translocation of phosphorylated NF- κ B to perinuclear and nuclear sites, estimated using immunofluorescence, occurred to a greater extent in H_2O_2 -treated cells than in untreated control cells and again this effect was abolished by melatonin. In addition, induction of Bcl-2 and Bax proteins was demonstrated in SH-SY5Y cultured cells treated with H_2O_2 , whereas the induction of Bax but not Bcl-2 was diminished by melatonin. Accordingly, MPP^+ significantly decreased cell viability. By contrast, an induction of phosphorylation of c-Jun, activation of caspase-3 enzyme activity, cleavage of DNA fragmentation factors 45 and DNA fragmentation were observed in MPP^+ -treated SK-N-SH cultured cells. In order to elaborate the functional significance of

melatonin in cerebral blood vessels, H₂O₂-induced induction in endothelial nitric oxide synthase (eNOS) protein level and phosphorylation of calcium/calmodulin-dependent protein kinase II (CaMKII) were demonstrated in the bovine isolated cerebral arteries with these effect being abolished by melatonin. These results demonstrate the cellular mechanisms of neuronal degeneration induced via NF- κ B, c-Jun-N-terminal kinases, CaMKII and caspase-dependent signaling, and the potential role of melatonin on protection of neuronal cell death induced by oxidative stress and neurotoxin. Furthermore, the regulatory role of melatonin in physiology of the cerebral vessels was also demonstrated. In light of these finding, it is possible that the neuroprotective effect of melatonin, may offer a means of treating neuronal degeneration and disease.

Keywords: Melatonin, Hydrogen peroxide, MPP⁺, Neuronal cells and Cerebral blood vessels