



# รายงานวิจัยฉบับสมบูรณ์

## โครงการ

1. การประเมินเทคโนโลยีและการใช้เทคโนโลยีอย่างเหมาะสมเพื่อคุณภาพการบริการ  
ในโรงพยาบาล (Technology assessment and quality use in hospital setting)
2. โครงการสหสาขาวิชาเพื่อแก้ปัญหาสาธารณสุขที่สำคัญของประเทศ  
(Multi disciplinary research on important public health problem)

โดย

ศาสตราจารย์ นายแพทย์จิตร สิทธีอมร

14 พฤษภาคม 2554

สัญญาเลขที่ RTA/08/2540

# รายงานวิจัยฉบับสมบูรณ์

## โครงการ

1. การประเมินเทคโนโลยีและการใช้เทคโนโลยีอย่างเหมาะสมเพื่อคุณภาพการบริการ  
ในโรงพยาบาล (Technology assessment and quality use in hospital setting)
2. โครงการสหสาขาวิชาเพื่อแก้ปัญหาสาธารณสุขที่สำคัญของประเทศ  
(Multi disciplinary research on important public health problem)

ศาสตราจารย์ นายแพทย์จิตร สิทธีอมร  
จุฬาลงกรณ์มหาวิทยาลัย

สนับสนุนโดย สำนักงานกองทุนสนับสนุนการวิจัย

(ความเห็นในรายงานนี้เป็นของผู้วิจัย สกว. ไม่จำเป็นต้องเห็นด้วยเสมอไป)

## บทคัดย่อ

โครงการทุนส่งเสริมการวิจัย เมธีวิจัยอาวุโส สกว. ศ. นพ. จิตร สิทธิอมร ประกอบด้วย 2 กลุ่มโครงการ คือ 1) กลุ่มโครงการการประเมินเทคโนโลยีและการใช้เทคโนโลยีอย่างเหมาะสมเพื่อคุณภาพการบริการในโรงพยาบาล และ 2) กลุ่มโครงการสหสาขาวิชาเพื่อแก้ปัญหาสาธารณสุขที่สำคัญของประเทศ โดยมีเป้าหมายรวม คือ การพัฒนาศักยภาพนักวิจัยในการทำงาน และบริหารงานวิจัย เพื่อนำกระบวนการวิจัยมาแก้ไขปัญหาด้านสาธารณสุขในประเทศ มีโครงการวิจัยที่ได้รับทุนสนับสนุน รวมทั้งสิ้น 17 โครงการ จากหลากหลายหน่วยงานในประเทศ

กลุ่มโครงการประเมินเทคโนโลยีเพื่อการบริการระดับโรงพยาบาล ประกอบด้วยโครงการวิจัย 10 โครงการ อันได้แก่ 1) The cost effectiveness of early detection of bancroftian filariasis in Myanmar Migrants using antigen-capsule ELISA, IgG4 antibody assays and PCR technique ระยะที่ 1 และ ระยะที่ 2 2) รูปแบบการตรวจพาหะธาลัสซีเมียฮีโมโกลบินผิดปกติ 3) ประสิทธิภาพของการดูแลร่วมกันสำหรับผู้ป่วยโรคลมชักในจังหวัดนครราชสีมา 4) การศึกษาเพื่อหาปัจจัยเสี่ยงของผู้ป่วยกล้ามเนื้อหัวใจตายเฉียบพลันกับกลุ่มควบคุม 5) การศึกษาครอบครัวไหลตาย 6) แนวโน้มการช่วยคืนชีพในผู้ป่วยระยะสุดท้าย และ ประสิทธิภาพของการทำพินัยกรรมชีวิต เรื่องการช่วยคืนชีพในผู้ป่วยระยะสุดท้ายในโรงพยาบาลมหาวิทยาลัยเชียงใหม่ 7) ทศนคติต่อการทำพินัยกรรมชีวิตและการให้ข้อมูลเรื่องผลของการช่วยคืนชีพในผู้ป่วยอายุรกรรม โรงพยาบาลมหาวิทยาลัย 8) โครงการตรวจสุขภาพประจำปี ของผู้สูงอายุในเขตชนบทของจังหวัดขอนแก่น 9) การศึกษาความเข้าใจ ความน่าเชื่อถือ ความสมบูรณ์ และความไวต่อการเปลี่ยนแปลงของแบบสอบถาม ฉบับภาษาไทย Health Assessment Questionnaire สำหรับผู้ป่วยโรคข้ออักเสบ และ 10) การศึกษาความชุก ภาวะความรุนแรง ปัจจัยและผลที่เกิดขึ้นจากเนื้อตายในผู้ป่วยถูกงูเห่าและงูกะปะกัด และ การศึกษาการป้องกันภาวะเนื้อตายที่เกิดขึ้นในสัตว์ทดลอง

กลุ่มโครงการสหสาขาวิชาเพื่อแก้ปัญหาสาธารณสุขที่สำคัญของประเทศ ประกอบด้วยโครงการวิจัย 7 โครงการ อันได้แก่ 1) ผลของการปรับเปลี่ยนการบริโภคมาเป็นข้าวกล้อง งา ผัก ผลไม้ ตามฤดูกาลของประชาชนในเขตเทศบาลจังหวัดสุโขทัย 2) ความต้องการและ ปัญหาด้านสุขภาพ การเข้าถึงบริการ และ คุณภาพชีวิตของวัยรุ่นกับการสร้างความร่วมมือ 3) การศึกษาผลกระทบของฝุ่นละอองขนาดเล็กต่อสุขภาพของประชาชนในจังหวัดเชียงใหม่ 4) สื่อสร้างสรรค์จากเพื่อนสู่เพื่อนเพื่อส่งเสริมพฤติกรรมป้องกันการเสพติดของวัยรุ่นในชุมชนแออัด 5) การมีส่วนร่วมของประชาสังคมในกลยุทธ์เพื่อส่งเสริมการใช้ยาอย่างเหมาะสมสำหรับการรักษาโรคติดต่อที่ระบบทางเดินหายใจส่วนบนของผู้ใหญ่ในระดับครัวเรือนและชุมชน: กรณีศึกษาชุมชนแออัดในกรุงเทพมหานคร 6) การวิเคราะห์ความเสี่ยงในการเกิดโรคหัวใจ ชัก และการเสียชีวิตจากการศึกษาระยะยาวในชุมชนแออัดคลองเตย และ 7) การติดต่อของเชื้อหวัดสายพันธุ์ A ระหว่างคนและสัตว์ในประเทศไทย

โครงการทุนส่งเสริมการวิจัย เมธีวิจัยอาวุโส สกว. ศ. นพ. จิตร สิทธิอมร ได้กำหนดตัวชี้วัดของการบรรลุเป้าหมาย อันได้แก่ ผลงานวิจัยที่ผลิต การตีพิมพ์ในวารสารระดับชาติและนานาชาติ การที่นักวิจัยสามารถเขียนโครงการขอทุนด้วยตัวเอง การทำวิจัยร่วมระหว่างสถาบันและระหว่างนักวิจัยต่างสาขา ตลอดจนการที่ผลงานวิจัยได้ถูกนำไปใช้เป็นแนวทางในการวางนโยบายระดับนโยบายสาธารณะ และในระดับสถานบริการ พบว่าโครงการส่วนใหญ่ที่ได้รับงบประมาณสนับสนุน ได้ดำเนินการบรรลุตามวัตถุประสงค์ และตัวชี้วัดที่ได้

กำหนดไว้ พบว่า จากจำนวนโครงการ 17 โครงการ มีการตีพิมพ์ในวารสารรวมทั้งสิ้น 16 บทความ เป็นการตีพิมพ์ในวารสารระดับนานาชาติ จำนวน 14 เรื่อง ตีพิมพ์ในวารสารระดับชาติ 2 จำนวน มีการนำเผยแพร่ผลงานวิชาการในการประชุมวิชาการ และการสัมมนาทั้งในระดับนานาชาติ และระดับชาติ รวม 5 เรื่อง มีการต่อยอดโครงการวิจัย โดยนักวิจัยสามารถเขียนโครงการขอทุนด้วยตัวเองได้ 4 เรื่อง มีการทำวิจัยร่วมระหว่างสถาบันและระหว่างนักวิจัยต่างสาขา จำนวน 4 เรื่อง มีนักวิจัย ที่มีความชำนาญ หรือเป็นผู้เชี่ยวชาญเพิ่มขึ้น 8 คน ในด้านการแพทย์ การพยาบาล การสาธารณสุข นักวิจัยจำนวน 3 คน ได้รับเข้าเป็นอาจารย์ในสถาบันการศึกษาของรัฐ ได้แก่ นพ.ธนินทร์ อัครวิเชียรจินดา เป็นอาจารย์คณะแพทยศาสตร์ จุฬาลงกรณ์มหาวิทยาลัย ดร.รัตนา สำโรงทอง เป็นอาจารย์วิทยาลัยวิทยาศาสตร์สาธารณสุข จุฬาลงกรณ์มหาวิทยาลัย และ ดร.ศิริตรี สุทธจิตต์ เป็นอาจารย์คณะเภสัชศาสตร์ มหาวิทยาลัยมหาสารคาม

นอกจากนี้ยังมีผลงานวิจัยจำนวนหนึ่ง ได้ถูกนำไปใช้เป็นแนวทางในตรวจโรคบางอย่างในโรงพยาบาล เช่น การพัฒนารูปแบบการตรวจพาหะธาลัสซีเมียฮีโมโกลบินผิดปกติและการศึกษาความคุ้มทุนของการตรวจ bancroftian filariasis ด้วยวิธีการต่างๆ และบางโครงการได้นำไปใช้เป็นแนวทางในแก้ปัญหาด้านสาธารณสุข ในเรื่องสุขภาพวัยรุ่น การส่งเสริมด้านโภชนาการ การใช้ยาอย่างเหมาะสม ได้มีนำผลการศึกษา ไปใช้เป็นแนวทางในการวางนโยบายระดับสาธารณะ เช่น เรื่องสุขภาพวัยรุ่นในชุมชนแออัด การใช้ยาอย่างเหมาะสมในชุมชน เป็นต้น

**คำสำคัญ** การใช้เทคโนโลยีอย่างเหมาะสม การบริการในโรงพยาบาล สหสาขาวิชา ปัญหาสาธารณสุข

## Abstract

The Thailand Research Fund (TRF) has supported research projects through Prof. Chitr Sitthi-amorn, TRF Senior Research Scholar. The projects were classified into 2 clusters: 1) Technology assessment and appropriate technology for quality of care of the hospitals and 2) Multidisciplinary research for solving public health problems. These research projects aimed to strengthen research capacity and research management for solving public health problems Thailand. There were 17 research projects undertaken by various institutions.

The cluster of the technology assessment projects to improve quality of care in the hospitals composed of 10 sub-projects:

- 1) The cost effectiveness of early detection of bancroftion filarasis in Myanmar Migrants using antigen-capsule ELISA, IgG4 antibody assays and PCR technique (Phase I, II)
- 2) Screening models for Thalassemia and Hemoglobin patients
- 3) The effectiveness of shared care for patients with epilepsy in Nakhonratchasima province
- 4) INTER-HEART: a global study of risk factors in Acute Myocardial Infraction
- 5) A study of Unexpected Death Syndrome
- 6) Trend of cardiopulmonary resuscitation patients with terminal illness in Chiang Mai
- 7) Attitude towards advance directives and the impact of prognostic information on the preference for cardiopulmonary resuscitation on medical inpatients in CMU Hospital, Thailand
- 8) Annual health examination among elderly in rural areas of Khon Kaen Province
- 9) Comprehensibility, reliability, validity, and responsiveness of the Thai version of the Health Assessment Questionnaire in Thai patients with rheumatoid arthritis
- 10) Prevalence, severity, determinants and consequences of tissue necrosis among victims envenomed by Naja kaouthia (Thai Cobra) and Calloselasma rhodostoma (Malayan Pit Viper) and its prevention in an experimental model

The cluster of the multidisciplinary research projects which sought to identify solutions to public health problems consisted of 7 sub-projects:

- 1) Healthy food choices in accordance with seasonal diets in Sukothai province
- 2) Adolescent health needs and accessibility of services in congested community
- 3) A study of effect of PM<sub>10</sub> and PM<sub>2.5</sub> of population in Chiang Mai
- 4) Friend media to prevent drug addict of adolescent in congested community
- 5) Partnership of civil society in a strategy to reduce the use of antibiotics in the treatment of adults with Upper Respiratory Infection (URIs) from viral origins at the household and community levels: A comparison study in congested community, Bangkok Thailand

- 6) An analysis of population cohort: Risk of Hypertension, Stroke, and Death of residence in Klong Toey Slum
- 7) Influenza A infections at the human animal interface.

The indicators of the achievement of these research projects included several research reports, national and international publications, fundraising and the utilization of findings to inform public policy and improve the quality of care. The results shown that many sub-projects supported by the TRF Senior Research Scholar (Prof. Chitr Sitthi-amorn) met the objectives. There were 16 publications from 17 research projects: 14 international publications and two national publications. In addition, five research projects were presented in international and national conferences; four research projects were awarded additional external funding to explore findings further; four projects have initiated institutional collaborations; eight principal investigators have become recognized experts in their area, and three principal investigators have since been appointed as university lecturers.

Furthermore, research findings of some TRF projects were applied to diagnose disease in hospitals, for example, the development of Thalassemia diagnosis and determination of the cost effectiveness of diagnosis bancroftion filarasis. In addition, some research projects have identified alternative methods to address community health issues, including improving adolescent health, improving nutritional status, better ensuring rational use of drugs and forming public health policy.

**Keywords:** Appropriate technology, Hospital care, Multidisciplinary, Health problem

# สารบัญ

	หน้า
บทคัดย่อภาษาไทย	i
บทคัดย่อภาษาอังกฤษ	iii
สารบัญ	v
รายงานผลการวิจัย	
กลุ่มโครงการประเมินเทคโนโลยีเพื่อการบริหารระดับโรงพยาบาล	
1 The cost effectiveness of early detection of bancroftion filarasis in Myanmar Migrants using antigen-capsule ELISA, IgG4 antibody assays and PCR technique ระยะที่ 1 และ ระยะที่ 2	1
2 รูปแบบการตรวจพาหะธาลัสซีเมียฮีโมโกลบินผิดปกติ	3
3 ประสิทธิภาพของการดูแลร่วมกันสำหรับผู้ป่วยโรคลมชักในจังหวัดนครราชสีมา	5
4 การศึกษาเพื่อหาปัจจัยเสี่ยงของผู้ป่วยกล้ามเนื้อหัวใจตายเฉียบพลันกับกลุ่มควบคุม	6
5 การศึกษาครอบครัวไหลตาย	8
6 แนวโน้มการช่วยคืนชีพในผู้ป่วยระยะสุดท้าย และประสิทธิผลของการทำพินัยกรรมชีวิต เรื่องการช่วยคืนชีพในผู้ป่วยระยะสุดท้ายในโรงพยาบาลมหาวิทยาลัยเชียงใหม่	10
7 ทศนคติต่อการทำพินัยกรรมชีวิตและการให้ข้อมูลเรื่องผลของการช่วยคืนชีพในผู้ป่วยอายุรกรรม โรงพยาบาลมหาวิทยาลัย	11
8 โครงการตรวจสุขภาพประจำปี ของผู้สูงอายุในเขตชนบทของจังหวัดขอนแก่น	12
9 การศึกษาความเข้าใจ ความน่าเชื่อถือ ความสมบูรณ์ และความไวต่อการเปลี่ยนแปลงของแบบสอบถาม ฉบับภาษาไทย Health Assessment Questionnaire สำหรับผู้ป่วยโรคข้ออักเสบ	13
10 การศึกษาความชุก ภาวะความรุนแรง ปัจจัยและผลที่เกิดขึ้นจากเนื้อตายในผู้ป่วยถูกงูเห่าและงูกะปะกิด และการศึกษาการป้องกันภาวะเนื้อตายที่เกิดขึ้นในสัตว์ทดลอง	14
กลุ่มโครงการสหสาขาวิชาเพื่อแก้ปัญหาสาธารณสุขที่สำคัญของประเทศ	
1 ผลของการปรับเปลี่ยนการบริโภคมาเป็นข้าวกล้อง งาม ผัก ผลไม้ ตามฤดูกาลของประชาชนในเขตเทศบาลจังหวัดสุโขทัย	16
2 ความต้องการและ ปัญหาด้านสุขภาพ การเข้าถึงบริการ และคุณภาพชีวิตของวัยรุ่นกับการสร้างความร่วมมือ	18
3 การศึกษาผลกระทบของฝุ่นละอองขนาดเล็กต่อสุขภาพของประชาชนในจังหวัดเชียงใหม่	20
4 สื่อสร้างสรรค์จากเพื่อนสู่เพื่อนเพื่อส่งเสริมพฤติกรรมป้องกันการเสพยาบ้าของวัยรุ่นในชุมชนแออัด	24
5 การมีส่วนร่วมของประชาสังคมในกลยุทธ์เพื่อส่งเสริมการใช้ยาอย่างเหมาะสมสำหรับการรักษาโรคติดต่อที่ระบบทางเดินหายใจ ส่วนบนของผู้ใหญ่ในระดับครัวเรือนและชุมชน: กรณีศึกษาชุมชนแออัดในกรุงเทพมหานคร	26
6 การวิเคราะห์ความเสี่ยงในการเกิดโรคหัวใจ ชัก และการเสียชีวิตจากการศึกษาระยะยาวในชุมชนแออัดคลองเตย	28
7 การติดต่อของเชื้อหวัดสายพันธุ์ A ระหว่างคนและสัตว์ในประเทศไทย	29
สรุปผลลัพธ์ (Output) ที่ได้จากโครงการ	34
ภาคผนวก ผลงานตีพิมพ์	37

## กลุ่มโครงการประเมินเทคโนโลยีเพื่อการบริหารระดับโรงพยาบาล 10 โครงการ

**ชื่อโครงการ 1** The cost effectiveness of early detection of bancroftion filarasis in Myanmar Migrants using antigen-capsule ELISA, IgG4 antibody assays and PCR technique

**นักวิจัยหลัก** ศ.พญ.สุรางค์ ไตรธีระประภาพ

**นักวิจัยในทีม** นพ.สมชาย วงศ์ยงเจริญ

**ที่ปรึกษา** ศ.นพ.จิตร สิทธิอมร

### หลักการและเหตุผล

ปัจจุบันการตรวจหาผู้ป่วยโรคเท้าช้างทำได้หลายวิธี วิธีที่นิยมใช้กันคือการใช้ฟิล์มส่องหาเชื้อ Microfilaria (Blood smear technique) ซึ่งให้ผลบวกต่อการตรวจ 3% (7/202) ในขณะที่วิธีตรวจ w. bancroftispecific circulating antigen (Ag) การตรวจ anti-filarial Ig4 antibody (Ab) และวิธี Polymerase Chain Reaction (PCR) ให้ผลบวกต่อการตรวจ 16% (34/202) และ 6% (13/202) ตามลำดับ โดยการตรวจด้วยวิธี Blood smear technique ยังไม่สามารถสรุปได้ว่าการตรวจไม่พบเชื้อ Microfilaria (Mf) จะไม่เป็นโรค เนื่องจากในการศึกษายังคงพบผู้ป่วยที่ตรวจไม่พบเชื้ออยู่จำนวนหนึ่ง ในการศึกษาในกลุ่มแรงงานชาวพม่าในเขตพื้นที่ อ.แม่สอด จ.ตาก จำนวน 202 ราย จึงกำหนดให้การตรวจพบ Microfilaria positive (Mf+) และ/หรือ Antigen positive (Ag+) เป็น Reference method ของการวินิจฉัยโรคเท้าช้าง ผลการศึกษาพบความชุกของโรค 17% (35/202) สำหรับการทดสอบความไวการตรวจหา Ag มีความไวสูงสุด (97%) รองลงมาเป็นการตรวจหา Ab (86%) วิธี PCR (37%) และวิธี Blood smear technique มีความไวในการตรวจพบผู้ป่วยต่ำสุด (20%) ในขณะที่ความจำเพาะต่อโรคของการตรวจหา Ab ต่ำกว่าวิธีอื่น (46%) ในการวิเคราะห์ต้นทุน-ประสิทธิผลแสดงให้เห็นว่าการใช้วิธี PCR โดยมีต้นทุน 52.99 บาทต่อคน 75.12 บาทต่อคน และ 205.03 บาทต่อคนตามลำดับ ดังนั้นการตรวจหา Ag จึงเป็นวิธีที่ cost effectiveness ที่สุด แม้จะมีค่าใช้จ่ายในการตรวจสูงกว่าวิธี Blood smear technique และการตรวจหา Ab ก็ตาม

### วัตถุประสงค์

ศึกษาความไวและความจำเพาะของวิธีวินิจฉัยโรคเท้าช้างซึ่งเกิดจากเชื้อ Wuchereria bancrofti (W. bancrofti) โดยใช้ปฏิกิริยาลูกโซ่โพลีเมอร์เรส (Polymerase Chain Reaction, PCR) และการตรวจระดับ Specific circulating antigen (Mf) ในเลือดต่ำมากไม่สามารถตรวจพบในเลือดได้โดยวิธีการดูด้วยกล้องจุลทรรศน์ และศึกษาถึงต้นทุน-ประสิทธิผลของวิธีวินิจฉัยดังกล่าว

### ขั้นตอนการศึกษา

1. การเก็บตัวอย่างสำหรับการวิเคราะห์ทางปฏิกิริยาลูกโซ่โพลีเมอร์เรสและการตรวจสอบทางน้ำเหลือง
2. ตัวอย่างสำหรับการตรวจทางปรสิตวิทยา
3. การตรวจด้วยวิธีปฏิกิริยาลูกโซ่โพลีเมอร์เรส
4. การตรวจ W. bancrofti-specific circulating antigen
5. การตรวจ anti-filarial IgG4 antibody

## 6. การวิเคราะห์ข้อมูล

### ผลการศึกษา

จากการศึกษาต้นทุน-ประสิทธิผลของวิธีการตรวจหาผู้ป่วยโรคเท้าช้าง โดยการตรวจหาเชื้อ Mf ด้วยวิธี Blood smear technique การตรวจหา Ag และ Ab รวมทั้งการตรวจหาด้วยวิธี PCR ในกลุ่มแรงงานพม่า จำนวน 202 รายในจ.ตาก เป็นเพศชาย 47 ราย (23%) และเพศหญิง 155 ราย (77%) พบว่าการตรวจหา Ab ให้ผลบวกต่อการตรวจสูงถึง 59% รองลงมาเป็นการตรวจหา Ag ให้ผลบวกต่อการตรวจ 16% วิธี Blood smear technique ให้ผลบวกต่อการตรวจ 6% และวิธี PCR ให้ผลบวกต่อการตรวจต่ำสุดคือ 3% เมื่อจำแนกตามช่วงอายุ พบว่ากลุ่มที่ให้ผลบวกต่อการตรวจหา Ag การตรวจหา Ab และวิธี PCR สูงสุดคือ กลุ่มช่วงอายุ >15-30 ปี ในขณะที่การตรวจหาเชื้อ Mf (Blood smear technique) ให้ผลบวกต่อการตรวจสูงสุดในกลุ่มช่วงอายุ >30-45 ปี

### การนำผลไปใช้ประโยชน์

การวิเคราะห์ต้นทุน-ประสิทธิผลได้ใช้ Mf+ และ/หรือ Ag+ เป็น reference method เนื่องจากแม้ว่า Mf จะเป็นวิธีดั้งเดิมที่ใช้กันแต่ไม่สามารถวินิจฉัยผู้ป่วยที่มีเชื้อจำนวนน้อย หรือผู้ป่วยที่อยู่ระยะแรกเริ่ม (prepatient) ได้ ส่วนการตรวจหา Ab นั้น แม้ว่าจะวินิจฉัยผู้ป่วยได้จำนวนมาก แต่ไม่ได้แยกว่าผู้ป่วยเป็นโรคอยู่หรือเคยเป็นโรคแต่หายแล้ว ในขณะที่วิธี PCR มีราคาตรวจ/case สูงที่สุด แต่วินิจฉัยผู้ป่วยได้จำนวนไม่มากนัก ดังนั้นการตรวจหา Ag ซึ่งบ่งภาวะ active infection ร่วมกับวิธี Mf ซึ่งถือเป็น Gold standard จึงเหมาะสมที่จะใช้ในการวิเคราะห์ต้นทุน-ประสิทธิผล ซึ่งเป็นต้นทุนที่คำนวณเพื่อให้ได้วิธีที่มีประสิทธิผลที่สุดในที่นี้หมายถึงความสามารถในการตรวจวินิจฉัยผู้ป่วยได้ถูกต้อง (true positive and true negative) โดยเสียค่าตรวจต่อ case ถูกที่สุด

อันึงการใช้วิธีการตรวจอื่นร่วมกับ Gold standard ส่งผลให้ค่า sensitivity, specificity, positive predictive value (PPV), negative predictive value (NPV) และ accuracy รวมถึงต้นทุนการตรวจต่อ case เปลี่ยนตาม reference method ที่ใช้ การใช้ Mf+ และ/หรือ Ab+ เป็นตัวกำหนดการวินิจฉัยโรคว่าเป็นโรคเท้าช้างทำให้เสียค่าตรวจต่อ case ลดลงเป็น 28.21 บาทต่อ case แต่ก็มี false positive สูง ในขณะที่การวินิจฉัยโรคโดยใช้ Mf+ และ/หรือ PCR+ และการใช้ร่วมกันทั้ง 4 วิธี (Mf/Ag/Ab/PCR) มีค่าตรวจต่อ case 184.53 บาทและ 697.24 บาท ซึ่งมีราคาสูงกว่าวิธีตรวจอื่น ดังนั้นการใช้ Mf+ และ/หรือ Ag+ เป็นตัวกำหนดการวินิจฉัยโรคจึงน่าจะคุ้มทุน และมีประสิทธิผลในการวินิจฉัยโรคได้ถูกต้องมากที่สุดโดยค่าตรวจต่ำสุดคือ 75.12 บาทต่อ case

### การนำไปใช้ประโยชน์และการเผยแพร่

ผลงานได้รับการเผยแพร่ในหน่วยงาน

## ชื่อโครงการไทย 2 รูปแบบการตรวจกรองหาพาหะของธาลัสซีเมียและฮีโมโกลบินผิดปกติ

## ชื่อโครงการอังกฤษ Screening models for Thalassemia and Hemoglobinopathies

นักวิจัยหลัก นพ.กิตติ ต่อจรัส

ที่ปรึกษา ศ.นพ.จิตร สิทธิอมร

### หลักการและเหตุผล

โรคเลือดจางธาลัสซีเมีย (thalassemia) เป็นปัญหาใหญ่และสำคัญทางด้านสาธารณสุขของประเทศ ไทย เนื่องจากเป็นโรคพันธุกรรมซึ่งมีอุบัติการณ์สูงมากในประเทศไทย ประมาณร้อยละ 20-30 ของประชากรมี ยีน  $\alpha$  - thalassemia ร้อยละ 3-9 มียีน  $\beta$ - thalassemia และพบยีนของฮีโมโกลบินผิดปกติ 2 ชนิด ฮีโมโกลบิน อี (Hb E) โดยเฉลี่ยพบประมาณร้อยละ 13 แต่ในชาวอีสานพบสูงถึงร้อยละ 30-40

คณะผู้วิจัยมีความประสงค์เสนอรูปแบบการตรวจกรองหาพาหะของธาลัสซีเมียและฮีโมโกลบินผิดปกติ วิธี OF test, DCIP (KKU-DCIP Clear) และ modified Hb inclusion โดยใช้การตรวจแบบ serial screening test ซึ่งสามารถประหยัดค่าใช้จ่าย มีขั้นตอนไม่ยุ่งยากสามารถทำได้ในโรงพยาบาลชุมชนและยังนำรูปแบบ การตรวจกรองที่เหมาะสมมาเป็นต้นแบบในการตรวจกรองสำหรับการป้องกันธาลัสซีเมียในประเทศไทยอย่าง มีประสิทธิภาพ

### วัตถุประสงค์การวิจัย

1. หา sensitivity ของ serial screening test ในการตรวจกรองหาพาหะของเบต้าธาลัสซีเมียอัลฟาธาลัสซีเมียและฮีโมโกลบินอี โดยวิธี OFT, DCIP และ modified Hb H inclusion test เปรียบเทียบกับวิธีมาตรฐาน
2. หารูปแบบที่เหมาะสมของการตรวจกรองผู้ที่เป็นพาหะของธาลัสซีเมียและฮีโมโกลบินผิดปกติ

### ขอบเขตการวิจัยและวิธีการวิจัย

การศึกษาแบบ diagnostic study กลุ่มประชากรเป้าหมายคือ กำลังพลของกองพลทหารบก ประชากรตัวอย่างได้แก่ กำลังพล และครอบครัวทหารราบที่ 11 ในจังหวัดฉะเชิงเทรา อายุตั้งแต่ 20 ปีขึ้นไป ทุกเพศ ระยะเวลาการศึกษา 1 ปี 6 เดือน ตั้งแต่เดือน มกราคม 2541 ถึง มิถุนายน 2542 โดยแบ่งการศึกษา เป็น 2 ส่วน คือ การตรวจกรองในภาคสนาม (Screening method) คืองานในภาคสนามเป็นการให้ความรู้เกี่ยวกับโรค การตรวจกรองและการป้องกันโรคโลหิตจางธาลัสซีเมีย ประชากรที่ได้รับข้อมูลเกี่ยวกับโรคและ มีความประสงค์จะเข้าร่วมโครงการจะได้รับการซักประวัติ ตรวจร่างกาย และเจาะเลือด และการตรวจหาพาหะ โดยวิธีมาตรฐาน (standard method) คือการเจาะเลือดจากภาคสนามจะนำมาทำการทดสอบยืนยันทาง ห้องปฏิบัติการที่หน่วยโลหิตวิทยา กองกุมารเวชกรรม การตรวจหาดัชนีเม็ดเลือดแดง ตรวจหาปริมาณ Hb E และ Hb A2 ตรวจหาชนิดของฮีโมโกลบิน ตรวจหาภาวะเลือดจางจากการขาดธาตุเหล็ก และการตรวจหา พาหะของอัลฟาธาลัสซีเมีย

### ประโยชน์ที่ได้จากการศึกษา

1. สามารถนำวิธีการตรวจกรองหาพาหะของธาลัสซีเมียและฮีโมโกลบินผิดปกติเป็นต้นแบบนำไปในการตรวจกรองระดับประเทศ
2. ใช้ประกอบกับแผนการควบคุมและป้องกันระดับประเทศได้อย่างมีประสิทธิภาพ

### การนำไปใช้ประโยชน์และการเผยแพร่

ผลงานได้รับการเผยแพร่ในหน่วยงาน

**ชื่อโครงการ ไทย 3 ประสิทธิภาพของการดูแลร่วมกันสำหรับผู้ป่วยโรคลมชักในจังหวัดนครราชสีมา**

**ชื่อโครงการอังกฤษ** The effectiveness of shared care for patients with epilepsy in Nakhonratchasima province

**นักวิจัยหลัก** นพ.ธนินทร์ อัสววิเชียรจินดา

**ที่ปรึกษา** ศ.นพ.จิตร สิทธิอมร, Prof. Edgar J. Love

### **หลักการและเหตุผล**

วัตถุประสงค์ของการศึกษา เพื่อที่จะประเมินประสิทธิภาพของระบบการดูแลร่วมกัน (Shared Care) สำหรับผู้ป่วยโรคลมชักในแง่ของการติดตามการรักษา การควบคุมอาการชัก คุณภาพชีวิต ความพึงพอใจของผู้ป่วย และการลดการดูแลรักษาที่ไม่เหมาะสม

**วิธีการวิจัย** การศึกษานี้เป็นการศึกษาเปรียบเทียบแบบสุ่มตัวอย่าง โดยสุ่มตัวอย่างโรงพยาบาลชุมชน การศึกษานี้ใช้เวลา 1 ปี โดยโรงพยาบาลชุมชนที่มีคุณสมบัติตามที่กำหนดจะถูกเชิญให้เข้าร่วมโครงการ โรงพยาบาลที่ยินดีเข้าร่วมโครงการจะถูกแบ่งกลุ่มออกเป็น 3 กลุ่ม จากนั้นในแต่ละกลุ่ม โรงพยาบาลจะถูกสุ่มให้เป็นโรงพยาบาลควบคุมและโรงพยาบาลที่มีการสอดแทรก จากนั้นผู้ป่วยในแต่ละโรงพยาบาลจะถูกเชิญให้เข้าร่วมโครงการ ผู้ป่วยที่มีคุณสมบัติตามที่กำหนด จะได้เข้าร่วมโครงการศึกษา

**ผลการศึกษา** โรงพยาบาลที่เข้าร่วมโครงการมีทั้งหมด 12 โรงพยาบาล และมีผู้ป่วยที่เข้าร่วมโครงการทั้งสิ้น 401 ราย ลักษณะพื้นฐานทางคลินิกของผู้ป่วยทั้งสองกลุ่มไม่มีความแตกต่างกันยกเว้นจำนวนผู้ป่วยที่รับประทานยากันชักหนึ่งตัวกับรับประทานยาอื่นร่วมด้วย แต่หลังจากแยกกลุ่มวิเคราะห์ตามความแตกต่างของลักษณะพื้นฐาน พบว่า ผู้ป่วยที่รับประทานยากันชักตัวเดียวในกลุ่มสอดแทรกมีอัตราการติดตามการรักษาได้สม่ำเสมอว่าผู้ป่วยในกลุ่มควบคุม แต่การควบคุมอาการชัก คุณภาพชีวิต และความพึงพอใจไม่มีความแตกต่างกันระหว่าง 2 กลุ่ม

**สำหรับสิ่งสอดแทรก** พบว่าผู้ป่วยประเมินว่า การให้ความรู้จากพยาบาลมีประโยชน์และสามารถนำไปใช้ในชีวิตประจำวันได้ และแพทย์ก็มีความรู้ในเรื่องการดูแลรักษาเพิ่มขึ้นและเป็นประโยชน์ต่อการปฏิบัติ การดูแลรักษาที่ไม่เหมาะสมก็ลดน้อยลงอย่างมีนัยสำคัญด้วย

**สรุป** การดูแลร่วมกัน (Shared Care) มีประสิทธิภาพในการทำให้ผู้ป่วยติดตามการรักษาสม่ำเสมอขึ้น โดยเฉพาะผู้ป่วยที่มีอาการชักไม่รุนแรง และสิ่งสอดแทรกก็เป็นประโยชน์อย่างมากสำหรับผู้ป่วยและแพทย์ในการดูแลรักษาผู้ป่วยโรคลมชัก ถ้าการศึกษานี้ดำเนินขึ้นเป็นประมาณ 2-5 ปีก็คิดว่า การควบคุมอาการชัก คุณภาพชีวิตและความพึงพอใจของผู้ป่วย น่าจะดีขึ้น

### **การนำผลไปใช้ประโยชน์/การเผยแพร่ผลงาน**

Asawavichienjinda, T., Sitthi-Amorn, C., & Tanyanont, W. (2002). Prevalence of epilepsy in rural Thailand: a population-based study. *J Med Assoc Thai*, 85(10), 1066-1073.

Asawavichienjinda, T., Sitthi-Amorn, C., & Tanyanont, W. (2003). Compliance with treatment of adult epileptics in a rural district of Thailand. *J Med Assoc Thai*, 86(1), 46-51.

**ชื่อโครงการ ไทย 4** การศึกษาเพื่อหาปัจจัยเสี่ยงของผู้ป่วยกล้ามเนื้อหัวใจตายเฉียบพลันกับกลุ่มควบคุม

**ชื่อโครงการอังกฤษ** INTER-HEART: a global study of risk factors in Acute Myocardial Infarction

**นักวิจัยหลัก:**

ศ.นพ.เกียรติชัย ภูริปัญญา	ศูนย์โรคหัวใจ โรงพยาบาลศิริราช
พญ.สมนพร บุญรัตเวช	ศูนย์โรคหัวใจ โรงพยาบาลจุฬาลงกรณ์
นพ.ศุภชัย ถนอมทรัพย์	ศูนย์โรคหัวใจ โรงพยาบาลรามาริบัติ
ศ.นพ.ปิยทัศน์ ทศนาวิวัฒน์	ศูนย์โรคหัวใจ มหาวิทยาลัยขอนแก่น
ผศ.นพ.วรวิทย์ จินตภากร	ศูนย์โรคหัวใจ มหาวิทยาลัยสงขลานครินทร์
นพ.บรรหาร กอนันต์กุล	ศูนย์ป้องกันและพัฒนาการรักษาโรคหัวใจ โรงพยาบาลภูมิพลอดุลยเดช
นพ.ปิยะ เกษมสุวรรณ	ศูนย์โรคหัวใจ วชิรพยาบาล
นพ.โสภณ กฤษณะรังสรรค์	ศูนย์โรคหัวใจ โรงพยาบาลตำรวจ
นพ.ประสาธน์ เหล่าถาวร	ศูนย์โรคหัวใจ โรงพยาบาลพระมงกุฎเกล้า
นพ.บุญจง แซ่จึ้ง	ศูนย์โรคหัวใจ โรงพยาบาลโรคทรวงอก
นพ.อภิชาติ สุนทรธรรม	ศูนย์โรคหัวใจ มหาวิทยาลัยเชียงใหม่

**ที่ปรึกษา** ศ.นพ.จิตร สิทธิอมร, ศ.นพ.ศุภชัย ไชยธีพันธ์, ศ.นพ.ธาดา ยิบอินซอย

### หลักการและเหตุผล

โครงการวิจัยนี้เกิดจากความร่วมมือของนักวิจัยของ 47 ประเทศทั่วโลก โดยมีวัตถุประสงค์เพื่อหาความสัมพันธ์ของการเกิดโรคหัวใจขาดเลือดเฉียบพลัน (Acute Myocardial Infarction) กับปัจจัยเสี่ยงที่ทราบกันอยู่แล้ว และที่กำลังเป็นที่สนใจใหม่ ที่จะมีส่วนต่อการพัฒนาความรู้และกระบวนการที่จะนำไปสู่การควบคุมป้องกันโรคหัวใจอย่างมีประสิทธิภาพต่อไป

สำหรับประเทศไทยได้เข้าร่วมโครงการในฐานะเป็นหน่วยเก็บข้อมูล ซึ่งมีทั้งหมด 12 หน่วยได้แก่ โรงพยาบาลศิริราช โรงพยาบาลจุฬาลงกรณ์ โรงพยาบาลรามาริบัติ มหาวิทยาลัยขอนแก่น โรงพยาบาลราชวิถี โรงพยาบาลภูมิพลอดุลยเดช วชิรพยาบาล โรงพยาบาลตำรวจ โรงพยาบาลพระมงกุฎเกล้า สถาบันโรคทรวงอก

### วัตถุประสงค์

เพื่อหาขนาดและความสัมพันธ์ระหว่างปัจจัยเสี่ยงที่ทราบกันแล้ว และปัจจัยเสี่ยงที่เพิ่งค้นพบใหม่ของการเกิดโรคกล้ามเนื้อหัวใจขาดเลือด และความแตกต่างของความเสี่ยงของปัจจัยเสี่ยงภายในประชากรกลุ่มต่างๆ

## วิธีการวิจัย

กลุ่มตัวอย่างเป็นผู้ป่วย nonfatal MI ทั้งชายและหญิง ที่ได้รับการรักษาในหน่วย Coronary Care Unit หรือ Cardiology Ward ภายใน 24 ชม.แรกของการเกิดของ MI โดย กำหนด Criteria ของการวินิจฉัย ประกอบด้วยอาการและการเปลี่ยนแปลงทางคลื่นไฟฟ้าหัวใจซึ่งประกอบด้วย New Pathologic Q wave 1mm ST elevation lead ที่ติดต่อกันเท่ากับหรือมากกว่า 2 leads New left bundle branch block new persistent ST-T wave change ซึ่งเป็นลักษณะของ non-Q wave MI โดยที่การวินิจฉัยจะได้รับการยืนยันด้วยการตรวจ Cardiac enzyme (สูงมากกว่า 2 เท่าของค่าปกติ) หรือมีการเปลี่ยนแปลงทางคลื่นไฟฟ้าหัวใจที่เข้ากับการวินิจฉัยของ MI (Evolution of ECG change)

สำหรับกลุ่มควบคุมเป็น Hospital control (Match for +/- 5 years and sex) โดยการศึกษาจากผู้ป่วยจำนวน 500 ราย เปรียบเทียบกับ Control 500 ราย กลุ่มตัวอย่างจะได้รับการซักประวัติ ตรวจร่างกาย วัดความดันโลหิต การสอบถามเกี่ยวกับวิถีชีวิต การออกกำลังกาย การรับประทานอาหาร การรับประทานยา และเจาะเลือด เพื่อส่งไปวิเคราะห์ในห้องปฏิบัติการกลางที่ Hamilton Sciences Corporation, Ontario, Canada

## ผลการวิเคราะห์กลุ่มตัวอย่างในประเทศไทย

ในกลุ่มตัวอย่างที่เก็บได้ในประเทศไทย มีผู้ป่วยโรคกล้ามเนื้อหัวใจตายเฉียบพลันจำนวน 181 คน และคนปกติที่ไม่เป็นโรคหัวใจจำนวน 216 คน ส่วนใหญ่ได้รับการศึกษาระหว่าง 1-8 ปี ปัจจัยที่ได้นำมาวิเคราะห์เพื่อหาความเสี่ยงในการเกิดโรคหัวใจขาดเลือด ได้แก่ การเป็นโรคความดันโลหิตสูง โรคเบาหวาน ค่าดัชนีมวลกาย สัดส่วนระหว่างเอวและสะโพก การออกกำลังกาย การดื่มเครื่องดื่มที่มี Alcohol ผสม พบว่า การเป็นโรคความดันโลหิตสูง โรคเบาหวาน และการมีสัดส่วนระหว่างเอวกับสะโพกมากกว่า 0.97 ในเพศชาย และมากกว่า 0.98 ในเพศหญิง เป็นปัจจัยเสี่ยงก่อให้เกิดโรคหัวใจขาดเลือดเฉียบพลัน ส่วนปัจจัยด้านระดับการศึกษา รายได้ ความเครียด และการออกกำลังกาย พบว่าไม่ได้เป็นปัจจัยเสี่ยงในการเกิดโรคหัวใจขาดเลือดเฉียบพลัน ส่วนการดื่มเครื่องดื่มที่มี Alcohol ปริมาณเล็กน้อย (1 ครั้ง/สัปดาห์) เป็นปัจจัยป้องกันการเกิดโรคหัวใจขาดเลือด

## การเผยแพร่ผลการวิจัย

Rosengren, A., Hawken, S., Ounpuu, S., Sliwa, K., Zubaid, M., Almahmeed, W. A., et al. (2004).

Association of psychosocial risk factors with risk of acute myocardial infarction in 11119 cases and 13648 controls from 52 countries (the INTERHEART study): case-control study. *Lancet*, 364(9438), 953-962.

Yusuf, S., Hawken, S., Ounpuu, S., Dans, T., Avezum, A., Lanas, F., et al. (2004). Effect of potentially modifiable risk factors associated with myocardial infarction in 52 countries (the INTERHEART study): case-control study. *Lancet*, 364(9438), 937-952.

## ชื่อโครงการไทย 5 การศึกษาครอบครัวโรคไหลตาย

นักวิจัยหลัก นพ.สมเกียรติ แสงวัฒนาโรจน์

ที่ปรึกษา ศ.นพ.จิตร สิทธิอมร

### หลักการและเหตุผล

โรคไหลตาย (Sudden Unexplained Death Syndrome or SUDS) เป็นปัญหาที่สำคัญทางการแพทย์ การสาธารณสุข เศรษฐกิจ และสังคม ของประเทศไทย สามเหตุการณ์เกิดไหลตายยังไม่เป็นที่ทราบแน่ โรคไหลตาย และ กลุ่มอาการ Brugada มีข้อเหมือนและแตกต่างกันหลายประการ คือ ทั้ง 2 กลุ่มอาการพบในผู้ชายมากกว่าผู้หญิง พบในคนอายุน้อย มีประวัติครอบครัว เกิด sudden death จาก VF โดยไม่พบความผิดปกติของหัวใจ (structure heart defect) ได้เหมือนกับ ECG พบลักษณะ RBBB และ ST elevation ใน lead V1 ถึง V3 ได้ทั้งคู่ ในกลุ่มอาการ Brugada จะตรวจพบในสมาชิกครอบครัวส่วนใหญ่ได้ หรือ ถ้าไม่พบ ก็สามารถกระตุ้นให้เกิด โดยการฉีด Ajmaline หรือ Procainamide ส่วนไหลตายจะตรวจพบได้ในบางรายที่รอดชีวิตมา และไม่แน่นอน บางครั้งตรวจพบ บางครั้งไม่พบ ถ้าตรวจพบ จะทำให้ ECG กลับมาปกติได้โดยการฉีด isoproterenol และ เมื่อฉีด Lidocaine ECG จะกลับผิดปกติอีก

### วัตถุประสงค์

ตรวจหา mutation ของ ion channel gene ในครอบครัวของผู้เสียชีวิตหรือนอดชีวิตจากไหลตาย ที่มี ECG แบบ RBBB และ ST elevation ใน lead V1 – 3 ด้วยวิธี DNA sequencing

To determine the mutation on ion channel gene in Thai SUDS or SUDS survivor families with RBBB and ST elevation in V1 – 3 pattern by direct DNA sequencing

### วิธีการดำเนินการวิจัย และการวิเคราะห์ข้อมูล

1. ผู้เสียชีวิตจากไหลตาย คือ ชาวอีสานที่แข็งแรงดี ไม่มีโรคประจำตัว เสียชีวิตกะทันหันโดยไม่คาดคิดขณะนอนหลับหรือ พักผ่อน และการตรวจชันสูตรศพไม่พบสาเหตุการตาย
2. ผู้รอดชีวิตจากไหลตาย คือ ชาวไทยที่เดิมแข็งแรงดี ไม่มีโรคประจำตัว เกิด cardiac arrest หรือ polymorphic ventricular tachycardia/ventricular fibrillation และได้ทำ cardiopulmonary resuscitation สำเร็จ ไม่มีประวัติการใช้ยาที่ทำให้เกิด polymorphic VT/VF ตรวจร่างกายปกติ และตรวจ ECG ไม่พบ long QT interval, blood chemistry, chest x-ray, echocardiography, exercise stress test และ coronary angiography อยู่ในเกณฑ์ปกติ
3. ครอบครัวผู้เสียชีวิตหรือ ผู้รอดชีวิตจากไหลตายที่มี ECG แบบ RBBB และ ST elevation ใน lead V1 – 3 คือ ญาติพี่น้องของผู้เสียชีวิตหรือรอดชีวิตจากไหลตาย ที่ตรวจคลื่นหัวใจ พบ RBBB และ ST segment (ส่วนของ ST segment ที่ห่างจาก J point 0.08 second) ยกสูงขึ้นมากกว่า 0.01 mV จาก baseline ใน lead V1 – 3 อย่างน้อย 2 ใน 3 leads ไม่ว่าจะเกิดขึ้นเอง หรือ ทำให้เกิด โดยใช้ยา ajmaline, procainamide เป็นต้น โดยการฉีดยาเข้าทางหลอดเลือดดำ ญาติพี่น้องที่ไม่มี ECG ดังกล่าว ถือว่าปกติ
4. คนปกติ หรือ กลุ่มเปรียบเทียบ คือ ชาวไทยอายุมากกว่า 50 ปีภูมิลำเนาเป็นคนอีสาน อาศัยอยู่ในหมู่บ้านเดียวกับผู้เสียชีวิตหรือรอดชีวิตจากไหลตาย ไม่มีประวัติการตายเฉียบพลันในครอบครัวอย่างน้อย 3 ชั่วอายุคน และไม่มี ECG แบบ RBBB และ ST elevation ใน lead V1 – 3 จำนวน 50 ราย

### สถานที่ในการทำวิจัย

โรงพยาบาล ในจังหวัดนครพนม จังหวัดสกลนคร จังหวัดอำนาจเจริญ จังหวัดขอนแก่น และโรงพยาบาลในภาคตะวันออกเฉียงเหนือที่ใกล้กับที่อยู่ของครอบครัวไหลตาย โดยคณะวิจัยจะออกไปตรวจประมาณเดือนละครั้งจำนวน 9 ครั้งๆ ละ 1-3 วัน ส่วนญาติพี่น้องของครอบครัวไหลตายที่มาทำงานในกรุงเทพฯ จะนัดมาตรวจที่สาขาวิชาโรคหัวใจและหลอดเลือด ภาควิชาอายุรศาสตร์ คณะแพทยศาสตร์ จุฬาลงกรณ์มหาวิทยาลัย

### การนำผลไปใช้ประโยชน์/การเผยแพร่ผลงาน

ผลการวิจัยได้เผยแพร่ในหน่วยงาน

**ชื่อโครงการไทย** 6 แนวโน้มการช่วยคืนชีพในผู้ป่วยระยะสุดท้าย และประสิทธิผลของการทำพินัยกรรมชีวิต เรื่องการช่วยคืนชีพในผู้ป่วยระยะสุดท้ายในโรงพยาบาลมหาวิทยาลัยเชียงใหม่

**ชื่อโครงการอังกฤษ** Trend of Cardiopulmonary resuscitation performed in Patients with Terminal Illness and the effectiveness of advance directive for terminal care in Chiang Mai University Hospital

**นักวิจัยหลัก** ผศ.ดร.สุตารัตน์ สิทธิสมบัติ

**หน่วยงาน** คณะพยาบาลศาสตร์ มหาวิทยาลัยเชียงใหม่

**หลักการและเหตุผล** The key hypothesis behind advance directives (ADs) proposes that, if an intervention enhances a person's right to choose, a dying person will not opt for expensive, life-prolonging medical care and an ethically acceptable saving of resources will result.

**วัตถุประสงค์** To assess the acceptability and effectiveness of ADs in reducing cardiopulmonary resuscitation (CPR) attempts and in-hospital death among terminally ill patients in a tertiary care hospital in northern Thailand

**ระเบียบวิธีวิจัย** Non-randomized controlled study

**ผลสรุป** The majority of the terminal ill patients and the surrogates preferred to employ ADs in expressing their preferences on CPR and there was a high level of agreement between the subjects and surrogates on the decision. The use of ADs appeared to be effective in reducing futile CPR attempts and the in-hospital mortality rate among subjects during the index hospitalization. Advance directives were accepted well in this study setting.

**การนำผลไปใช้ประโยชน์/การเผยแพร่ผลงาน**

Sittisombut, S., Love, E. J., & Sitthi-Amorn, C. (2001). Cardiopulmonary resuscitation performed in patients with terminal illness in Chiang Mai University Hospital, Thailand. *Int J Epidemiol*, 30(4), 896-898.

Sittisombut, S., Maxwell, C., Love, E. J., & Sitthi-Amorn, C. (2008). Effectiveness of advance directives for the care of terminally ill patients in Chiang Mai University Hospital, Thailand. *Nurs Health Sci*, 10(1), 37-42.

**ชื่อโครงการไทย** 7 ทศนคติต่อการทำพินัยกรรมชีวิตและการให้ข้อมูลเรื่องผลของการช่วยคืนชีพในผู้ป่วย  
อายุรกรรม โรงพยาบาลมหาวิทยาลัยเชียงใหม่

**ชื่อโครงการอังกฤษ** Attitude towards advance directives and the impact of prognostic information on  
the preference for cardiopulmonary resuscitation on medical inpatients in CMU Hospital, Thailand

**นักวิจัยหลัก** ผศ.ดร.สุตารัตน์ สิทธิสมบัติ

**หน่วยงาน** คณะพยาบาลศาสตร์ มหาวิทยาลัยเชียงใหม่

**นักวิจัยในทีม -**

**หลักการและเหตุผล** Our previous study revealed that cardiopulmonary resuscitation (CPR) was  
performed in 65.7% of 411 terminally ill patients who died in a tertiary-care university hospital in  
northern Thailand. Advance directives (ADs) are needed to ensure that life-sustaining therapies are  
used more appropriately.

**วัตถุประสงค์** To investigate inpatients' attitudes regarding ADs for CPR and the impact of providing  
prognostic information on treatment preferences for CPR

**ระเบียบวิธีวิจัย** Descriptive study

**ผลสรุป** Most patients had a positive attitude towards ADs for CPR. The majority preferred to have  
CPR when no information was provided on the chance of survival. However, this proportion  
decreased depending on the prognostic scenarios. Our investigation suggested that the preference  
of patients for CPR should be assessed individually and gradually, with adequate information given  
on the chance of survival.

**การนำผลไปใช้ประโยชน์/การเผยแพร่ผลงาน**

Sittisombut, S., Love, E. J., & Sitthi-Amorn, C. (2005). Attitudes toward advance directives and the  
impact of prognostic information on the preference for cardiopulmonary resuscitation in  
medical inpatients in Chiang Mai University Hospital, Thailand. *Nurs Health Sci*, 7(4), 243-  
250.

Sittisombut, S., Maxwell, C., Love, E. J., & Sitthi-Amorn, C. (2009). Physicians' attitudes and  
practices regarding advanced end-of-life care planning for terminally ill patients at Chiang  
Mai University Hospital, Thailand. *Nurs Health Sci*, 11(1), 23-28.

## ชื่อโครงการ ไทย 8 การตรวจสุขภาพประจำปีของผู้สูงอายุในเขตชนบทของจังหวัดขอนแก่น

## ชื่อโครงการอังกฤษ Annual Health Examination among Elderly in Rural Areas of Khon Kaen Province

นักวิจัยหลัก ผศ.ดร.จิราพร เขียวอยู่

นักวิจัยในทีม ศิริพร คำสะอาด

ที่ปรึกษา ศ.นพ.จิตร สิทธิอมร

**หลักการและเหตุผล** การวิจัยนี้มีวัตถุประสงค์เพื่อศึกษาการไปตรวจสุขภาพประจำปีของผู้สูงอายุในเขตชนบทของจังหวัดขอนแก่น ตัวอย่างเป็นผู้ที่มีอายุ 60 ปีขึ้นไปจำนวน 576 ราย สุ่มตัวอย่างแบบหลายขั้นตอน เก็บข้อมูลโดยการสัมภาษณ์โดยตรงตามแบบสอบถามซึ่งสร้างขึ้นด้วยวิธีมาตรฐาน

**รูปแบบการวิจัย** เป็นการวิจัยเชิงพรรณนา

**ผลสรุป** ที่ผ่านมามีผู้สูงอายุร้อยละ 79.9 ได้รับตรวจสุขภาพอย่างใดอย่างหนึ่งต่อไปนี้ การซักประวัติการสูบบุหรี่ การซักประวัติการดื่มสุรา การซักประวัติการบริโภคอาหาร การชั่งน้ำหนัก วัดส่วนสูง การวัดความดันโลหิต การวัดชีพจร การตรวจเลือด การตรวจปัสสาวะ การตรวจสายตา และ การตรวจฟัน เมื่อพิจารณาแต่ละประเภทของการตรวจสุขภาพดังกล่าวข้างต้น พบ ร้อยละของผู้สูงอายุที่ไปตรวจสุขภาพเท่ากับ 23.0, 21.0, 36.4, 71.7, 32.5, 75.7, 55.4, 43.0, 41.6, 18.8, และ 13.3 ตามลำดับ ในกลุ่มผู้สูงอายุสตรี 391 ราย มีผู้ไปรับการตรวจเต้านม และมะเร็งปากมดลูก ร้อยละ 9.0 และ 11.5 ตามลำดับ

**การนำผลไปใช้ประโยชน์/การเผยแพร่ผลงาน**

ได้เผยแพร่ผลงานในหน่วยงาน

**ชื่อโครงการไทย** 9 การศึกษาความเข้าใจ ความน่าเชื่อถือ ความสมบูรณ์ และความไวต่อการเปลี่ยนแปลงของแบบสอบถาม ฉบับภาษาไทย Health Assessment Questionnaire สำหรับผู้ป่วยโรคข้ออักเสบ

**ชื่อโครงการอังกฤษ** Comprehensibility, reliability, validity, and responsiveness of the Thai version of the Health Assessment Questionnaire in Thai patients with rheumatoid arthritis

**นักวิจัยหลัก** พญ.มนาริป โอศิริ

**ที่ปรึกษา** ศ.นพ.จิตร สิทธิอมร

**หลักการและเหตุผล** The Health Assessment Questionnaire Disability Index (HAQ-DI) is a commonly used instrument to assess functional status of patients with rheumatoid arthritis (RA). Translations and adaptations of the HAQ-DI have been carried out for use with RA patients in several countries. The objective is to evaluate the psychometric properties of the Thai version of the HAQ-DI (Thai HAQ) in Thai patients with RA.

**รูปแบบการวิจัย** Comprehensibility of the Thai HAQ was assessed by 126 patients with RA from 6 medical centers in Thailand. Another group of 115 patients with active RA was enrolled to test the reliability (internal reliability and 1-week test-retest reliability), construct validity (correlations with other measures of RA disease activity), floor and ceiling effects, and sensitivity to change of the Thai HAQ at 3 months of treatment with disease-modifying antirheumatic drugs.

**ผลสรุป** The Thai HAQ is comprehensible, reliable, valid and sensitive to change in the evaluation of functional status of Thai patients with RA. The Thai HAQ is an essential tool to measure treatment effects and progression of disability in RA patients and should be applied in both clinical trials and routine clinical care settings.

**การนำผลไปใช้ประโยชน์/การเผยแพร่ผลงาน**

Osiri, M., Wongchinsri, J., Ukritchon, S., Hanvivadhanakul, P., Kasitanon, N., & Siripaitoon, B. (2009). Comprehensibility, reliability, validity, and responsiveness of the Thai version of the Health Assessment Questionnaire in Thai patients with rheumatoid arthritis. *Arthritis Res Ther*, 11(4), R129.

**ชื่อโครงการไทย** 10 การศึกษาภาวะความชุก ความรุนแรง บั๊จัย และผลที่เกิดขึ้นจากการเกิดเนื้อตายในผู้ป่วยที่ถูกงูเห่า และการศึกษาการป้องกันภาวะเนื้อตายที่เกิดขึ้นในสัตว์ทดลอง

**ชื่อโครงการอังกฤษ** Prevalence, severity, determinants and consequences of tissue necrosis among victims envenomed by *Naja kaouthia* (Thai Cobra) and *Calloselasma rhodostoma* (Malayan Pit Viper) and its prevention in an experimental model

**นักวิจัยหลัก** ดร.นวลน้อย วงศ์ทองคำ

**ที่ปรึกษา** ศ.นพ.จิตร สิทธิอมร

### หลักการและเหตุผล

การถูกงูพิษกัดในประเทศไทย ยังคงเป็นปัญหาที่สำคัญทางการแพทย์โดยเฉพาะงูเห่า และงูกะปะ ซึ่งพิษสามารถก่อให้เกิดผลทั่วร่างกายและผลเฉพาะที่ ในผู้ป่วยที่มีอาการรุนแรงอาจเสียชีวิตหรืออาจเสียอวัยวะที่ถูกกัดอย่างถาวร หรืออาจต้องตัดอวัยวะส่วนนั้นทิ้งและก่อให้เกิดความพิการพหุระบบบริเวณที่ถูกกัด เกิดจากการทำงานร่วมกันของเอนไซม์ phospholipase A<sub>2</sub> (PLA<sub>2</sub>) ซึ่งสามารถทำลายกล้ามเนื้อ เอนไซม์ metalloproteinase ซึ่งทำให้เกิดภาวะเลือดออก และเอนไซม์ hyaluronidase ซึ่งช่วยทำให้พิษงูแพร่กระจาย รวมทั้งสารพิษอื่นๆ ในพิษงู

### วัตถุประสงค์

1. การศึกษาด้านระบาดวิทยา ทั้งส่วนการศึกษาไปข้างหน้า และการศึกษาย้อนหลัง ในผู้ป่วยที่ถูกงูเห่าหรืองูกะปะกัด
2. การทดลองเพื่อดูประสิทธิภาพของตัวยับยั้งเอนไซม์ metalloproteinase และ PLA<sub>2</sub> โดยมีจุดมุ่งหมายเพื่อหาวิธีลดภาวะเกิดเนื้อตายบริเวณที่ถูกกัด และลดการเกิดพิษทั่วร่างกาย

### ผลสรุป

ผลการศึกษาทางด้านระบาดวิทยาพบว่า อุบัติการณ์ของการเกิดเนื้อตายบริเวณที่ถูกงูกะปะและงูเห่ากัดสูงถึง 94.50% และ 91.10% ตามลำดับ โดยส่วนใหญ่จะพบภาวะเนื้อตายที่ไม่รุนแรงและจะแสดงออกในช่วง 12 ชั่วโมงแรกของการเข้ารับการรักษา การรักษาผู้ป่วยส่วนใหญ่เป็นการทำแผลเพื่อป้องกันและลดการติดเชื้อ มีผู้ป่วยที่ถูกงูเห่ากัดเพียง 1 รายที่ต้องถูกตัดนิ้ว

ผลการศึกษาในการทดลอง พบว่า 2 mM Na<sub>2</sub>EDTA หรือ 20 mM N-Phenylglycine สามารถยับยั้งผลของเอนไซม์ metalloproteinase และ PLA<sub>2</sub> ตามลำดับได้อย่างสมบูรณ์ ทั้งในพิษงูเห่าและงูกะปะ การทดลองในสัตว์ทดลองพบว่า Na<sub>2</sub>EDTA (93.05 ไมโครกรัม/หนู) หรือ N-Phenylglycine (37.80 ไมโครกรัม/หนู) เมื่อทำการ preincubate กับงูก่อนที่จะฉีดในหนูพบว่าสามารถลดการเกิดภาวะการฉีกขาด และการเกิดเนื้อตายอันเกิดจากพิษงูได้อย่างมีนัยสำคัญทางสถิติ ส่วน "Inhibitor mixture" ที่ประกอบด้วย N-Phenylglycine (37.80 ไมโครกรัม/หนู) Na<sub>2</sub>EDTA (93.05 ไมโครกรัม/หนู) และตัวยับยั้งเอนไซม์ hyaluronidase คือ sodium aurothiomalate (195 ไมโครกรัม/หนู) พบว่าสารผสมนี้สามารถลดการเกิดเนื้อ

ตายเฉพาะที่เมื่อฉีดภายในเวลา 1, 3 และ 10 นาที หลังจากฉีดพิษงู และยังพบว่าผลของตัวยับยั้งเอนไซม์ metalloproteinase และ PLA<sub>2</sub> สามารถยืดระยะเวลาตายของหนูที่ฉีดพิษงูเห่าและงูกะปะ ในขนาดที่ทำให้หนูตาย ผลการทดลองนี้บ่งชี้ว่าสารผสม Inhibitor mixture มีประสิทธิภาพดีในการลดภาวะการเกิดเนื้อตาย ถ้าฉีดทันทีที่บริเวณที่ถูกกัด และลดความเป็นพิษที่ออกฤทธิ์ทั่วร่างกายของพิษงูเห่าและงูกะปะ

### การนำผลไปใช้ประโยชน์

ผลการศึกษานี้ทำให้ได้ข้อมูลที่เป็นประโยชน์ในการรักษาผู้ป่วยที่มีแผลถูกกัด โดยผลการทดลองบ่งชี้ว่าสารผสม Inhibitor mixture มีประสิทธิภาพดีในการลดภาวะการเกิดเนื้อตาย ถ้าฉีดทันทีที่บริเวณที่ถูกกัด และลดความเป็นพิษที่ออกฤทธิ์ทั่วร่างกายของพิษงูเห่าและงูกะปะ

### การนำผลไปใช้ประโยชน์/การเผยแพร่ผลงาน

- Wongtongkam, N., Wilde, H., Sitthi-Amorn, C., & Ratanabanangkoon, K. (2005a). A study of 225 Malayan pit viper bites in Thailand. *Mil Med*, 170(4), 342-348.
- Wongtongkam, N., Wilde, H., Sitthi-Amorn, C., & Ratanabanangkoon, K. (2005b). A study of Thai cobra (*Naja kaouthia*) bites in Thailand. *Mil Med*, 170(4), 336-341.

## โครงการสหสาขาวิชาเพื่อแก้ปัญหาสาธารณสุขที่สำคัญของประเทศ 7 โครงการ

**ชื่อโครงการไทย 1** ผลของการปรับเปลี่ยนการบริโภคมาเป็นข้าวกล้อง งาม ผัก ผลไม้ ตามฤดูกาลของประชาชนในเขตเทศบาลในเขตเทศบาลในจังหวัดสุโขทัย

**นักวิจัยหลัก** สมนึก สุขัยธนาวิช

**ที่ปรึกษา** ศ.นพ.จิตร สิทธิอมร

### หลักการและเหตุผล

ปัจจุบันเทคโนโลยีเจริญก้าวหน้าอย่างรวดเร็ว โดยเฉพาะอย่างยิ่งการแปรรูปผลิตภัณฑ์ด้านอาหารและการกระตุ้นความอยากด้วยอิทธิพลสื่อโฆษณา ทำให้แบบแผนการบริโภคของคนไทยเปลี่ยนแปลงไปจากอดีต สถานการณ์การเจ็บป่วยของประชาชนที่เปลี่ยนแปลงจากโรคติดต่อมาเป็นโรคที่ไม่ติดต่อ การเจ็บป่วยด้วยโรคเหล่านี้ถูกเชื่อมโยงให้ประชาชนหันมาให้ความสนใจต่อการบริโภคมากขึ้น พฤติกรรมการบริโภคอาหารที่ถูกปนเปื้อนสารเคมีต่างๆ โดยที่ผู้บริโภคไม่มีโอกาสเลือกมากนักถูกมองว่าเป็นสาเหตุหนึ่งที่ทำให้เกิดโรคที่รักษาไม่หายเหล่านี้ หลายองค์กรให้ความสนใจในการบริโภคแบบยั่งยืน สุขภาพแบบองค์รวม เช่น อาหารมังสวิรัต แมคโครไบโอติก หรืออาหารชีวจิต การปรับเปลี่ยนแบบแผนการบริโภคของคนไทยเพื่อเอื้อต่อสุขภาพดี มีชีวิตที่ยั่งยืนโดยไม่เคร่งครัดกับหลักการบริโภคแบบบังคับมากนัก น่าจะเป็นทางเลือกที่เหมาะสมให้สอดคล้องกับสภาวะเศรษฐกิจของประชาชนและของประเทศ

### วัตถุประสงค์

1. เพื่อประเมินรูปแบบการรณรงค์การบริโภคข้าวกล้อง งาม ผัก ผลไม้ตามฤดูกาลสามารถทำให้ประชาชนเปลี่ยนมาบริโภคอาหารดังกล่าวมากกว่าพื้นที่ที่ไม่ได้รณรงค์
2. เพื่อศึกษาประสิทธิผลของการบริโภคข้าวกล้อง งาม ผัก ผลไม้ตามฤดูกาล ที่มีผลต่อสุขภาพของประชาชน
3. เพื่อเปรียบเทียบค่า fasting blood sugar อัตราการไ้ยา และค่าใช้จ่ายด้านสุขภาพของผู้ป่วยเบาหวานประเภท NIDDM ที่ปรับเปลี่ยนมาบริโภคข้าวกล้อง งาม ผัก ผลไม้ตามฤดูกาล

ขอบเขตการวิจัย พื้นที่ทดลอง คือ เขตเทศบาลเมืองสวรรคโลก พื้นที่ควบคุม คือ เขตเทศบาลเมืองสุโขทัย

### การเก็บรวบรวมข้อมูล

- ประเมินรูปแบบการรณรงค์และประสิทธิผลของการปรับเปลี่ยนการบริโภค
- พัฒนารูปแบบการรณรงค์เพื่อให้ความรู้ในพื้นที่

### ประโยชน์ที่คาดว่าจะได้รับ

1. เพื่อประเมินรูปแบบการรณรงค์การบริโภคข้าวกล้อง งาม ผัก ผลไม้ตามฤดูกาลสามารถทำให้ประชาชนเปลี่ยนมาบริโภคอาหารดังกล่าวมากกว่าพื้นที่ที่ไม่ได้รณรงค์
2. เพื่อศึกษาประสิทธิผลของการบริโภคข้าวกล้อง งาม ผัก ผลไม้ตามฤดูกาล ที่มีผลต่อสุขภาพของประชาชน

3. เพื่อเปรียบเทียบค่า fasting blood sugar อัตราการใช้ยา และค่าใช้จ่ายด้านสุขภาพของผู้ป่วยเบาหวานประเภท NIDDM ที่ปรับเปลี่ยนมาบริโภคข้าวกล้อง งามัก ผลไม้ตามฤดูกาล

#### การนำผลไปใช้ประโยชน์/การเผยแพร่ผลงาน

เผยแพร่ผลการศึกษาในหน่วยงาน

**ชื่อโครงการไทย 2** ความต้องการและปัญหาด้านสุขภาพ การเข้าถึงบริการ และคุณภาพชีวิตของวัยรุ่น กับ การสร้างภาคีความร่วมมือ

**ชื่อโครงการอังกฤษ** Adolescent Health Needs, Accessibility of Services and Quality of Life by Assisting in the Development of Community Partnerships.

**นักวิจัยหลัก** ดร.รัตนา สำโรงทอง

**ที่ปรึกษา** ศ.นพ.จิตร สิทธิอมร

### หลักการและเหตุผล

วัยรุ่นคือวัยแรกเจริญพันธุ์จนถึงวัยเริ่มต้นของการเติบโตเป็นผู้ใหญ่ เป็นวัยที่ต้องการการดูแลเป็นพิเศษต่างไปจากวัยอื่นๆ เนื่องจากธรรมชาติของวัยรุ่นเป็นสิ่งที่ซับซ้อนยากที่จะเข้าใจการค้นหาสาเหตุของปัญหาและดำเนินการแก้ไขจึงเป็นเรื่องละเอียดอ่อน การดูแลจำเป็นจะต้องครอบคลุมไปทุกๆ ด้าน ซึ่งวัตถุประสงค์ของการดูแลนั้น ไม่ได้เฉพาะเจาะจงในเรื่องสุขภาพอย่างเดียว แต่ยังรวมถึงการพัฒนาการทางด้านกาย จิต สังคมของแต่ละบุคคลด้วย ดังนั้น นอกเหนือจากตัววัยรุ่นเองแล้ว การดูแลและแสวงหาทางแก้ไข จึงจำเป็นต้องรวมถึงครอบครัว ชุมชน สังคมและสิ่งแวดล้อมของวัยรุ่นด้วย

**วัตถุประสงค์** ศึกษาลักษณะของการให้บริการด้านสุขภาพวัยรุ่นของหน่วยงานต่างๆ ในชุมชน

1. ศึกษาถึงความต้องการและปัญหาด้านสุขภาพ การเข้าถึงบริการของวัยรุ่นชาย-หญิง และในแต่ละกลุ่มอายุ
2. เปรียบเทียบความต้องการบริการด้านสุขภาพกับการไปใช้บริการของวัยรุ่นชาย-หญิง และในแต่ละกลุ่มอายุ
3. ศึกษาถึงการให้ความหมาย “คุณภาพชีวิต” ของวัยรุ่น
4. ประเมินคุณภาพชีวิตโดยใช้ แบบวัด WHO-BREF
5. สร้างภาคีความร่วมมือเพื่อพัฒนาการให้บริการด้านสุขภาพที่วัยรุ่นสามารถเข้าถึง อันนำไปสู่การลดปัญหาทางสุขภาพ ความซึมเศร้า อันนำไปสู่การมีคุณภาพชีวิตที่ดี

การเก็บข้อมูลเชิงคุณภาพใช้แบบแนวสัมภาษณ์เจาะลึก และแนวสนทนากลุ่ม การเก็บข้อมูลเชิงปริมาณใช้แบบสอบถามกับ กลุ่มตัวอย่างเป็นวัยรุ่นอายุ 12-22 ปี ที่อาศัยอยู่ในสลัมแห่งหนึ่งในกรุงเทพฯ การวิเคราะห์สถิติใช้ Chi-square และ Regression

### ผลการศึกษา

พบว่ากลุ่มตัวอย่างเป็นเพศหญิงมากกว่าชาย (57.2% , 42.8%) แบ่งเป็นวัยรุ่นตอนต้น (อายุ 12-13 ปี 23.3%) วัยรุ่นตอนกลาง (14-17 ปี 48.2%) และวัยรุ่นตอนปลาย (18-22 ปี 28.5%) ส่วนใหญ่ (78.3%) อยู่ในสถานศึกษา มีเพียง 5.9% เจ็บป่วยเรื้อรัง วัยรุ่นเกือบ 1 ใน 5 คน เคยมีประสบการณ์ทางเพศ และส่วนใหญ่มิเพศสัมพันธ์กับคูรักร โดยมีเพียงร้อยละ 12.6 ใช้ถุงยางอนามัยทุกครั้ง วัยรุ่นให้ความหมายเรื่อง “คุณภาพชีวิต” ใกล้เคียงกับคำจำกัดความขององค์การอนามัยโลก หากแต่วัยรุ่นไม่ได้กล่าวถึงเรื่องเพศและ

การเข้าถึงบริการ วัยรุ่นส่วนใหญ่ (71.8%) มีคุณภาพชีวิตในระดับกลาง สิบอันดับแรกของปัญหาสุขภาพวัยรุ่น ได้แก่ 1) สิว 2) ตั้งครรภ์ไม่พึงประสงค์ 3) ยาบ้า 4) เฮโรอีน 5) ทำแท้ง 6) เครียด 7) ต่อสู้/ชกตี 8) บุหรี่ 9) สุรา และ 10) ข่มขืน ซึ่งพบความแตกต่างอย่างมีนัยสำคัญทางสถิติในระหว่างเพศ กับปัญหาสุขภาพที่กล่าวมาแล้ว 8 จาก 10 เรื่อง (ยกเว้นเรื่องบุหรี่กับแอลกอฮอล์) และพบความแตกต่างอย่างมีนัยสำคัญทางสถิติในระหว่างกลุ่มอายุกับปัญหาสุขภาพบุหรี่

การศึกษาเชิงคุณภาพพบว่ายังมีช่องว่าง การแบ่งแยก และความซับซ้อนของการบริการด้านสุขภาพ ปัจจัยการเข้าถึงบริการด้านสุขภาพของวัยรุ่น ได้แก่ “อาชีพของผู้ปกครอง” และ “โอกาสได้พบแพทย์ที่สถานบริการสุขภาพ” การวิเคราะห์ความแตกต่างในระหว่างเพศและกลุ่มอายุของวัยรุ่น สำหรับปัจจัยเหล่านี้ พบว่า (1) ความแตกต่างในระหว่างเพศ อาชีพของผู้ปกครองมีความสำคัญสำหรับเพศชาย ในขณะที่โอกาสการได้พบแพทย์ที่สถานบริการสุขภาพกลับมีความสำคัญสำหรับเพศหญิง (2) ความแตกต่างในระหว่างกลุ่มอายุ พบว่าปัจจัยที่มีความสำคัญที่สุดถึงการเข้าถึงบริการในวัยรุ่นตอนต้น ได้แก่ “ช่วงเวลาที่จะสะดวกในการไปสถานบริการสุขภาพ” “ระดับการศึกษาในปัจจุบัน” และ “การรู้จักสถานบริการสุขภาพในชุมชน” สำหรับวัยรุ่นตอนกลาง ได้แก่ “รู้เกี่ยวกับค่าใช้จ่ายในบ้าน” และสำหรับวัยรุ่นตอนปลาย ได้แก่ “อาชีพของผู้ปกครอง” “รู้จักสถานบริการสุขภาพในชุมชน” และ “อายุ”

การไปใช้บริการด้านสุขภาพ พบว่าในสิบอันดับแรกของปัญหาวัยรุ่นนั้น กลุ่มวัยรุ่นไปใช้บริการน้อยกว่าร้อยละ 60 เพศชายไปใช้บริการน้อยกว่าเพศหญิง โดยเพศชายไปใช้บริการเนื่องจากบาดเจ็บจากการชกตี/ต่อสู้ การสูบบุหรี่ การดื่มแอลกอฮอล์ และความเครียด สำหรับเพศหญิงไปใช้บริการเนื่องจากความเครียดและภาวะความซึมเศร้า

จากการศึกษาพบว่า ผู้ที่ทำงานเรื่องวัยรุ่นในชุมชนตกลงที่จะร่วมกันเป็นภาคีเพื่อให้วัยรุ่นเข้าถึงบริการสุขภาพ และพัฒนาคุณภาพชีวิตวัยรุ่น โดยร่วมกันวางวัตถุประสงค์หลัก และนำผลการศึกษาของงานวิจัยครั้งนี้จะเป็นข้อมูลสำคัญ ในการวางแผนการดำเนินสำหรับจัดทำกิจกรรม/โครงการในชุมชน ในการพัฒนาการเข้าถึงสุขภาพของวัยรุ่น โดยเน้นด้านระบบสืบพันธุ์/เพศสัมพันธ์ สุขภาพจิต และสารเสพติด สำหรับการวิจัยในอนาคตนั้นควรพิจารณาในเรื่องของการเข้าถึงบริการด้านสังคมและปัจจัยทางด้านสังคมอื่นๆ ภาควิชาความร่วมมือและการวิจัยเชิงปฏิบัติการจะนำมาสู่ความร่วมมือและใช้ทรัพยากรร่วมกัน โดยความร่วมมือในการเลือกผู้นำเพื่อเป็นกลไกที่สำคัญในการทำงานร่วมกัน

### การนำผลไปใช้ประโยชน์/การเผยแพร่ผลงาน

Somrongthong, R., Panuwatsuk, P., Amarathithada, D., Chaipayom, O., & Sitthi-amorn, C. (2003). Sexual behaviors and opinions on sexuality of adolescents in a slum community in Bangkok. *Southeast Asian J Trop Med Public Health*, 34(2), 443-446.

หมายเหตุ ดร.รัตนา สำโรงทอง ได้รับตำแหน่งเป็นอาจารย์ประจำวิทยาลัยวิทยาศาสตร์สาธารณสุข จุฬาลงกรณ์มหาวิทยาลัย

## ชื่อโครงการ ไทย 3 การศึกษาผลกระทบของฝุ่นละอองขนาดเล็กต่อสุขภาพประชาชน จังหวัดเชียงใหม่

### นักวิจัยหลัก ดร.นันทวรรณ วิจิตรวาทการ

ที่ปรึกษา ศ.นพ.จิตร สิทธิอมร

### หลักการและเหตุผล

ประเทศไทยเป็นประเทศกำลังพัฒนา มีการนำวิทยาการและเทคโนโลยีใหม่ ๆ เข้ามาใช้อย่าง มากมาย ส่งผลให้เกิดการเจริญเติบโตทางเศรษฐกิจที่รวดเร็ว โดยเฉพาะในช่วงปี พ.ศ. 2538 – 2539 ทำให้เกิดกิจกรรมการก่อสร้างอาคาร ถนน สาธารณูปโภค รวมทั้งปริมาณยานพาหนะที่เพิ่มมากขึ้น และส่งผลกระทบต่อสิ่งแวดล้อมคือ ปริมาณฝุ่นละอองที่สูงขึ้นจนอยู่ในระดับที่เป็นปัญหา และมีผลกระทบต่อสุขภาพ ซึ่งปัญหานี้ไม่เพียงแต่เฉพาะในเขตกรุงเทพมหานครเท่านั้น ในพื้นที่ต่างจังหวัดก็เกิดผลกระทบเช่นกัน ในภาคเหนือที่จังหวัดลำปาง เชียงใหม่ นครสวรรค์ พบว่าสารมลพิษที่มีปริมาณเกินมาตรฐาน ได้แก่ ฝุ่นขนาดเล็กกว่า 10 ไมครอน ( $PM_{10}$ ) ก๊าซโอโซน ( $O_3$ ) และก๊าซซัลเฟอร์ไดออกไซด์ ( $SO_2$ ) ส่วนในภาคตะวันออกเฉียงเหนือพบว่ามีปัญหาจากฝุ่นละอองเพียงชนิดเดียว จากการตรวจวัดที่จังหวัดขอนแก่นและนครราชสีมา ในภาคกลาง ภาคตะวันออกและภาคใต้ พบว่าฝุ่นละอองเป็นปัญหาหลักและก๊าซโอโซนเป็นปัญหารองลงมา

ฝุ่นละอองเป็นปัญหาอยู่ในทุกภาคของประเทศไทย แต่ในการศึกษาผลกระทบของฝุ่นละอองต่อสุขภาพ ส่วนใหญ่เป็นการศึกษาในกรุงเทพมหานคร เพื่อเป็นการขยายผลการศึกษาให้มีภาพกว้างขึ้น เห็นผลกระทบต่อสุขภาพที่ชัดเจนขึ้นจึงน่าจะมีการศึกษาในจังหวัดอื่น ๆ ที่นอกเหนือจากกรุงเทพมหานคร จากการพิจารณาถึงปริมาณฝุ่นละอองขนาดเล็ก ( $PM_{10}$ ) และลักษณะของชุมชนที่ศึกษา พบว่าในจังหวัดเชียงใหม่มีลักษณะของชุมชนใกล้เคียงกับพื้นที่วงเวียนโอเดียในกรุงเทพมหานคร ซึ่งทำการศึกษาในปี พ.ศ.2538 – 2539 ในแง่ของผลกระทบของฝุ่นละอองต่อสุขภาพทางระบบหายใจ ถึงแม้ว่าปริมาณ  $PM_{10}$  ที่ตรวจวัดได้ในจังหวัดเชียงใหม่จะมีค่าไม่สูงมากนัก คือมีค่าเฉลี่ยต่อปีเท่ากับ 75.3 และ 85.2 ไมโครกรัมต่อลูกบาศก์เมตร ( $\mu g/m^3$ ) ในปี พ.ศ.2539 และ พ.ศ.2540 ตามลำดับ แต่จากการตรวจวัดปริมาณ  $PM_{2.5}$  โดยใช้เครื่องมือของ US-EPA ที่จังหวัดเชียงใหม่ในช่วงฤดูแล้ง พบว่ามีค่าเฉลี่ย 24 ชั่วโมง อยู่ในช่วง 200–300  $\mu g/m^3$  ซึ่งเกินค่ามาตรฐานของสหรัฐอเมริกาที่ตั้งไว้ที่ 65  $\mu g/m^3$  และจากผลการศึกษาในต่างประเทศทั่วโลก พบว่าปริมาณฝุ่นละอองขนาดเล็ก ( $PM_{10}$  และ  $PM_{2.5}$ ) มีความสัมพันธ์กับอัตราป่วยและอัตราตายเนื่องจากโรคระบบหายใจและโรคระบบหัวใจ แต่เนื่องจากสภาพภูมิประเทศและภูมิอากาศที่แตกต่างกัน อาจส่งผลถึงองค์ประกอบของฝุ่นละอองที่แตกต่างกัน และก่อให้เกิดผลกระทบต่อสุขภาพในระดับที่แตกต่างกันด้วย

ดังนั้นจากการที่ตรวจพบว่าปริมาณ  $PM_{2.5}$  ในจังหวัดเชียงใหม่ มีค่าที่สูงเกินกว่ามาตรฐานมากจึงเป็นจุดที่น่าสนใจว่าในปัญหาฝุ่นละอองก่อให้เกิดผลกระทบต่อสุขภาพของคนในภาคเหนืออย่างไร และแตกต่างจากที่พบในกรุงเทพมหานครหรือในประเทศต่าง ๆ หรือไม่ และก่อให้เกิดความสูญเสียทางเศรษฐกิจมากน้อยเพียงใด

## วัตถุประสงค์ของโครงการ

วัตถุประสงค์หลัก เพื่อศึกษาผลของฝุ่นละอองขนาดเล็ก ( $PM_{10}$  และ  $PM_{2.5}$ ) ที่มีผลต่อสุขภาพของประชาชนในภาคเหนือของไทย

## วัตถุประสงค์เฉพาะ

1. เพื่อศึกษาความสัมพันธ์ระหว่างปริมาณฝุ่นละอองกับการเจ็บป่วยด้วยอาการระบบหายใจในเด็กนักเรียนในจังหวัดเชียงใหม่
2. เพื่อศึกษาความสัมพันธ์ระหว่างปริมาณฝุ่นละอองกับการเจ็บป่วยด้วยอาการระบบหายใจในกลุ่มผู้ใหญ่ ในจังหวัดเชียงใหม่
3. เพื่อศึกษาความสัมพันธ์ระหว่างปริมาณฝุ่นละอองกับสมรรถภาพปอด
4. เพื่อประเมินถึงความเสี่ยงทางเศรษฐกิจอันเกิดจากโรคระบบหายใจเนื่องจากฝุ่นละออง

**รูปแบบการศึกษา** การศึกษาครั้งนี้ได้วางรูปแบบการศึกษา โดยแบ่งเป็น 2 การศึกษาย่อย คือ

การศึกษาแบบ panel study เพื่อศึกษาผลกระทบต่ออัตราป่วยของระบบหายใจในแต่ละระดับของการสัมผัสกับอนุภาคมลสาร ซึ่งครอบคลุมถึง  $PM_{10}$ , และ  $PM_{2.5}$  โดยการวิเคราะห์ข้อมูลที่ได้จากการติดตามการเจ็บป่วยของกลุ่มตัวอย่างเด็กและผู้ใหญ่อายุนานกลุ่มละประมาณ 100 คน เป็นเวลา 100 วัน และการสัมผัสอนุภาคมลสารของกลุ่มตัวอย่างเหล่านี้

การประเมินผลกระทบทางเศรษฐศาสตร์ โดยการออกแบบสำรวจเพื่อรวบรวม ข้อมูลเชิงเศรษฐศาสตร์ในกลุ่มตัวอย่างเดียวกับการศึกษาแบบ panel study เพื่อประเมินผลกระทบทางเศรษฐศาสตร์ที่เกิดขึ้นเนื่องจากปริมาณอนุภาคมลสารในจังหวัดเชียงใหม่

## ผลการศึกษา Panel study ของอาการป่วยทางระบบหายใจ

การศึกษา Panel Study เป็นการติดตามอาการป่วยทางระบบหายใจของกลุ่มตัวอย่างแต่ละคนที่ได้บันทึกอาการทางระบบหายใจในช่วง 24 ชั่วโมงที่ผ่านมาของตนในแต่ละวันตามแบบบันทึกอาการที่ใช้ในการศึกษา ทำการบันทึกทุกวันติดต่อกันเป็นระยะเวลา ประมาณ 100 วัน ในขณะที่เดียวกันก็มีการเก็บข้อมูลอนุภาคมลสารในพื้นที่ใกล้เคียงกับที่กลุ่มตัวอย่างอาศัยอยู่ กลุ่มตัวอย่างของการศึกษานี้ประกอบด้วย 2 กลุ่ม คือ 1) กลุ่มผู้ใหญ่ ที่มีบ้านพักอาศัยอยู่ริมถนน 2) กลุ่มเด็กนักเรียนชั้นประถมปีที่ 3-6 จากการเก็บข้อมูลได้กลุ่มตัวอย่างผู้ใหญ่ 108 คน อายุเฉลี่ย 46 ปี ได้กลุ่มตัวอย่างเด็ก 101 คน อายุเฉลี่ย 11 ปี ทำการเก็บข้อมูลเป็นระยะเวลา 105 วัน ในแต่ละวันเฉลี่ยกลุ่มตัวอย่างให้ความร่วมมือในการเก็บข้อมูล (Response rate ) ร้อยละ 92 ในกลุ่มเด็ก และร้อยละ 95 ในกลุ่มผู้ใหญ่

ในการวิเคราะห์ข้อมูลได้แบ่งกลุ่มตัวอย่างทั้งเด็กและผู้ใหญ่เป็น 2 กลุ่ม คือ กลุ่มปกติ และกลุ่มที่มีภาวะไวต่อมลภาวะ โดยในกลุ่มเด็กคือกลุ่มที่เคยมีประวัติเกี่ยวกับโรคปอดหรือโรคหัวใจ ส่วนในกลุ่มผู้ใหญ่คือผู้ที่เคยมีประวัติโรคหอบหืด โรคถุงลมโป่งพอง หรือโรคหัวใจ หรืออายุมากกว่า 65 ปี และทำการวิเคราะห์

เปรียบเทียบอัตราเสี่ยงต่อการเกิดอาการทางระบบหายใจระหว่างกลุ่มปกติและกลุ่มที่มีความไวต่อมลภาวะ และวิเคราะห์ความสัมพันธ์กับปริมาณมลพิษที่ตรวจวัดได้ในช่วงที่ทำการศึกษา

ระดับของมลพิษอากาศในช่วงที่ศึกษา ระดับฝุ่นละอองขนาดเล็ก ( $PM_{10}$ ) ณ จุดตรวจวัดที่โรงเรียนเทศบาลวัดพวงช้าง เฉลี่ย  $146.71 \mu g/m^3$   $PM_{2.5}$  เฉลี่ย  $105.47 \mu g/m^3$  และระดับฝุ่น  $PM_{10}$  ณ จุดตรวจวัดที่เทศบาลเชียงใหม่ เฉลี่ย  $128.32 \mu g/m^3$   $PM_{2.5}$  เฉลี่ย  $90.21 \mu g/m^3$

ผลการวิเคราะห์ในกลุ่มเด็กนักเรียน พบว่า กลุ่มเด็กที่มีความไวต่อมลภาวะมีความเสี่ยงต่อการสัมผัส  $PM_{2.5}$  มากกว่าเด็กปกติอย่างมีนัยสำคัญทางสถิติ เช่นอาการหายใจไม่สะดวก แน่นหน้าอก (OR = 4.6, 95%CI 1.92, 10.93) หายใจไม่อิ่ม (OR = 4.5 95%CI 1.43, 5.65) หายใจมีเสียงวี๊ด (OR = 2.3 95% CI 0.98, 5.53 ) และเด็กผู้หญิงมีความเสี่ยงสูงกว่าเด็กผู้ชาย แต่ไม่พบความสัมพันธ์ระหว่างปริมาณ  $PM_{2.5}$  และ  $PM_{10}$  ต่อค่าความจุปอด (PEFR) แต่พบว่า ค่าทัศนวิสัย (Visibility) มีความสัมพันธ์กับค่าความจุปอด อย่างมีนัยสำคัญทางสถิติ คือเมื่อค่า Visibility เพิ่มขึ้น 10 หน่วย จะทำให้ค่า PEFR ลดลง 2.8 มิลลิลิตร

ในกลุ่มผู้ใหญ่ พบว่ากลุ่มที่มีความไวต่อมลภาวะมีความเสี่ยงต่อการสัมผัสมลพิษ ทั้ง  $PM_{10}$  และ  $PM_{2.5}$  มากกว่ากลุ่มผู้ใหญ่ปกติอย่างมีนัยสำคัญทางสถิติ เช่นอาการหายใจไม่อิ่ม (OR = 23 ) อาการหายใจไม่สะดวก แน่นหน้าอก ( OR = 13.04-12.28) และอาการระคายคอ (OR = 1.94) และพบว่าเพศหญิงมีความเสี่ยงต่อการเกิดอาการต่างๆสูงกว่าเพศชาย

ในการวิเคราะห์ความสัมพันธ์ระหว่างค่าความจุปอด พบว่า  $PM_{2.5}$  มีความสัมพันธ์ต่อค่าความจุปอด คือ เมื่อ  $PM_{2.5}$  เพิ่มขึ้น จะมีผลทำให้ค่าความจุปอดลดลงอย่างมีนัยสำคัญทางสถิติ ส่วน  $PM_{10}$  ไม่พบว่ามีผลกระทบต่อค่าความจุปอด

การศึกษานี้แสดงให้เห็นว่าฝุ่นละอองขนาดเล็กทั้ง  $PM_{10}$  และ  $PM_{2.5}$  มีความสัมพันธ์กับอาการทางระบบหายใจทั้งในผู้ใหญ่และในเด็กโดยเฉพาะกลุ่มที่มีความไวต่อการสัมผัสมลพิษ โดยเฉพาะ  $PM_{2.5}$  ที่สะสมมีผลกระทบต่อค่าความจุปอดของผู้ใหญ่

### ผลการประเมินผลกระทบทางเศรษฐศาสตร์

การศึกษาค่าความพึงพอใจที่จะจ่าย (Willingness to Pay, WTP) ได้ถูกออกแบบเพื่อใช้ควบคู่ไปกับการวิจัยการบันทึกอาการทางระบบหายใจประจำวันเพื่อให้ได้ข้อมูลว่ากลุ่มตัวอย่างจะให้ค่าในการหลีกเลี่ยงการมีอาการทางระบบหายใจเป็นเวลา 1 วัน มากเพียงใด ความท้าทายของการออกแบบสำรวจ WTP สำหรับอาการทางสุขภาพคือ การสร้างคำถามในทางที่จะช่วยให้กลุ่มตัวอย่างคิดในสภาพความเป็นจริงเกี่ยวกับการแลกเปลี่ยน (tradeoffs) ระหว่างรายได้หรือค่าใช้จ่ายกับการเจ็บป่วย ความคิดในการใช้จ่ายเพื่อหลีกเลี่ยงหรือลดการเจ็บป่วย ซึ่งผลจากการศึกษานี้สามารถหาค่าของผลกระทบต่อสุขภาพที่สัมพันธ์กับความเข้มข้นของมลภาวะทางอากาศในกรุงเทพมหานคร ซึ่งจะนำไปสู่การคำนึงถึงประโยชน์ของการควบคุมมลภาวะทางอากาศในกรุงเทพมหานคร

ในการประเมินผลกระทบทางด้านเศรษฐศาสตร์ในการศึกษานี้ ได้ประเมินค่าใช้จ่ายจริงที่เกิดขึ้นเมื่อเกิดอาการทางระบบหายใจในช่วง 3 เดือนที่ผ่านมา และความเต็มใจที่จะจ่าย เพื่อป้องกันไม่ให้เกิดอาการทางระบบหายใจ ค่าใช้จ่ายจริงในกลุ่มผู้ใหญ่สูงกว่าในกลุ่มเด็กทั้งในอาการเล็กน้อย อาการปานกลางและอาการ

มาก และจำนวนเงินที่ยินดีที่จะจ่ายเพื่อป้องกันอาการป่วยทางระบบหายใจน้อยกว่าค่าเฉลี่ยของค่าใช้จ่ายจริง  
ทุกระดับอาการทั้งในเด็กและผู้ใหญ่ ยกเว้นในกลุ่มเด็กสำหรับกรณีอาการมากที่พ่อแม่ยินดีจะจ่ายเงินมากกว่า  
ค่าใช้จ่ายจริง

การศึกษาจากตัวอย่างผู้ใหญ่ 101 คน มีจำนวนครั้งที่ป่วยด้วยอาการทางระบบหายใจรวม 135 ครั้ง  
ในช่วง 3 เดือน เท่ากับ 540 ครั้งใน 1 ปี หรือประมาณ 5 ครั้ง ต่อคนต่อปี และในแต่ละครั้งที่ป่วยสูญเสีย  
ค่าใช้จ่ายในการรักษาพยาบาลประมาณ 186 บาท ดังนั้น ใน 1 คนที่ป่วยจะสูญเสียค่าใช้จ่ายในการ  
รักษาพยาบาลประมาณ 930 บาทต่อปี และถ้าหากประชากรในเชียงใหม่ที่อายุตั้งแต่ 15 ปีขึ้นไป มีอาการ  
ป่วยทางระบบหายใจทุกคน ในระยะเวลา 1 ปี คิดเป็นความสูญเสียทางเศรษฐกิจทั้งหมดประมาณ 2,630.5  
ล้านบาท

โดยสรุปฝุ่นละอองขนาดเล็กทั้ง  $PM_{10}$  และ  $PM_{2.5}$  มีผลกระทบต่อสุขภาพทั้งในประชากรเด็กและ  
ประชากรผู้ใหญ่ที่อาศัยอยู่ในจังหวัดเชียงใหม่และค่าใช้จ่ายที่สูญเสียเนื่องจากการรักษาอาการป่วยทางระบบ  
หายใจที่เกิดขึ้นมีมูลค่ามหาศาล ทั้งนี้ยังไม่ได้รวมถึงมูลค่าความสูญเสียทางสุขภาพของประชาชนเข้าไปด้วย  
ถ้าหากสามารถดำเนินนโยบายต่างๆ เพื่อการป้องกันและลดปริมาณฝุ่นละออง โดยเฉพาะฝุ่นละอองขนาดเล็ก  
ในจังหวัดเชียงใหม่ ก็น่าจะเป็นผลดีอย่างมากต่อสุขภาพของประชาชน

### การนำผลไปใช้ประโยชน์/การเผยแพร่ผลงาน

เผยแพร่ผลการศึกษาในหน่วยงาน

**ชื่อโครงการไทย 4** สื่อสร้างสรรค์จากเพื่อนสู่เพื่อนเพื่อส่งเสริมพฤติกรรมป้องกันการเสพยาบ้าของวัยรุ่นในชุมชนแออัด

**นักวิจัยหลัก** นางสาวภัทรพรรณ เล่านิรมัย

**นักวิจัยในทีม** นางสาวอรพินท์ ไชยพยอม

**ที่ปรึกษา** ศ.นพ.จิตร สิทธิอมร

### หลักการและเหตุผล

จากการเพิ่มของปัญหาที่มาจากครอบครัว เช่น การแยกทางกันของพ่อแม่ การเลี้ยงดูลูก สภาพคล่องทางการเงินของครอบครัว และปัญหาในชุมชน เช่น การเสพยาเสพติด การพนัน ล้วนเป็นปัญหาใหญ่ที่กำลังทวีความรุนแรงมากขึ้น ปัญหาเหล่านี้มีผลกระทบโดยตรงกับวัยรุ่น ที่ส่วนใหญ่ใช้เวลาอยู่กับเพื่อน ทำให้มีโอกาสพึ่งยาเสพติดจากเพื่อน ด้วยความซับซ้อนของปัญหาดังกล่าว จำเป็นต้องให้วัยรุ่นตระหนักถึงความเข้าใจในตนเอง โครงการนี้เป็นจุดเริ่มต้นในการกระตุ้นให้วัยรุ่นเห็นถึงศักยภาพของตนเองผ่านกลุ่มเพื่อนในการแก้ไขปัญหาได้อย่างถูกต้อง

### วัตถุประสงค์

เพื่อสร้างและทดลองใช้กิจกรรมสื่อสร้างสรรค์ในการส่งเสริมความรู้เกี่ยวกับยาบ้า ความตระหนักถึงความรุนแรงของยาบ้า และทักษะชีวิตที่ส่งเสริมพฤติกรรมป้องกันการเสพยาบ้า ให้แก่วัยรุ่นในชุมชนแออัดแห่งหนึ่งในกรุงเทพฯ

### วิธีการวิจัย

ผู้ที่เข้ามามีส่วนร่วมประกอบด้วย ตัวแทนวัยรุ่นในชุมชน เจ้าหน้าที่องค์กรพัฒนาเอกชนในชุมชน และคณะผู้วิจัย โดยมุ่งเน้นให้วัยรุ่นได้มีส่วนร่วมในการจัดกิจกรรมต่างๆ เพื่อให้ได้กิจกรรมที่เหมาะสมกับสภาพชุมชน และตรงตามความต้องการของวัยรุ่นในชุมชนที่เลือกเป็นกรณีศึกษา โดยใช้เวลาในการทดลองดำเนินกิจกรรมประมาณ 1 ปี โครงการฯ ได้จัดกิจกรรมต่างๆ ประกอบด้วย การจัดตั้งมุมเพื่อนใจวัยรุ่น การเปิดให้บริการโทรศัพท์สายด่วนเพื่อนใจวัยรุ่น การจัดค่ายเยาวชน การฝึกอบรมการจัดทำสื่อ การจัดทำจดหมายข่าว และกิจกรรมย่อยอื่นๆ และติดตามประเมินผลการจัดกิจกรรมโดยใช้การสังเกตการณ์แบบมีส่วนร่วม การสัมภาษณ์โดยใช้แบบสอบถาม การสัมภาษณ์แบบเจาะลึก และการสนทนากลุ่ม

### ผลการวิจัย

กิจกรรมในรูปแบบค่ายเยาวชนหรือค่ายอบรมที่จัดขึ้นภายนอกชุมชนเป็นรูปแบบกิจกรรมที่วัยรุ่นส่วนใหญ่ในชุมชนอยากจะเข้าร่วมมากที่สุด แต่มีข้อจำกัดบางประการ เช่น ใช้งบประมาณสูง และวัยรุ่นในชุมชนไม่สามารถเข้าร่วมได้อย่างทั่วถึง ส่วนการจัดตั้งมุมเพื่อนใจวัยรุ่น และการเปิดให้บริการโทรศัพท์สายด่วน ได้รับการตอบรับจากวัยรุ่นในระดับปานกลาง และยังจำกัดกลุ่มผู้ใช้บริการอยู่เพียงบางกลุ่ม แต่เป็นประโยชน์อย่างยิ่งในการใช้เป็นช่องทางในการประชาสัมพันธ์กิจกรรม สำหรับสื่อสิ่งพิมพ์ที่เหมาะสมกับวัยรุ่น

ในชุมชน ควรให้ความสำคัญกับรูปแบบของสื่อที่ดึงดูดความสนใจ ทั้งสีสัน การตกแต่ง รูปภาพ และศิลปกรรม นอกจากนั้น ควรคำนึงถึงความสม่ำเสมอในการเผยแพร่ ความทั่วถึงในการเผยแพร่ และประโยชน์ใช้สอยของสื่อด้วย เพื่อให้วัยรุ่นได้รับทั้งความรู้ และความเพลิดเพลินในการอ่านในเวลาเดียวกัน แต่สิ่งที่สำคัญของการจัดกิจกรรม คือ การเปิดโอกาสให้วัยรุ่นในชุมชนได้มีทางเลือกในการใช้เวลาว่างให้เกิดประโยชน์ แทนที่จะนำเวลาว่างไปใช้ในทางที่ผิด ส่วนประโยชน์อื่นๆ เช่น ความรู้ ความตระหนัก และทักษะชีวิต ถือเป็นผลที่ได้จากการเข้าร่วมกิจกรรมซึ่งจะมองเห็นผลในระยะยาวต่อไป

#### การนำผลไปใช้ประโยชน์/การเผยแพร่ผลงาน

Laoniramai, P., Laosee, O. C., Somrongthong, R., Wongchalee, S., & Sitthi-Amorn, C. (2005). Factors affecting the experiences of drug use by adolescents in a Bangkok slum. *Southeast Asian J Trop Med Public Health*, 36(4), 1014-1019.

**ชื่อโครงการไทย** 5 การมีส่วนร่วมของประชาสังคมในกลยุทธ์เพื่อส่งเสริมการใช้ยาอย่างเหมาะสมเพื่อรักษาโรคติดต่อที่ทางเดินหายใจส่วนบน ของผู้ใหญ่ในระดับครัวเรือนและชุมชน: กรณีศึกษาชุมชนแออัดในกรุงเทพมหานคร

**ชื่อโครงการอังกฤษ** Partnership of civil society in a strategy to reduce the use of antibiotics in the treatment of adults with Upper Respiratory Infection (URIs) from viral origins at the household and community levels: A comparison study in congested community, Bangkok Thailand

**นักวิจัยหลัก** ดร.ศิริตรี สุทธิจิตต์

**ที่ปรึกษา** ศ.นพ.จิตร สิทธิอมร

### หลักการและเหตุผล

โรคติดเชื้อในระบบทางเดินหายใจเป็นโรคที่เกิดขึ้นได้บ่อย ในประเทศไทยพบว่าโรคติดเชื้อในระบบทางเดินหายใจส่วนบนเป็นโรคที่พบได้มากที่สุด สถานพยาบาล และยาปฏิชีวนะที่มีส่วนใหญ่อำนาจรักษาโรคที่เกิดจากการติดเชื้อในระบบทางเดินหายใจส่วนบน จากการใช้อย่างผิดวิธีในการรักษาโรคทางเดินหายใจเพิ่มขึ้นทุกปี ผู้ป่วยเองบางคนซื้อยามารับประทานเอง รับประทานไม่ต่อเนื่อง และเกิดความเข้าใจผิดๆ เกี่ยวกับการรักษาโรคทางเดินหายใจ โดยเฉพาะหวัด ก่อให้เกิดปัญหาการใช้ยาที่ไม่เหมาะสม

### วัตถุประสงค์

เพื่อประเมินประสิทธิภาพของกลยุทธ์ในการลดการใช้ยาปฏิชีวนะที่ไม่เหมาะสมในการรักษาโรคติดเชื้อทางเดินหายใจส่วนบน สำหรับผู้ใหญ่ ในระดับครัวเรือนและชุมชน โดยผ่านการมีส่วนร่วมของประชาสังคม

### วิธีการวิจัย

Pre-post study intervention study with a comparison group ดำเนินการศึกษาในชุมชนแออัดสองแห่งในกรุงเทพฯ โดยครอบคลุมการปรับเปลี่ยนพฤติกรรมของกลุ่มเป้าหมายทั้งสามกลุ่มที่เกี่ยวข้องกับการใช้ยาปฏิชีวนะ คือ ประชาชน ร้านยา และแพทย์ในสถานบริการสาธารณสุขทุกแห่ง Intervention คือการสนับสนุนให้เกิดประชาสังคมในกลุ่มเป้าหมาย ประกอบกับ educational และ managerial strategies ประเมินผลด้วยการสัมภาษณ์ครัวเรือน ล่อซื้อและสัมภาษณ์ร้านยา และการติดตามการสั่งจ่ายยาในสถานบริการสาธารณสุข

### ผลการวิจัย

ความพยายามในการสนับสนุนให้เกิดประชาสังคมด้านยาและสุขภาพในชุมชนแออัด โดยให้ประกอบด้วย 3 ภาคีทั้งผู้ส่งใช้ ผู้จ่าย และผู้ขาย ยังไม่สามารถดำเนินการได้จริง ด้วยการขาดความพร้อมขององค์ประกอบในการประชาสังคมหลายด้าน เช่น มี knowledge gap ระหว่างกลุ่ม ขาดจิตสำนึกประชาสังคม และขาดพื้นที่สำหรับการแลกเปลี่ยนเสวนา เป็นต้น อย่างไรก็ตาม โครงการได้สนับสนุนให้เกิดคู่มือการรักษาโรคติดเชื้อทางเดินหายใจส่วนบนและให้มีการใช้ในแต่ละกลุ่มเป้าหมาย มีกิจกรรมในแต่ละกลุ่มเป้าหมาย เกิดคู่มือและสื่อสำหรับการให้ความรู้ ปรับทัศนคติต่อยาปฏิชีวนะและพฤติกรรมการใช้ยาอย่าง

เหมาะสม ทั้งนี้พบการลดลงในการจ่ายยาปฏิชีวนะของแพทย์ในศูนย์บริการสาธารณสุขทุกหม. แต่ยังไม่พบการปรับพฤติกรรมการใช้ยาในกลุ่มประชาชนและร้านยาที่ชัดเจน จึงควรต้องศึกษาเพิ่มเติมถึงการประยุกต์ใช้แนวคิดเรื่องประชาสังคมกับสุขภาพ และการใช้ยา โดยเฉพาะกับการประยุกต์ในกลุ่มธุรกิจเช่นร้านยา

#### การนำผลไปใช้ประโยชน์/การเผยแพร่ผลงาน

Suttajit, S., Wagner, A. K., Tantipidoke, R., Ross-Degnan, D., & Sitthi-amorn, C. (2005). Patterns, appropriateness, and predictors of antimicrobial prescribing for adults with upper respiratory infections in urban slum communities of Bangkok. *Southeast Asian J Trop Med Public Health*, 36(2), 489-497.

**ชื่อโครงการไทย 6** การวิเคราะห์ความเสี่ยงในการเกิดโรคหัวใจ ชักและการเสียชีวิตจากการศึกษาระยะยาว  
ในชุมชนแออัดคลองเตย

**ชื่อโครงการอังกฤษ** An Analysis of Population Cohort: Risk of Hypertension, Stroke, and Death of  
Residence in Klong Toey Slum

**นักวิจัยหลัก** นางสาวอรพินท์ ไชยพยอม

**ที่ปรึกษา** ศ.นพ. จิตร สิทธิอมร

### หลักการและเหตุผล

จากสถิติโรงพยาบาลระดับศูนย์ในกรุงเทพมหานครพบว่า เกือบครึ่งของผู้ป่วยที่มีเลือดออกในสมองเกิดจากภาวะชัก ผู้ป่วยจำนวนมากมาจากชุมชนแออัดคลองเตย และจากสถิติสาธารณสุขในปีพ.ศ. 2539 พบว่าจำนวนผู้เข้ามารับการรักษาเพิ่มขึ้น โดยเฉพาะผู้ป่วยโรคความดันโลหิตสูงเพิ่มขึ้นอย่างต่อเนื่อง ทำให้รัฐต้องปรับคุณภาพการรักษาให้มีประสิทธิภาพมากขึ้น เพื่อลดการป่วยรุนแรงและการเสียชีวิตอันเป็นผลมาจากภาวะแทรกซ้อนของโรคความดันโลหิตสูงนี้

การศึกษานี้เก็บข้อมูลระยะยาวจากผู้ป่วยในชุมชนแออัดคลองเตยและแฟลตคลองเตยจำนวน 1885 คน ในปี พ.ศ. 2531 และมีการติดตามกลุ่มตัวอย่างดังกล่าวในปี พ.ศ. 2536, 2541 โดยการวัดความดันโลหิต ชั่งน้ำหนัก ประเมินภาวะชัก และการเสียชีวิต อย่างไรก็ตามโครงการนี้ยังขาดการวิเคราะห์เพื่อหาปัจจัยเสี่ยงต่างๆ เหล่านี้

### วัตถุประสงค์

เพื่อวิเคราะห์ปัจจัยเสี่ยงที่ทำให้เกิดการเสียชีวิตด้วยโรคหัวใจ ปัจจัยที่ทำให้เกิดการเกิดภาวะชัก และปัจจัยเสี่ยงที่ทำให้เกิดการเสียชีวิต ในกลุ่มประชาชนที่อาศัยในชุมชนแออัดคลองเตยและแฟลตคลองเตย

### ผลการวิเคราะห์

จากกลุ่มตัวอย่างจำนวน 1,876 คน เพศชาย 867 คน (46%) หญิง 1,009 คน (54%) ความชุกของการเกิดโรคความดันโลหิตสูงมีแนวโน้มเพิ่มขึ้น จาก 20% ในปี พ.ศ.2531 และ 26% ในปีพ.ศ. 2536 ตัวแปรที่มีผลอย่างมีนัยสำคัญทางสถิติ ต่อการเสียชีวิตด้วยโรคหัวใจ คือ เพศ, อายุ, ระยะเวลาที่ดื่ม alcohol ตัวแปรที่มีผลอย่างมีนัยสำคัญทางสถิติ ต่ออาการชัก คือ อายุ กล่าวคือ เมื่อมีอายุเพิ่มขึ้น 1 ปี จะมีโอกาสเกิดชักเพิ่มขึ้น 1.3 เท่า และตัวแปรที่มีอิทธิพลต่อการเสียชีวิตอย่างมีนัยสำคัญทางสถิติ ( $P < 0.05$ ) ต่อการเสียชีวิตโดยรวม คือ อายุ, ระยะเวลาที่ดื่ม alcohol, HDL, Triglyceride

### การนำผลไปใช้ประโยชน์/ การเผยแพร่

เผยแพร่ผลการศึกษาในหน่วยงาน

**ชื่อโครงการ ไทย 7** การติดต่อของเชื้อหวัดสายพันธุ์ A ระหว่างคนและสัตว์ในประเทศไทย

**ชื่อโครงการอังกฤษ** Influenza A Infections at the Human Animal Interface

**นักวิจัยหลัก** ผศ. ดร.รัตนา สำโรงทอง

**ที่ปรึกษา** ศ.นพ. จิตร สิทธิอมร

**นักวิจัยในทีม**

1. อาจารย์ สพ.ญ.ประวีณา กิตติคุณ คณะสัตวแพทยศาสตร์ จุฬาลงกรณ์มหาวิทยาลัย พญาไท แขวงวังใหม่ เขตปทุมวัน กทม. 10330 โทรศัพท์ 02-2189578, 02-2189577
2. นางลินดา วิมลเก็จ ศูนย์เชี่ยวชาญเฉพาะทางด้านไวรัสวิทยาคลินิก คณะแพทยศาสตร์ จุฬาลงกรณ์มหาวิทยาลัย โทรศัพท์ 02-2564909, 02-2564929
3. นางสาวสุนิตรา ภาคอินทรีย์ วิทยาลัยวิทยาศาสตร์สาธารณสุข จุฬาลงกรณ์มหาวิทยาลัย เขตปทุมวัน กรุงเทพฯ โทรศัพท์ 02-2188197 Fax: 02-255 6046

#### หลักการและเหตุผล

ในระหว่างปี 2003-2004 พบการระบาดครั้งใหญ่ของเชื้อดังกล่าวในสัตว์ปีกเป็นจำนวนมาก และมีการทำลายสัตว์ปีกในประเทศไทยมากกว่า 60 ล้านตัว ปัญหาดังกล่าวเกิดติดต่อกันมาเรื่อยๆจนถึงปี 2008 ตั้งแต่ปี 2003 จนถึงปัจจุบันพบว่า มีผู้ติดเชื้อทั่วโลก 359 คน และ เสียชีวิต 226 คน ([www.who.int](http://www.who.int)-accessed February 5, 2008) มีรายงานการระบาดของเชื้อโรคไข้หวัดนกมากถึง 48 ประเทศ ([www.oie.int](http://www.oie.int)-accessed February 6, 2008) เช่น ประเทศ กัมพูชา จีน ไครเอเชีย ฮองกง อินโดนีเซีย ญี่ปุ่น คาซัคสถาน เกาหลี ลาว มาเลเซีย มองโกเลีย รัสเซีย ไทย ตุรกี เวียดนาม เป็นต้น

การระบาดของไข้หวัดนกในประเทศไทย มีการระบาดเป็นระลอก (wave) จำนวน 6 ระลอกแม้ว่าทุกฝ่ายได้ร่วมมือกันแก้ไขปัญหา แต่ยังไม่สามารถรับมือกับการระบาดของโรคได้ ถ้ามีการระบาดของหวัดนกขนาดใหญ่ (pandemic) เข้าสู่คน และแพร่กระจายจากคนสู่คนในภูมิภาคเอเชียจะทำให้มีผู้เสียชีวิตจำนวนมาก นับล้านคน ผู้เสียชีวิตส่วนใหญ่จะเป็นผู้ที่ยังไม่เคยมี ภูมิคุ้มกันของโรคมาก่อน โดยเฉพาะในเด็กหรือวัยรุ่นหากไวรัสดังกล่าวยังคงระบาดในสัตว์ปีก ก็จะทำให้มีโอกาสที่จะเกิดการติดเชื้อในคนและสัตว์อื่นเพิ่มขึ้นซึ่งอาจก่อให้เกิดการระบาดที่รุนแรงมากขึ้นได้

การศึกษาครั้งนี้เป็นส่วนหนึ่งในการพัฒนาศักยภาพของบุคลากร ที่ทำการศึกษาที่เกี่ยวข้องกับการระบาดของไข้หวัดนก โดยจะมีการเชื่อมโยงข้อมูลที่ได้จากห้องปฏิบัติการ (ตัวอย่าง specimen ในคนและสัตว์) กับพฤติกรรมของประชาชนในชุมชนที่มีการระบาดซ้ำซาก เพื่อค้นหาสาเหตุทั้งทางระบาดวิทยา สังคม วัฒนธรรม เพื่อตอบโจทย์การระบาดซ้ำซาก โดยจะนำไปสู่การเกิดความร่วมมือกับภาครัฐ (เช่น มหาวิทยาลัย กระทรวงสาธารณสุข กรมปศุสัตว์ ฯลฯ) ภาคเอกชน (เช่น เจ้าของฟาร์มสัตว์ปีก ผู้ที่ทำธุรกิจเกี่ยวข้องกับสัตว์ปีก ฯลฯ) ส่วนท้องถิ่น (เช่น องค์การปกครองส่วนท้องถิ่น) และชุมชน (เช่น เกษตรกรผู้เลี้ยงไก่บ้าน, ไก่ชน, เป็ด และสัตว์ปีกอื่นๆ) เพื่อให้ได้แนวทางและคำตอบในการเฝ้าระวัง ดูแล แก้ไขปัญหา และการรับมือไข้หวัดใหญ่สายพันธุ์ A และไข้หวัดนก โดยความร่วมมือจากทุกภาคส่วน ทั้งภาครัฐ สถาบันการศึกษา ท้องถิ่น และชุมชน รวมถึงการพัฒนาศักยภาพบุคลากรทางด้านการแพทย์ การสาธารณสุข สัตวแพทย์ และ

นักวิทยาศาสตร์ห้องปฏิบัติการให้มีประสิทธิภาพในการดำเนินโครงการวิจัยให้ได้รับการยอมรับในระดับสากล เกิดทักษะในการพัฒนาโครงการเพื่อแสวงหาทุนวิจัยจากแหล่งทุนภายนอกทั้งในและต่างประเทศ เพื่อศึกษาค้นคว้า และแก้ปัญหาการระบาดของเชื้อหวัดใหญ่สายพันธุ์ A และไข้หวัดนกต่อไป

### วัตถุประสงค์

1. เพื่อหาข้อสรุปสถานการณ์ของการติดต่อของโรค โดยวิเคราะห์ข้อมูลที่มีอยู่ เพื่อแสดงข้อเท็จจริงตามหลักวิชาการเกี่ยวกับไข้หวัดนก ในประเด็นต่างๆ ในพื้นที่ที่มีไก่ตายซ้ำซากในจังหวัดสุพรรณบุรี
2. เพื่อค้นหาพฤติกรรมและปัจจัยเสี่ยงที่มีผลต่อการระบาดของโรคไข้หวัดใหญ่และไข้หวัดนกในพื้นที่ที่กำหนด

### ประโยชน์ที่คาดว่าจะได้รับ

1. ทราบถึงสถานการณ์เกี่ยวกับไข้หวัดนก ในประเด็นต่างๆ จากข้อมูลที่มีอยู่
2. ทราบเกี่ยวกับการติดต่อในไก่พื้นบ้าน ไก่ชน และเป็ดไล่ทุ่ง รวมทั้งไก่ฟาร์ม การติดต่อจากสัตว์ปีกสู่คน โดยเน้นเฉพาะพื้นที่ในอำเภอเมือง และ/หรืออำเภออู่ทอง จังหวัดสุพรรณบุรี
3. โครงการวิจัยเพื่อป้องกันการระบาดของโรคไข้หวัดนก โดยร่วมมือกันของ กรมปศุสัตว์ สำนักงานสาธารณสุขจังหวัด มหาวิทยาลัย องค์กรต่างๆ ทั้งในและต่างประเทศ

ขอบเขตการดำเนินงาน ดำเนินโครงการวิจัยใน 2 อำเภอ ได้แก่ อ.เมือง และ อ.อู่ทอง จังหวัดสุพรรณบุรี

วิธีการวิจัย นักวิจัยได้พื้นที่วิจัยนำร่อง 2 แห่งโดยมีข้อกำหนดในการเลือกพื้นที่ คือ

1. พื้นที่ที่มีอัตราการตายของสัตว์ปีกสูง
2. มีความร่วมมือของปศุสัตว์และสาธารณสุขจังหวัด
3. ประชาชน ชุมชนและ NGO ให้ความร่วมมือ
4. หมู่บ้านที่มีการระบาดของหวัดนกซ้ำซาก

รูปแบบการวิจัยใช้วิธีการวิจัยแบบมีส่วนร่วมเพื่อตั้งคำถาม และใช้ระเบียบวิธีการวิจัยในการหาคำตอบ ความร่วมมือดังกล่าวจะมาจากส่วนท้องถิ่น NGO ในพื้นที่ หน่วยงานทางวิชาการ เพื่อที่จะวางแผนดำเนินการ ประเมินผล

### การนำประโยชน์ไปใช้/การเผยแพร่ผลงาน

Somrongthong, R., Beaudomin, A.L., Pakinsee, S., & Sitthi-amorn, C. (2010). Folk Knowledge about Avian Influenza and the Use of Personal Protective Equipment: A Qualitative Study. *J Health Res*, 24(Suppl 1), 27-32.

### สรุปผลลัพธ์ (Output) ที่ได้จากโครงการ

1. โครงการย่อย 17 โครงการ    1) ผลงานตีพิมพ์ ทั้งหมด 17 บทความ
  - ระบับนานาชาติ จำนวน 16 บทความ
  - ระบับชาติ จำนวน 1 บทความ  
2) นำเสนอในผลงานวิจัยในที่ประชุม
  - ระบับนานาชาติ 7 ครั้ง
  - ระบับชาติ 5 ครั้ง
2. ผลงานตีพิมพ์เพิ่มเติม จำนวน 7 บทความ
3. การบรรยายพิเศษในที่ประชุมระบับนานาชาติ จำนวน 3 ครั้ง

## 1. โครงการย่อย

ลำดับ	ชื่อโครงการย่อย	ผู้วิจัยหลักโครงการย่อย	การเผยแพร่ผลงาน				
			ชื่อผลงาน	การตีพิมพ์		การนำเสนอผลงานวิจัยในที่ประชุม	
				ระดับนานาชาติ	ระดับชาติ	ระดับนานาชาติ	ระดับชาติ
1.	The cost effectiveness of early detection of bancroftion filarasis in Myanmar Migrants using antigen-capsule ELISA, IgG4 antibody assays and PCR technique	ศ.พญ.สุรางค์ ไตรวีระประภาพร					
2.	รูปแบบการตรวจหาเชื้อพยาธิเม็ดเลือดในไมโครบินชนิดปกติ	นพ.กิตติ ต่อจรัส					
3.	The effectiveness of Shared Care for patients with Epilepsy in Nakornratchasima	นพ.ธนินทร์ อัครวิเชียรจินดา	1) Prevalence of epilepsy in rural Thailand: a population-based study 2) Compliance with treatment of adult epileptics in a rural district of Thailand	✓ ✓	✓	✓	✓
4.	การศึกษาเพื่อหาปัจจัยเสี่ยงของผู้ป่วยกล้ามเนื้อหัวใจตายเฉียบพลันในกลุ่มควบคุม - INTERHEART	ศ.นพ.จิตร์ ลิขชีมวร	1) Association of psychosocial risk factors with risk of acute myocardial infarction in 11,119 cases and 13,648 controls from 52 countries: case-control study 2) Effect of potentially modifiable risk factors associated with myocardial infarction in 52 countries (the INTERHEART study): case-control study	✓ ✓	✓	✓	
5.	การศึกษาคอร์บอร์วี่ไหลตาย	นพ.สมเกียรติ แสงวัฒนาโรจน์	The Right Ventricular Electrocardiographic leads for detections of Brugada Syndrome in Sudden Unexplained Death Syndrome survivors and their relatives	✓			
6.	Trend of Cardiopulmonary Resuscitation Patients with Terminal Illness in Chiang Mai	ผศ.ดร.สุตารัตน์ ลิขชีสมบัติ	1)Cardiopulmonary resuscitation performed in patients with terminal illness in Chiang Mai University Hospital, Thailand 2)Effectiveness of advance directives for the care of terminally ill patients in Chiang Mai Hospital, Thailand	✓ ✓		✓	✓

ลำดับ	ชื่อโครงการย่อย	ผู้วิจัยหลัก/โครงการย่อย	การเผยแพร่ผลงาน				
			ชื่อผลงาน	การตีพิมพ์		การนำเสนอผลงานวิจัยในที่ประชุม	
				ระดับนานาชาติ	ระดับชาติ	ระดับนานาชาติ	ระดับชาติ
7.	Attitudes towards advance directives and the impact of prognostic information on the preference for cardiopulmonary resuscitation on medical inpatients in CMU Hospital, Thailand	ผศ.ดร.สุภารัตน์ สิกขิสมบัติ	1) Attitudes toward advance directives and the impact of prognostic information on the preference for cardiopulmonary resuscitation in medical inpatients in Chiang Mai University Hospital, Thailand 2) Physicians' attitudes and practices regarding advanced end-of-life care planning for terminally ill patients at Chiang Mai University Hospital, Thailand	✓			
8.	โครงการการตรวจสุขภาพประจำปี และปัจจัยที่เกี่ยวข้องของผู้สูงอายุในเขตชนบทของจ.ขอนแก่น	ผศ.จิราพร เทียวอยู่					
9.	การศึกษาความเข้าใจ ความน่าเชื่อถือ ความสมบูรณ์ และความไวต่อการเปลี่ยนแปลงของแบบสอบถาม Health Assessment Questionnaire ฉบับภาษาไทย สำหรับผู้ป่วยโรคข้ออักเสบ	รศ.พญ.มนาริปี โอศิริ	Comprehensibility, reliability, validity and responsiveness of the Thai version of the Health Assessment Questionnaire in Thai patients with rheumatoid arthritis	✓			
10.	การศึกษาความชุก ภาวะความรุนแรง ปัจจัยและผลที่เกิดขึ้นจากเนื้อตายในผู้ป่วยถูกงูเห่าและงูกะปะกัด	ดร.พวลน้อง วงศ์ทองคำ	1) A study of 225 Malayan pit viper bites in Thailand. 2) A study of Thai cobra (Naja kaouthia) bites in Thailand.	✓ ✓		✓	✓

ลำดับ	ชื่อโครงการย่อย	ผู้วิจัยหลักโครงการย่อย	การเผยแพร่ผลงาน				
			ชื่อผลงาน	การตีพิมพ์		การนำเสนอผลงานวิจัยในที่ประชุม	
				ระดับนานาชาติ	ระดับชาติ	ระดับนานาชาติ	ระดับชาติ
1.	ผลของการปรับเปลี่ยนการบริโภคมาเป็นข้าวกล้อง งา ผัก ผลไม้ ตามฤดูกาลของประชาชนในเขตเทศบาลในเขตเทศบาลในจังหวัดสุโขทัย	ภก.สมนึก สุขยธนาวิช					
2.	ความต้องการและปัญหาด้านสุขภาพ การเข้าถึงบริการ และคุณภาพชีวิตของวัยรุ่นกับการสร้างความร่วมมือ	ผศ.ดร.รัตนา ลำโรงทอง	1) Sexual behaviors and opinions on sexuality of adolescents in a slum community in Bangkok 2) Factors affecting the experiences of drug use by adolescents in a Bangkok slum	✓ ✓	✓	✓	✓
3.	การศึกษาผลกระทบของผู้และของชนชาติเล็กต่อสุขภาพของประชาชนใน จ.เชียงใหม่	รศ.ดร.นันทวรรณ วิจิตรวาทการ					
4.	สื่อสร้างสรรค์จากเพื่อนสู่เพื่อนเพื่อส่งเสริมพฤติกรรมการป้องกันการใช้ยาของวัยรุ่นในชุมชนแออัด	นส.ภัทราพรณ เล้าภิรัมย์	Factors affecting the experiences of drug use by adolescents in a Bangkok slum	✓			
5.	การมีส่วนร่วมของประชาสังคมในกลยุทธ์เพื่อส่งเสริมการใช้ยาอย่างเหมาะสมสำหรับการรักษาโรคติดต่อที่ทางเดินหายใจส่วนบนของผู้ใหญ่ในระดับครัวเรือนและชุมชน: กรณีศึกษาชุมชนแออัดในกรุงเทพมหานคร	ดร.ศิริตรี สุทธิจิตต์	Patterns, appropriateness, and predictors of antimicrobial prescribing for adults with upper respiratory infections in urban slum communities of Bangkok	✓	✓		✓
6.	การวิเคราะห์ความเสียหายในการเกิดโรคหัวใจ ชัก และการเสียชีวิตจากการศึกษาระยะยาวในชุมชนแออัดคลองเตย	นางสาวอรพินท์ ไชยพยอม					
7.	Influenza A Infections at the Human Animal Interface	ผศ.ดร.รัตนา ลำโรงทอง	Folk Knowledge about Avian Influenza and the Use of Personal Protective Equipment: A Qualitative Study		✓	✓	

## 2. ผลงานตีพิมพ์เพิ่มเติม

1. Priority setting for health research: lessons from developing countries, Health Policy and Planning; 15(2): 130-136 Oxford University Press 2000.
2. Strengthening health research capacity in developing countries: a critical element for achieving health equity. BMJ 2000; 321; 813-817.
3. Globalization and health viewed from three parts of the world. Bulletin of the World Health Organization 2001; 79 (9); 889-890.
4. The Asian Voice in building equity in health for development from the Asian Forum for Health Research, Manila, February 2000. Health Policy and Planning; 17 (2): 213-217 Oxford University Press 2002.
5. Rebuilding Health Systems towards Health Security: Some essential Indicators. In: Health Development in the South-East Asia Region: An Overview. World Health Organization 2004; 75-91.
6. Information systems and community diagnosis in developing countries. Oxford textbook of Public Health, 2001.
7. Prescription pattern for treatment of hemorrhoids under the universal coverage policy of Thailand. Southeast Asian J Trop Med Public Health. Vol. 36 No. 4 July 2005.

## 3. การบรรยายพิเศษในที่ประชุมระดับนานาชาติ

1. Medicines as emerging technologies: unprecedented opportunity, continuing disparity. Keynote Address to the International Conference on Improving Use of Medicine (ICIUM 2004). Chiangmai: March 2004.
2. Childhood Morbidity and Mortality in Thailand. Keynote Address to the UNICEF/TASC Conference on Child Injury. Bangkok: April 2004.
3. Injury & Equity. Keynote Address to the XVII<sup>th</sup> International Epidemiological Association World Congress of Epidemiology. Bangkok: August 2005.

### List of Publications

1. Asawavichienjinda, T., Sitthi-Amorn, C., & Tanyanont, W. (2002). Prevalence of epilepsy in rural Thailand: a population-based study. *J Med Assoc Thai*, 85(10), 1066-1073.
2. Asawavichienjinda, T., Sitthi-Amorn, C., & Tanyanont, W. (2003). Compliance with treatment of adult epileptics in a rural district of Thailand. *J Med Assoc Thai*, 86(1), 46-51.
3. Rosengren, A., Hawken, S., Ounpuu, S., Sliwa, K., Zubaid, M., Almahmeed, W. A., et al. (2004). Association of psychosocial risk factors with risk of acute myocardial infarction in 11119 cases and 13648 controls from 52 countries (the INTERHEART study): case-control study. *Lancet*, 364(9438), 953-962.
4. Yusuf, S., Hawken, S., Ounpuu, S., Dans, T., Avezum, A., Lanas, F., et al. (2004). Effect of potentially modifiable risk factors associated with myocardial infarction in 52 countries (the INTERHEART study): case-control study. *Lancet*, 364(9438), 937-952.
5. Sittisombut, S., Love, E. J., & Sitthi-Amorn, C. (2001). Cardiopulmonary resuscitation performed in patients with terminal illness in Chiang Mai University Hospital, Thailand. *Int J Epidemiol*, 30(4), 896-898.
6. Sittisombut, S., Maxwell, C., Love, E. J., & Sitthi-Amorn, C. (2008). Effectiveness of advance directives for the care of terminally ill patients in Chiang Mai University Hospital, Thailand. *Nurs Health Sci*, 10(1), 37-42.
7. Sittisombut, S., Love, E. J., & Sitthi-Amorn, C. (2005). Attitudes toward advance directives and the impact of prognostic information on the preference for cardiopulmonary resuscitation in medical inpatients in Chiang Mai University Hospital, Thailand. *Nurs Health Sci*, 7(4), 243-250.
8. Sittisombut, S., Maxwell, C., Love, E. J., & Sitthi-Amorn, C. (2009). Physicians' attitudes and practices regarding advanced end-of-life care planning for terminally ill patients at Chiang Mai University Hospital, Thailand. *Nurs Health Sci*, 11(1), 23-28.
9. Osiri, M., Wongchinsri, J., Ukritchon, S., Hanvivadhanakul, P., Kasitanon, N., & Siripaitoon, B. (2009). Comprehensibility, reliability, validity, and responsiveness of the Thai version of the Health Assessment Questionnaire in Thai patients with rheumatoid arthritis. *Arthritis Res Ther*, 11(4), R129.
10. Wongtongkam, N., Wilde, H., Sitthi-Amorn, C., & Ratanabanangkoon, K. (2005a). A study of 225 Malayan pit viper bites in Thailand. *Mil Med*, 170(4), 342-348.
11. Wongtongkam, N., Wilde, H., Sitthi-Amorn, C., & Ratanabanangkoon, K. (2005b). A study of Thai cobra (*Naja kaouthia*) bites in Thailand. *Mil Med*, 170(4), 336-341.
12. Somrongthong, R., Panuwatsuk, P., Amarathithada, D., Chaipayom, O., & Sitthi-amorn, C. (2003). Sexual behaviors and opinions on sexuality of adolescents in a slum community in Bangkok. *Southeast Asian J Trop Med Public Health*, 34(2), 443-446.
13. Somrongthong, R. & Sitthi-amorn, C. (2000). Existing health needs and related health services for adolescents in a slum community in Thailand. *Int J Adolesc Med Health*, 12 (2-3): 191-203.

14. Laoniramai, P., Laosee, O. C., Somrongthong, R., Wongchalee, S., & Sitthi-Amorn, C. (2005). Factors affecting the experiences of drug use by adolescents in a Bangkok slum. *Southeast Asian J Trop Med Public Health*, 36(4), 1014-1019.
15. Suttajit, S., Wagner, A. K., Tantipidoke, R., Ross-Degnan, D., & Sitthi-amorn, C. (2005). Patterns, appropriateness, and predictors of antimicrobial prescribing for adults with upper respiratory infections in urban slum communities of Bangkok. *Southeast Asian J Trop Med Public Health*, 36(2), 489-497.
16. Somrongthong, R., Beaudomin, A.L., Pakinsee, S., & Sitthi-amorn, C. (2010). Folk Knowledge about Avian Influenza and the Use of Personal Protective Equipment: A Qualitative Study. *J Health Res*, 24(Suppl 1), 27-32.
17. Sangwatanaroj, S., Prechawat, S., Sunsaneewitayakul, B., Sittthisook, S., Tosukhowong, P., & Tungsanga, K. (2001). Right Ventricular Electrocardiographic Leads for Detection of Brugada Syndrome in Sudden Unexpected Death Syndrome Survivors and Their Relatives. *Clin. Cardiol*, 24, 776-781.

---

## Prevalence of Epilepsy in Rural Thailand : A Population-Based Study

---

THANIN ASAWAVICHJENJINDA, M.D.\*,  
CHITR SITTHI-AMORN, M.D., Ph.D.\*\*,  
WACHARIN TANYANONT, MS.\*\*\*

### Abstract

**Background :** A study of the community-based burden of illness based upon prevalence is needed to plan intervention strategy.

**Purpose :** To determine the prevalence of epilepsy in a rural population in Thailand.

**Method :** From January to June, 2000, all of the people in Talardkav sub-district were invited to be interviewed and examined by a neurologist who visited their village.

**Results :** Of the 2,069 people in 553 households who gave information (72.2% of the total households), 43 had a history of seizure and of the 43, 15 were epileptics. The prevalence of epilepsy was estimated at 7.2 per 1,000 population. The highest two peaks were in the age groups of 5-9 and 25-34 years (17.0, 17.4/1,000, respectively).

**Conclusion :** The prevalence of epilepsy in rural Thailand is low, although probably underestimated, but it is the best to date for rural Thailand.

**Key word :** Epilepsy, Epidemiology, Population-Based Study, Prevalence, Thailand

ASAWAVICHJENJINDA T,  
SITTHI-AMORN C, TANYANONT W  
J Med Assoc Thai 2002; 85: 1066-1073

\* Neurological Unit, Department of Medicine, Maharat Nakhon Ratchasima Hospital, Nakhon Ratchasima 30000,

\*\* Department of Medicine, Faculty of Medicine and College of Public Health, Chulalongkorn University, Bangkok 10330,

\*\*\* College of Public Health, Chulalongkorn University, Bangkok 10330, Thailand.

Epilepsy is a disorder with significant social and economic consequences<sup>(1)</sup>. In Thailand, it is the third commonest neurological disease after cerebrovascular disease and headache<sup>(2)</sup>.

Prevalence, as a measure of the disease burden in the community, should be considered when planning the health needs at local, regional and national levels<sup>(3)</sup>. The only previous study of the prevalence of epilepsy in Thailand reported a very low prevalence rate (2.8/1,000 population in those aged 15 years or more)<sup>(4)</sup>.

The objective of this study was to determine the prevalence of epilepsy in a rural community in Thailand.

## MATERIAL AND METHOD

This study was conducted in Nakhon Ratchasima province, the second largest of the 76 provinces in Thailand, located 254 km northeast of the capital, Bangkok. The province has an area of 20,494 km<sup>2</sup>, divided into 26 districts, 287 sub-districts and has a total population of 1.77 million. The sub-district Talardkav, an average rural area, was chosen to survey people with a history of epilepsy because of its proximity (within 40 kilometers) to the Provincial Hospital. This sub-district has a population of 3,258, residing in 766 households in 11 villages in an area of 25 km<sup>2</sup>. This population estimate is based upon a survey done by health volunteers and sub-district health officers every six months.

In this study, EPILEPSY was defined as two or more clinical afebrile seizures

- Unrelated to acute metabolic derangement or to withdrawal from alcohol or drugs<sup>(5)</sup>.

- Excluding those which occur within a 24-hour period<sup>(5)</sup>.

- Excluding those in which the diagnosis of epilepsy was questionable.

Each village in Talardkav sub-district was visited between January and June, 2000. In preparation, the authors met with all sub-district health officials, health volunteers and teachers in each village to discuss the objective and rationale of the survey. For seven days before the study, the sub-district health officers announced the date of study by the village intercom and the health volunteers went from door to door to tell all of the people whether they had or did not have a health problem, the date that the neurologist would visit their village to examine the people health, providing free common drugs and

education related to their health problems and to invite them to have an examination on that date.

On the study day, a team consisting of sub-district health officials, health volunteers conducted general examinations (body weight and blood pressure) of all of the people and then the neurologist administered the screening questionnaire (Placentia et al)<sup>(6)</sup> in groups, consisting of six to eight persons from two to three families. They provided information about themselves and their children, and about those who were mentally retarded, demented or psychotic or who were not present. If the screening was suggestive of epilepsy, the neurologist conducted an in-depth interview with the individual; for those retarded, demented, psychotic or absent and for young children, with the parents or caregivers; for school aged children with both their parents and teachers. All of them were examined by the neurologist and provided with common drugs and education related to their health problems.

## RESULTS

### Population and sample characteristics

During the survey, 2,069 (69.5%) in 553 households of the total 3,258 people in 766 households gave the information. The remaining 1,189 did not because nobody in the household came and it was not possible to determine why they did not.

The gender and age specific response rates of receiving the information are shown in Table 1. Overall and in all but the age group 25-34 years, the response rates were higher among the females; significantly so only for the overall response rate and for the response rates in the age groups 10-14 and 55-64 years. There was a reversal of this observation in the age group 25-34 years, where the response rate was higher but not significantly so among the males. The response rates were lowest among those aged 15-44 years, (56.9%), with higher rates in the children (74.5%) and in the older adults (65.9%); p-value <0.001.

### Seizure and epilepsy distribution

Among the 2,069 subjects who gave information, there were 43 (20.8/1000) with a history of seizure (Table 2). Of these 43, only 15 were diagnosed as epileptics. More of the 28 non-epileptics were male (52.9%) and most (82.1%) had had a febrile convulsion. Seizures related to withdrawal from alcohol were only found in the males.

Table 1. Gender and age specific response rates.

Age group (year)	Males		Females		Both genders		P-value (age specific)
	n/N	%	n/N	%	n/N	%	
0-4	83/110	75.5	96/115	83.5	179/225	79.6	ns
5-9	82/120	68.3	94/129	72.9	176/249	70.7	ns
10-14	94/137	68.6	97/122	79.5	191/259	73.7	<0.05
15-24	159/278	57.2	164/272	60.3	323/550	58.7	ns
25-34	148/250	59.2	140/267	52.4	288/517	55.7	ns
35-44	128/234	54.7	150/262	57.3	278/496	56.0	ns
45-54	100/163	61.3	123/179	68.7	223/342	65.2	ns
55-64	76/126	60.3	105/141	74.5	181/267	67.8	<0.05
65+	89/149	59.7	141/204	69.1	230/353	65.2	ns
All age groups	959/1,567	61.2	1,110/1,691	65.6	2,069/3,258	63.5	<0.01

n = Respondents

N = Population

ns = not significant

Table 2. Distribution by gender in those with seizures and epilepsy.

Seizure classification	Male		Female		Total	
	N	%	N	%	N	%
<b>A. Total Epileptics</b>	<b>5</b>	<b>100.0</b>	<b>10</b>	<b>100.0</b>	<b>15</b>	<b>100.0</b>
I. Generalized type	3	60.0	3	30.0	6	40.0
- Generalized tonic clonic	3	60.0	1	10.0	4	26.6
- Absence	0	0	1	10.0	1	6.7
- Generalized tonic	0	0	1	10.0	1	6.7
II. Localization related type	1	20.0	6	60.0	7	46.7
- Simple partial	0	0	1	10.0	1	6.7
- Complex partial	0	0	1	10.0	1	6.7
- Partial with secondarily generalized	1	20.0	4	40.0	5	33.3
III. Unclassified	1	20.0	1	10.0	2	13.3
<b>B. Total Non-epileptics</b>	<b>17</b>	<b>100.0</b>	<b>11</b>	<b>100.0</b>	<b>28</b>	<b>100.0</b>
I. Febrile convulsion	14	82.4	9	81.8	23	82.1
II. Single seizure	0	0	1	9.1	1	3.6
III. Seizure related to						
a) Acute Metabolic condition	2	11.8	1	9.1	3	10.7
b) Withdrawal from alcohol	1	5.9	0	0	1	3.6

In contrast, most of the 15 epileptics were female (66.7%). Slightly more of the epileptics were classified as having a localization related type of epilepsy (46.7%).

#### Prevalence rates

Of the 2,069 who gave information, only 15 were considered to be epileptics. This would yield a crude prevalence rate of epilepsy of 7.2 per 1,000 population (15/2,069) or a rate of 4.6/1,000 (15/3,258) if all the non-givers were assumed to have no history of epilepsy.

As shown in Table 3, the prevalence in females (9.0/1,000) was nearly twice that of the males (5.2/1,000); with the difference greatest in three age groups: 5-9 years (21.3 vs 12.2/1,000); 25-34 years (28.6 vs 6.8/1,000) and 45-54 years (16.3 vs 0/1,000).

#### DISCUSSION

Although examination was offered to all residents of Talardkav Sub-District, the response rate of receiving information was only 69.5 per cent. Most Thai people in a rural area are likely to visit a mobile medical service to obtain medication even if they

Table 3. Gender and age specific prevalence of epilepsy.

Age group (year)	Males		Females		Total		P-value
	n <sub>1</sub> /n	n <sub>1</sub> /1,000	n <sub>1</sub> /n	n <sub>1</sub> /1,000	n <sub>1</sub> /n	n <sub>1</sub> /1,000	
0-4	0/83	0	0/96	0	0/179	0	-
5-9	1/82	12.2	2/94	21.3	3/176	17.0	ns
10-14	0/94	0	1/97	10.3	1/191	5.2	ns
15-24	1/159	6.3	0/164	0	1/323	3.1	ns
25-34	1/148	6.8	4/140	28.6	5/288	17.4	ns
35-44	1/128	7.8	0/150	0	1/278	3.6	ns
45-54	0/100	0	2/123	16.3	2/223	9.0	ns
55-64	1/76	13.2	1/105	9.5	2/181	11.0	ns
65+	0/89	0	0/141	0	0/230	0	-
Total	5/959	5.2	10/1,110	9.0	15/2,069	7.2	ns

n = Respondents

n<sub>1</sub> = Epileptics

ns = not significant

have no definite medical complaint. They would only miss this opportunity when they thought their health was good and had something more important to do. It is possible that those not attending were concerned that their history of epilepsy would become known to either health officers or the community and, as a result, they or their family may become stigmatized. The same may be true for those who attended because the questionnaire was administered in a group setting. Epilepsy is a clinical diagnosis. Patient and eyewitness's information, therefore, is the key to making the diagnosis. However, most types of epilepsy involved loss of consciousness and then amnesia. Most of the time, the diagnosis, hence, can be made from only eyewitness's information.

There were 43 people with a history of seizures but only 15 were considered to be epileptic. In the non-epileptics, there was no marked difference between the males and females in the percentage of febrile convulsion and of seizure related to an acute metabolic condition.

Localization-related type of epilepsy was more common (46.7%) than the generalized type of epilepsy (40.0%). Neurocysticercosis, birth and accident related to head trauma might contribute to this high prevalence of partial epilepsy<sup>(3)</sup>. Generalized epilepsy has been the most common type reported in most community-based surveys<sup>(3,7-18)</sup>. This variation may be related to a lower level of case ascertainment of partial epilepsy, misclassification of a proportion of seizure<sup>(3,9)</sup>, the time<sup>(3)</sup> and the method of diagnosis<sup>(13)</sup>.

The crude prevalence rate of epilepsy in this study ranged from 4.6 to 7.2 per 1,000 population, which was quite low. However, similar prevalence rates have been reported by all community-based studies in Asia<sup>(3,4,7,19)</sup> (APPENDIX 1). Most studies in developing countries excluding Asia have found prevalence rates ranging from 8.5-57/1,000<sup>(8-13,20-23)</sup>. In developed countries, prevalence rates range from 7.5-10.4/1,000<sup>(14,24,25)</sup>. The exceptions are the studies done in Igbo-Ora, Nigeria (5.3/1,000)<sup>(20)</sup> and Riposto, Italy (2.7/1,000)<sup>(15)</sup>. Moreover, the difference in prevalence rates between developing and developed countries as shown in APPENDIX 1, are not marked. In the hospital-based studies, the prevalence rates showed wide variations ranging from 2.0-20.3/1,000<sup>(16-18,26,27)</sup>.

In a previous study in Thailand, the prevalence of diagnosed epilepsy in the population aged  $\geq$  fifteen years was 2.8/1,000<sup>(4)</sup>, which is considerably lower than that found in the present study (4.6-7.2/1,000). The reason might be that the previous study was done by asking people whether they had been diagnosed as epileptic and that different definitions and criteria for making the diagnosis of epilepsy were used by the physicians who examined the patients<sup>(4)</sup>.

The reasons for the low prevalence ratio of epilepsy in the present study might be the method of questionnaire administration and of receiving information. Epileptics with no obvious convulsion might be missed. Recall bias is possible; maybe intentional, possibly due to the fear of stigmatization or unintentional because of the forgetfulness or ignorance of

people to recall their previous symptoms. Incomplete information for making the diagnosis and the diagnostic method [e.g. no access to electroencephalography (EEG)] may also be responsible.

Most studies<sup>(3,7,11,13,15,19,20,24)</sup> have reported a higher prevalence of epilepsy in men, probably due to a higher frequency of head injury<sup>(13)</sup>. In the present study, gender-specific prevalence in women was nearly twice as high as in men (9.0/1,000 *versus* 5.2/1,000) without statistical difference. One of the reasons for the difference from many reported studies is the lower response rates among the males, possibly because even if they have epilepsy they need to work to support their household. The other maybe men do not admit the occurrence of seizures as frequently as women in community surveys<sup>(13)</sup>.

The age-specific prevalence of epilepsy in this study classified in accordance with some other studies demonstrated that the prevalence of epilepsy was highest in two peaks, 5-9 and 25-34 age groups (17.0 and 17.4/1,000, respectively), which was similar to the data from the community of Riposto, Italy<sup>(15)</sup>. A possible cause of the highest two peaks' prevalence rate is the common onset of epilepsy in childhood and in young adult life<sup>(7)</sup>. In developing coun-

tries, most peaked in the second decade<sup>(3,9-11,13,14,19-21)</sup>. However, in developed countries, it has been shown to be higher with increasing age<sup>(14,16-18,27)</sup>. The lower prevalence rate of epilepsy in the elderly in developing countries may be related to multiple factors such as lower life expectancy<sup>(3)</sup>, higher seizure-related mortality<sup>(3)</sup>, and under-ascertainment of seizure disorders in this population<sup>(3)</sup>.

The prevalence of epilepsy in rural Thailand was quite low compared to studies in developing countries<sup>(8,10,12,13,21-23)</sup>. Although this figure is the best to date, it is likely an underestimate of the true prevalence.

#### ACKNOWLEDGEMENT

The authors wish to thank the Thailand Research Fund for funding the study, and also the Director of Maharat Nakhon Ratchasima Hospital, the Provincial Chief Medical Officer Nakhon Ratchasima, the Head of the Department of Medicine and the Department of Social Health, the Director of Chokchai Community Hospital, the Head of the Chokchai District Health Office, and all the Talardkav sub-district health officials and health volunteers.

## REFERENCES

- Garnett WR. Antiepileptic drug treatment: Outcomes and adherence. *Pharmacotherapy* 2000; 20: 191-9.
- Boongird P, Soranastaporn S, Menken M, Vejajiva A. Spectrum of neurological diseases in Thailand. *Neurol J Southeast Asia* 1996; 1: 65-7.
- Radhakrishnan K, Pandian JD, Santhoshkumar T, et al. Prevalence, knowledge, attitude, and practice of epilepsy in Kerala, South India. *Epilepsia* 2000; 41: 1027-35.
- Chuprapawan C. First report of health status nationwide survey of Thai people by Interview and physical examination in 1991-1992. Chuprapawan C; ed. Bangkok: Design Company Limited, 1996: 78-81.
- Guidelines for epidemiologic studies on epilepsy: Commission on epidemiology and prognosis. International League Against Epilepsy. *Epilepsia* 1993; 34: 592-6.
- Placencia M, Suarez J, Crespo F, et al. A large-scale study of epilepsy in Ecuador: Methodology aspects. *Neuroepidemiology* 1992; 11: 74-84.
- Li SC, Schoenberg BS, Wang CC, Cheng XM, Zhou SS, Bolis CL. Epidemiology of epilepsy in urban areas of the People's Republic of China. *Epilepsia* 1985; 26: 391-4.
- Gracia F, Lao SL, Castillo L, et al. Epidemiology of epilepsy in Guaymí Indians from Bocas del Toro Province, Republic of Panama. *Epilepsia* 1990; 31: 718-23.
- Mendizabal JE, Salguero LF. Prevalence of epilepsy in a rural community of Guatemala. *Epilepsia* 1996; 37: 373-6.
- Gomez JG, Arciniegas E, Torres J. Prevalence of epilepsy in Bogota, Columbia. *Neurology* 1978; 28: 90-4.
- Rwiza HT, Kilonzo GP, Haule L, et al. Prevalence and incidence of epilepsy in Ulanga, a rural Tanzania district: A community-based study. *Epilepsia* 1992; 33: 1051-6.
- Placencia M, Shorvon SD, Paredes V, et al. Epileptic seizures in an Andean region of Ecuador: Incidence and prevalence and regional variation. *Brain* 1992; 115: 771-82.
- Nicoletti A, Reggio A, Bartoloni A, et al. Prevalence of epilepsy in rural Bolivia: A door-to-door survey. *Neurology* 1999; 53: 2064-9.
- Haerer AF, Anderson DW, Schoenberg BS. Prevalence and clinical features of epilepsy in a biracial United States population. *Epilepsia* 1986; 27: 66-75.
- Reggio A, Failla G, Patti F, et al. Prevalence of epilepsy. A door-to-door survey in the Sicilian community of Riposto. *Ital J Neurol Sci* 1996; 17: 147-51.
- Cockerell OC, Eckle I, Goodridge DMG, Sander JWA, Shorvon SD. Epilepsy in a population of 6000 re-examined: Secular trends in first attendance rates, prevalence, and prognosis. *J Neurol Neurosurg Psychiatry* 1995; 58: 570-6.
- Olafsson E, Hauser WA. Prevalence of epilepsy in rural Iceland: A population-based study. *Epilepsia* 1999; 40: 1529-34.
- Wright J, Pickard N, Whitfield A, Hakin N. A population-based study of the prevalence, clinical characteristics and effect of ethnicity in epilepsy. *Seizure* 2000; 9: 309-13.
- Singh A, Kaur A. Epilepsy in rural Haryana-Prevalence and treatment seeking behaviour. *J Indian Med Assoc* 1997; 95: 37-47.
- Osuntokun BO, Adeuja AOG, Nottidge VA, et al. Prevalence of the epilepsies in Nigerian Africans: A community-based study. *Epilepsia* 1987; 28: 272-9.
- Osuntokun BO, Schoenberg BS, Nottidge VA, et al. Research protocol for measuring the prevalence of neurologic disorders in developing countries. *Neuroepidemiology* 1982; 1: 143-53.
- Debrock C, Preux PM, Houinato D, et al. Estimation of the prevalence of epilepsy in the Benin region of Zinvie using the capture-recapture method. *Int J Epidemiol* 2000; 29: 330-5.
- Kaamugisha J, Feksi AT. Determining the prevalence of epilepsy in the semi-urban population of Nakuru, Kenya, comparing two independent methods not apparently used before in epilepsy studies. *Neuroepidemiology* 1988; 7: 115-21.
- Karaagac N, Yeni SN, Senocak M, et al. Prevalence of epilepsy in Silivri, a rural area of Turkey. *Epilepsia* 1999; 40: 637-42.
- Beran RG, Hall L, Pesch A, et al. Population prevalence of epilepsy in Sydney, Australia. *Neuroepidemiology* 1982; 1: 201-8.
- Morgan CLI, Ahmed Z, Kerr MP. Social deprivation and prevalence of epilepsy and associated health usage. *J Neurol Neurosurg Psychiatr* 2000; 69: 13-7.
- Wallace H, Shorvon S, Tallis R. Age-specific incidence and prevalence rates of treated epilepsy in an unselected population of 2,052,922 and age-specific fertility rates of women with epilepsy. *Lancet* 1998; 352: 1970-3.

## APPENDIX 1

## Summary of studies on the prevalence of epilepsy by type of study and country.

## A. Community-based studies

Country	Year of study	Prevalence rates/1,000			Authors	Reference number
		Male	Female	Total		
<b>I. In developing countries</b>						
<b>Africa</b>						
- Igbo-Ora, Nigeria	1982	5.1	5.6	5.3	Osuntokun BO, et al	20
- Aiyete, Nigeria	*	28.0	44.0	37.0	Osuntokun BO, et al	21
- Rural area, Tanzania	1989	9.2	11.1	10.2	Rwiza HT, et al	11
- Zinvie, Africa	1997	*	*	33.5-35.1	Debrock C, et al	22
- Nakuru, Kenya	1985-86	*	*	18.2	Kaamugisha J, et al	23
<b>Latin America</b>						
- Republic of Panama	1988	*	*	57.0	Gracia F, et al	8
- Guatemala	*	*	*	8.5	Mendizabal JE, et al	9
- Bogota, Columbia	1974	15.5	22.9	19.5	Gomez JG, et al	10
- Ecuador	1986-87	*	*	12.2-19.5	Placencia M, et al	12
- Cordillera Province, Bolivia	1996	11.4	13.1	12.3	Nicoleeti A, et al	13
<b>Asia</b>						
- Republic of China	1983	5.0	4.1	4.6	Li SC, et al	7
- Kerala, South India	1996	5.2	4.6	4.9	Radhakrishnan K, et al	3
- Thailand, Nationwide	1991-92	*	*	2.8	Chuprapawan C	4
- Haryana, North India	1992-94	4.9	3.3	4.2	Singh A, et al	19
<b>II. In developed countries</b>						
- Sydney, Australia	1980	*	*	7.5	Beran RG, et al	25
- Copiah County, Mississippi	1978	12.4	8.6	10.4	Haerer AF, et al	14
- Silivri, Turkey	1994	10.4	10.0	10.2	Keraagac N, et al	24
- Riposto, Italy	1987	2.6	2.8	2.7	Reggio A, et al	15

## B. Hospital-based studies

Country	Year of study	Prevalence rates/1,000			Authors	Reference number
		Male	Female	Total		
Bradford, England	1996-98	*	*	7.3	Wright J, et al	18
South Glamorgan, UK	1996	*	*	2.0-13.4	Morgan CLI	26
Rural Iceland	1993	4.6	9.4	7.3	Olafsson E, et al	17
Tonbridge, UK	1983	15.4	26.5	20.3	Cockerell OC, et al	16
England and Wales	1995	*	*	5.15	Wallace H, et al	27

\* = Not mentioned

## ความชุกของโรคลมชักในชนบทของประเทศไทย : การศึกษาบนพื้นฐานของชุมชน

ธนิษฐ์ อัครวิเชียรจินดา, พ.บ.\*,

จิตร ลิทธิธรรม, พ.บ.\*\* , วชิร ธิญญานนท์, วท.ม.\*\*

การศึกษาโรคที่เป็นปัญหาในชุมชนจำเป็นต้องทราบถึงความชุกของโรคนั้นเพื่อที่จะวางแผนในการหามาตรการในการควบคุม

**วัตถุประสงค์ :** เพื่อหาความชุกของโรคลมชักในชนบทในประเทศไทย

**ผู้ป่วยและวิธีการ :** การศึกษาเริ่มตั้งแต่เดือนมกราคมถึงมิถุนายน 2543 โดยชาวบ้านทั้งหมดที่อาศัยอยู่ในตำบลท่าลาดขาว อำเภอโซ่พิสัย จังหวัดนครราชสีมาได้รับเชิญให้เข้ารับการตรวจโดยประสาทแพทย์ที่หมู่บ้าน

**ผล :** ชาวบ้าน 2069 ราย ใน 533 ครัวเรือน (คิดเป็นร้อยละ 72.2 ของครัวเรือนทั้งหมด) ถูกสัมภาษณ์และตรวจโดยประสาทแพทย์ 43 รายมีประวัติเคยชักมาก่อน มีเพียง 15 รายที่เป็นโรคลมชักซึ่งได้ค่าความชุกประมาณ 7.2/1,000 ประชากร ความชุกสูงสุด อยู่ในช่วงอายุ 5-9 ปี (17/1,000 ประชากร) และ 25-34 ปี (17.4/1,000 ประชากร)

**สรุป :** ความชุกของโรคลมชักในชนบทค่อนข้างต่ำกว่าความเป็นจริง แต่ก็ข้อมูลที่มีอยู่ในปัจจุบันนี้เท่านั้น

**คำสำคัญ :** โรคลมชัก, ระบาดวิทยา, การศึกษาบนพื้นฐานของชุมชน, ความชุก, ประเทศไทย

ธนิษฐ์ อัครวิเชียรจินดา, จิตร ลิทธิธรรม, วชิร ธิญญานนท์

จดหมายเหตุทางแพทย์ ๙ 2545; 85: 1066-1073

\* หน่วยประสาทวิทยา, กลุ่มงานอายุรกรรม, โรงพยาบาลมหาราชนครราชสีมา, นครราชสีมา 30000

\*\* ภาควิชาอายุรศาสตร์, คณะแพทยศาสตร์ และ คณะวิทยาลัยการสาธารณสุข, จุฬาลงกรณ์มหาวิทยาลัย, กรุงเทพฯ ๙ 10330

\*\*\* วิทยาลัยการสาธารณสุข, จุฬาลงกรณ์มหาวิทยาลัย, กรุงเทพฯ ๙ 10330

---

## Compliance with Treatment of Adult Epileptics in a Rural District of Thailand

---

THANIN ASAWAVICHJENJINDA, MD\*,  
CHITR SITTHI-AMORN, MD, PhD\*\*,  
WACHARIN TANYANONT, MS\*\*

### Abstract

**Background :** Epilepsy, a disease when seizures can occur from antiepileptic drug withdrawal, requires regular drug taking. Non-compliance, therefore, is a major factor contributing to sub-optimal control of the seizures.

**Purpose :** To determine the factors associated with noncompliance in epileptics in rural Thailand.

**Method :** All epileptics, registered in the Pak Thong Chai District and their caregivers were invited to be interviewed and examined by a neurologist in their village.

**Results :** Of a total of 93 epileptics registered, 83 with their caregivers were interviewed and examined by the neurologist (T.A.) and of those 72 were adults. Of the 72 adult epileptics, 41 (56.9%) were 100 per cent compliant and factors found to be significantly associated with compliance were gender, household income and patient's health insurance ( $p$ -value  $< 0.05$ ). The major reasons for non-compliance were misunderstanding (48.4%), forgetfulness (16.1%) and economic problem (12.9%).

**Conclusion :** To improve patient-compliance, the real factors for non-compliance, which are unique to patients in a specific area, need to be identified.

**Key word :** Epilepsy, Compliance, Adult, Reason for Non-Compliance, Thailand

ASAWAVICHJENJINDA T,  
SITTHI-AMORN C, TANYANONT W  
J Med Assoc Thai 2003; 86: 46-51

\* Unit of Neurology, Department of Medicine, Maharat Nakhon Ratchasima Hospital, Nakhon Ratchasima 30000,

\*\* The College of Public Health, Chulalongkorn University, Bangkok 10330, Thailand.

Epilepsy is one of the commonest problems encountered in neurological clinics in developing countries<sup>(1)</sup> and needs long-term treatment<sup>(2)</sup>. Compliance, including attendance at appointments and adherence to antiepileptic drug regimens (AED) is very important in the management of epilepsy<sup>(3)</sup>. The reason being that neglect of medical treatment may cause epileptic seizures<sup>(4)</sup>, which, in turn, may result in increased direct medical costs<sup>(5)</sup>, physical injury to self and others<sup>(5)</sup> and a decrease in the quality of life<sup>(6)</sup>. The frequency of noncompliance with treatment in epileptics has been reported ranging from 19 per cent to 75 per cent<sup>(7-9)</sup>.

The problem of compliance with treatment in chronic diseases is not selective but rather is pervasive and affects a broad range of illnesses<sup>(10)</sup>. However, compliance in epilepsy has not been studied extensively<sup>(4)</sup>.

The objectives of this study were to determine the frequency of non-compliance and the factors associated with non-compliance in epileptics in a rural district of Thailand.

## MATERIAL AND METHOD

The study was conducted in Pak Thong Chai, one of the 26 districts of Nakhon Ratchasima Province, the principal province of Northeastern Thailand. The district is around 32 kilometers Southwest of the provincial capital and 286 kilometers from Bangkok. It covers 1,352 square kilometers with a population (in 1997) of 74,177 and is divided into 16 sub-districts with a total of 20 Sub-District Health Care Offices (SDHCO) and one Community Hospital.

In the district, like all districts in Nakhon Ratchasima Province, there is a Registry of Epileptics who have visited either the SDHCO or the Community Hospital; the Registry is updated yearly by sub-district health care officials. In 1997, 93 epileptics were registered in the district. All of the patients with their caregivers were informed of the study and invited, after informed consent, for an interview and examination by sub-district health care officials and a neurologist (T.A) from January-May 1998.

The data collected included: age; gender; socio-economic data; whether the patient had health care insurance which would entitle them to free health care including AEDs.

The data related to epilepsy included: age at first onset of epilepsy; duration of epilepsy and of treatment; frequency of seizures; the AED regimens;

alcoholic intake and the need to be escorted for treatment.

The outcome variable measured was whether the level of compliance with AED regimens over the past year was 100 per cent (defined as taking the AED on time, without fail and without manipulating their dosage)<sup>(5)</sup>, as determined during an in-depth interview of the patient and their caregiver. During the same interview, the major reason for non-compliance was determined.

The authors defined Epilepsy as two or more clinical afebrile seizures unrelated to acute metabolic derangements or to withdrawal from alcohol or drugs<sup>(11)</sup>. Seizures, which occur within a 24-hour period, were excluded<sup>(11)</sup>.

In the data analysis, the sample based upon the interview of the epileptic and of their caregivers, each group was further stratified into those considered compliance and non-compliance. The differences were, then, analyzed by univariate and multivariate analysis.

## RESULTS

Of the 93 epileptics registered in Pak Thong Chai District, only 83 (89.2%) and their caregivers were interviewed and examined. The gender and age response rate is shown in Table 1. The non-respondents of ten epileptics, although it could not be verified, were presumably less affected by their epilepsy.

Of the 83 epileptics examined, only 11 were children (age 0-14 years); so, further analyses was restricted to the adults. Of the 72 adult epileptics, the average age was 36.3 years (SD = 14.0; median = 36; range: 16-75 years) and 59.7 per cent were female. The patient's characteristics and clinical characteristics are shown in Table 2.

The patient-compliance (100%) over the past one-year was 56.9 per cent. The differences between those considered to be compliant or non-compliant by each of the demographic, socio-economic and clinical variables are shown in Table 2.

By univariate analysis, none of the analyzed variables were significantly associated with compliance. However, by multivariate analysis, gender, household income and patient's health insurance had a significant association with compliance (shown in Table 2).

The major reasons for non-compliance were misunderstanding of the need for long-term treatment (48.4%); forgetfulness (16.1%) and economic pro-

blem (12.8%). The other reasons were continuity of seizures despite treatment (9.7%); misbeliefs (6.5%) and not having a caregiver to escort them to hospital (6.5%).

## DISCUSSION

In this study, compliance with AED regimens over the past one year was measured by self-report of the patients or their caregivers. The data collection method and the cross-sectional design employed precluded the possibility of any pill counting procedures or blood level testing. Although, it was quite difficult for them to recall the degree of compliance with their physician's prescriptions, the authors used 100 per cent compliance rates; even though compliance of 80 per cent is sufficient for optimal treatment in patients with epilepsy<sup>(12)</sup>. The frequency of the patients with 100 per cent compliance in the present study was only 56.9 per cent, which is similar to reports from other studies<sup>(6,13)</sup>.

On multivariate analysis, gender, household income and patient's health insurance had a statistically significant relationship to compliance. The possible reasons for this finding is that, in Thailand any patient who holds any kind of health insurance card can get medical service including drugs free of charge at any level of government health care offices. In addition, most poor patients trust their physicians and strictly follow the physicians' suggestions. Economic status is still a major factor for patient compliance<sup>(10)</sup>. Male gender is another variable because male patients usually work outside.

Other factors can affect compliance. However, the relationship of these factors to compliance has not been consistent. Age and number of AED

taken had a statistically significant relationship with compliance in Buck's study<sup>(14)</sup> but were not in Gopinath's study<sup>(5)</sup>. Seizure frequency and duration of epilepsy were related to compliance in three studies (Kyngas<sup>(4)</sup>, Gopinath<sup>(5)</sup> and Loiseau<sup>(15)</sup>) but were not in Buck's study<sup>(14)</sup>. However, gender and seizure type have consistently been shown to have no relationship with compliance<sup>(5,14,15)</sup>. The patient's education in Gopinath's study<sup>(5)</sup> and of age at first onset of epilepsy in Loiseau's study<sup>(15)</sup> showed no significant relationship to compliance. Alcoholic intake in Kyngas's study<sup>(4)</sup> and of higher income in Shope's study<sup>(16)</sup> revealed a significant association with compliance.

Not only the factors mentioned earlier but also several personal reasons may contribute to non-compliance including: patient's life style<sup>(14,17)</sup>; patient's feeling of stigma<sup>(14)</sup>; denial of their illness<sup>(18)</sup>; individual's perception of causality<sup>(5)</sup>; relationship between patient and health care staff<sup>(4,5,14)</sup>; misconceptions regarding the goal of the treatment and the consequences of missing a prescribed drug dose<sup>(4)</sup>; misunderstanding the doctor's instructions<sup>(18)</sup>, fear of addiction<sup>(19)</sup>; general dissatisfaction with treatment<sup>(5,19)</sup>; trouble with remembering<sup>(18)</sup> and forgetfulness<sup>(18)</sup>. Some studies have found that the following altered drug taking behavior: adverse side effects<sup>(4,18,19)</sup>; patient's personal experience<sup>(14)</sup>; family relationships<sup>(3)</sup>; the support of parents<sup>(4)</sup>.

Finally, no single measure represents a completely valid indicator of the risk of non-compliance. Mitchell *et al* stated in their paper that dealing with compliance in children that "It is multidimensional and includes related but somewhat distinct facets of socio-cultural, medical, behavioral, environmental

Table 1. Gender and age specific response rates.

Age group (year)	Males		Females		Both genders	
	n/N	%	n/N	%	n/N	%
0-4	1/1	100.0	0/0	-	1/1	100.0
5-9	2/2	100.0	2/2	100.0	4/4	100.0
10-14	3/3	100.0	3/3	100.0	6/6	100.0
15-24	9/11	81.8	7/7	100.0	16/18	88.9
25-34	7/10	70.0	10/11	90.9	17/21	80.9
35-44	6/10	60.0	18/18	100.0	24/28	85.7
45-54	5/5	100.0	2/2	100.0	7/7	100.0
55-64	1/1	100.0	4/4	100.0	5/5	100.0
65+	1/1	100.0	2/2	100.0	3/3	100.0
All age groups	35/44	79.5	48/49	97.9	83/93	89.2

n = Respondents, N = Registered epileptics

**Table 2. Factors and their relationship with patient-compliance in adult epileptics.**

Factors	C (41)	NC (31)	Total		P-value (Univariate)	P-value (Multivariate)
			n	%		
Gender						
Male	16	13	29	40.3	0.99	< 0.05
Female	25	18	43	59.7		
Patient's education						
≤ 6 yrs	36	28	64	88.9	1.00	0.68
> 6 yrs	5	3	8	11.1		
Caregiver's education						
≤ 6 yrs	39	31	70	97.2	0.50	0.89
> 6 yrs	2	0	2	2.8		
Patient's employment						
Unemployed	19	13	32	44.4	0.89	0.91
Employed	22	18	40	55.6		
Household income (monthly) in Baht						
≤ 2,800	28	15	43	59.7	0.14	< 0.05
> 2,800	13	16	29	40.3		
Alcohol intake						
Regular or occasional	8	10	18	25.0	0.34	0.49
Not at all or quit	33	21	54	75.0		
Need to be escorted						
Yes	16	14	30	41.7	0.78	0.23
No	25	17	42	58.3		
AED regimens						
Monotherapy	28	24	52	72.2	0.55	0.45
Multiple AED regimens	13	7	20	27.8		
Patient's health insurance						
Yes (one or more cards)	36	21	57	79.2	0.07	< 0.05
No	5	10	15	20.8		
Age at first onset of epilepsy						
< 20 yrs	25	18	43	59.7	0.99	0.66
≥ 20 yrs	16	13	29	40.3		
Duration of epilepsy						
≤ 6 yrs	5	2	7	9.7	0.69	0.77
> 6 yrs	36	29	65	90.3		
Duration of treatment						
≤ 6 yrs	5	7	12	16.7	0.39	0.79
> 6 yrs	36	24	60	83.3		
Seizure frequency						
< once a week	36	24	60	83.3	0.39	0.06
≥ once a week	5	7	12	16.7		

C = Compliance, NC = Non-compliance

(i.e. family), and individual characteristics"<sup>(10)</sup>. In the present study, the reasons for non-compliance with AED regimens; misunderstanding, forgetfulness and economic problem were compatible with the results reported by other studies<sup>(15,16,18)</sup>. This suggests how patient-compliance can be improved.

#### ACKNOWLEDGEMENT

The authors wish to thank the Health System Research Institute for funding the study. We

also wish to thank the Director of Maharat Nakhon Ratchasima Hospital, the Provincial Chief Medical Officer Nakhon Ratchasima, the Head of the Department of Medicine, the Head of the Department of Social Health and the staff, the Director of Pak Thong Chai Community Hospital, the Head and all of the Sub-District Health Officials of the Pak Thong Chai District Health Office who assisted us during this study.

## REFERENCES

1. Osuntokun BO, Adeuja AOG, Nottidge VA, et al. Prevalence of the epilepsies in Nigerian Africans: A community-based study. *Epilepsia* 1987; 28: 272-9.
2. Medical Research Council Antiepileptic Drug Withdrawal Study Group. Randomized study of antiepileptic drug withdrawal in patients in remission. *Lancet* 1991; 337: 1175-80.
3. Otero S, Hodes M. Maternal expressed emotion and treatment compliance of children with epilepsy. *Dev Med Child Neurol* 2000; 42: 604-8.
4. Kyngas H. Compliance with health regimens of adolescents with epilepsy. *Seizure* 2000; 9: 598-604.
5. Gopinath B, Radhakrishnan K, Sarma PS, Jayachandran D, Alexander A. A questionnaire survey about doctor-patient communication, compliance and locus of control among South Indian people with epilepsy. *Epilepsy Res* 2000; 39: 73-82.
6. Leppik IE. How to get patients with epilepsy to take their medication: The problem of non-compliance. *Postgrad Med J* 1990; 88: 253-6.
7. Shope JT. Medication compliance. *Pediatr Clin North Am* 1981; 28: 5-8.
8. Wright EC. Non-compliance-or how many aunts has Matilda? *Lancet* 1993; 342: 909-13.
9. Cramer JA, Mattson RH, Prevey ML, Scheyer RD, Ouellette VL. How often is medication taken as prescribed? A novel assessment technique. *JAMA* 1989; 261: 3273-7.
10. Mitchell WG, Scheier LM, Baker SA. Adherence to treatment in children with epilepsy: Who follow "doctor's orders"? *Epilepsia* 2000; 41: 1616-25.
11. Waaler PE, Blom BH, Skeidsvoll H, Mykletun A. Prevalence, classification, and severity of epilepsy in children in Western Norway. *Epilepsia* 2000; 41: 802-10.
12. Lisk DR, Greene SH. Drug compliance and seizure control in epileptic children. *Postgrad Med J* 1985; 61: 401-5.
13. Leppik IE, Schmidt D. Summary of the first international workshop on compliance in epilepsy. *Epilepsy Res* 1988; 1: 179-82.
14. Buck D, Jacoby A, Baker GA, Chadwick DW. Factors influencing compliance with antiepileptic drug regimens. *Seizure* 1997; 6: 87-93.
15. Loiseau P, Marchal C. Determinants of compliance in epileptic patients. In: Schmidt D, Leppik IE, eds. *Compliance in epilepsy*. Amsterdam: Elsevier Science Publishers B.V, 1988: 135-40.
16. Shope JT. Compliance in children and adults: Review of studies. In: Schmidt D, Leppik IE, eds. *Compliance in epilepsy*. Amsterdam: Elsevier Science Publishers B.V, 1988: 23-47.
17. Leppik IE. Compliance during treatment of epilepsy. *Epilepsia* 1988; 29: 79-84.
18. Garnett WR. Antiepileptic drug treatment: Outcomes and adherence. *Pharmacotherapy* 2000; 20 (Suppl 2): S191-9.
19. Scambler G, Hopkins A. Accommodating epilepsy in families. In: Anderson R and Bury M, eds. *Living with chronic illness: The experience of patients and their families*. London: Allen & Unwin, 1988: 156-76.

## การรับประทานยาตามแพทย์สั่งของผู้ป่วยโรคลมชักในชุมชน

ธนินทร์ อัครวิเชียรจินดา, พบ\*,  
จิตร ลิทธิอมร, พบ, ปรด\*\*, วชิริน ธัญญานนท์, พม\*\*

โรคลมชักเป็นโรคชนิดหนึ่งที่อาการชักจะเกิดขึ้นได้จากการสั่นประทุกันชัก การรับประทานยาตามแพทย์สั่งจึงมีความจำเป็น

**วัตถุประสงค์ :** เพื่อหาปัจจัยที่มีส่วนเกี่ยวข้องกับการไม่รับประทานยาตามแพทย์สั่งของผู้ป่วยโรคลมชักในชุมชน

**ผู้ป่วยและวิธีการ :** โดยการเชิญญาติและผู้ป่วยโรคลมชักที่ได้ขึ้นทะเบียนที่อำเภอปรางค์กู่ จังหวัดนครราชสีมาเพื่อที่จะสัมภาษณ์และตรวจโดยประสาทแพทย์ที่หมู่บ้านของผู้ป่วย

**ผล :** ผู้ป่วย 83 รายจากทั้งหมด 93 รายที่ได้ขึ้นทะเบียนที่อำเภอปรางค์กู่และญาติได้รับการสัมภาษณ์และตรวจโดยประสาทแพทย์ ในจำนวน 83 รายนี้ 72 รายเป็นผู้ป่วยที่มีอายุตั้งแต่ 15 ปีขึ้นไป และมีเพียง 41 ราย (ร้อยละ 56.9) ที่รับประทานยาตามแพทย์สั่ง ปัจจัยที่มีส่วนเกี่ยวข้องกับการรับประทานยาของผู้ป่วยได้แก่ เพศ รายได้ของครอบครัว และการมีบุตรสุขภาพ ส่วนเหตุผลหลักที่ผู้ป่วยไม่รับประทานยาได้แก่ การขาดความรู้ความเข้าใจในเรื่องการรักษา (48.4%) การลืมรับประทานยา (16.1%) และปัญหาทางการเงิน (12.9%)

**สรุป :** การที่จะทำให้ผู้ป่วยรับประทานยาตามแพทย์สั่งเพื่อผลการรักษา จำเป็นที่จะต้องทราบถึงเหตุผลที่แท้จริงของผู้ป่วยซึ่งจะแตกต่างกันไปตามพื้นที่

**คำสำคัญ :** โรคลมชัก, การรับประทานยาตามแพทย์สั่ง, ผู้ป่วยที่อายุตั้งแต่ 15 ปีขึ้นไป, เหตุผลของการไม่รับประทานยา, ประเทศไทย

ธนินทร์ อัครวิเชียรจินดา, จิตร ลิทธิอมร, วชิริน ธัญญานนท์

จดหมายเหตุมหาวิทยาลัย ๙ 2546; 86: 46-51

\* หน่วยประสาทวิทยา, กลุ่มงานอายุรกรรม, โรงพยาบาลมหาราชนครราชสีมา, นครราชสีมา 30000

\*\* วิทยาลัยการสาธารณสุข, จุฬาลงกรณ์มหาวิทยาลัย, กรุงเทพฯ ๙ 10330

# Association of psychosocial risk factors with risk of acute myocardial infarction in 11 119 cases and 13 648 controls from 52 countries (the INTERHEART study): case-control study

Annika Rosengren, Steven Hawken, Stephanie Öunpuu, Karen Sliwa, Mohammad Zubaid, Wael A Almahmeed, Kathleen Ngu Blackett, Chitr Sitthi-amorn, Hiroshi Sato, Salim Yusuf, for the INTERHEART investigators\*

## Summary

**Background** Psychosocial factors have been reported to be independently associated with coronary heart disease. However, previous studies have been in mainly North American or European populations. The aim of the present analysis was to investigate the relation of psychosocial factors to risk of myocardial infarction in 24 767 people from 52 countries.

**Methods** We used a case-control design with 11 119 patients with a first myocardial infarction and 13 648 age-matched (up to 5 years older or younger) and sex-matched controls from 262 centres in Asia, Europe, the Middle East, Africa, Australia, and North and South America. Data for demographic factors, education, income, and cardiovascular risk factors were obtained by standardised approaches. Psychosocial stress was assessed by four simple questions about stress at work and at home, financial stress, and major life events in the past year. Additional questions assessed locus of control and presence of depression.

**Findings** People with myocardial infarction (cases) reported higher prevalence of all four stress factors ( $p < 0.0001$ ). Of those cases still working, 23.0% ( $n=1249$ ) experienced several periods of work stress compared with 17.9% (1324) of controls, and 10.0% (540) experienced permanent work stress during the previous year versus 5.0% (372) of controls. Odds ratios were 1.38 (99% CI 1.19–1.61) for several periods of work stress and 2.14 (1.73–2.64) for permanent stress at work, adjusted for age, sex, geographic region, and smoking. 11.6% (1288) of cases had several periods of stress at home compared with 8.6% (1179) of controls (odds ratio 1.52 [99% CI 1.34–1.72]), and 3.5% (384) of cases reported permanent stress at home versus 1.9% (253) of controls (2.12 [1.68–2.65]). General stress (work, home, or both) was associated with an odds ratio of 1.45 (99% CI 1.30–1.61) for several periods and 2.17 (1.84–2.55) for permanent stress. Severe financial stress was more typical in cases than controls (14.6% [1622] vs 12.2% [1659]; odds ratio 1.33 [99% CI 1.19–1.48]). Stressful life events in the past year were also more frequent in cases than controls (16.1% [1790] vs 13.0% [1771]; 1.48 [1.33–1.64]), as was depression (24.0% [2673] vs 17.6% [2404]; odds ratio 1.55 [1.42–1.69]). These differences were consistent across regions, in different ethnic groups, and in men and women.

**Interpretation** Presence of psychosocial stressors is associated with increased risk of acute myocardial infarction, suggesting that approaches aimed at modifying these factors should be developed.

## Introduction

Popular opinion holds that stress is an important risk factor for coronary heart disease. However, compared with other major risk factors, psychosocial variables such as stress are difficult to define objectively, and stress consists of several different (and inter-related) elements. Therefore, measurement of stress is complex and difficult. Despite this drawback, several constructs within the broad conceptual framework of stress are increasingly regarded as being causally related to coronary heart disease.<sup>1–15</sup>

The concept of stress encompasses several factors, from external stressors such as job stress,<sup>5,8,10,13</sup> adverse life events<sup>7</sup> and financial problems, to potential reactions to stress such as depression,<sup>1–3</sup> vital exhaustion,<sup>15</sup> anxiety,<sup>12,14</sup> psychological distress,<sup>11</sup> and sleeping difficulties. The same construct might not be applicable

in different countries and ethnic groups, because cultural influences can vary. Perceived mental stress, measured by response to a single-item question, was associated with increased mortality from coronary disease in a large study of Japanese men and women.<sup>9</sup> Apart from this study,<sup>9</sup> previous investigations have been done in mainly North American or European populations. Thus, limited data are available about psychological variables and coronary heart disease in other countries and ethnic groups. The aim of the INTERHEART study, undertaken in a large number of patients with a first acute myocardial infarction and controls matched for age and sex, was to investigate the associations of several psychosocial stressors with the risk of acute myocardial infarction globally, and in different populations characterised by age, sex, geographic region, and ethnic origin.



Lancet 2004; 364: 953–62

Published online  
September 3, 2004  
<http://image.thelancet.com/extras/04art8002web.pdf>

See [Comment](#) page 912

\*See [Articles](#) page 937

McMaster University, Hamilton Health Sciences, Hamilton, ON, Canada (Prof S Yusuf FRCP, S Hawken MSc, S Öunpuu PhD); Sahlgrenska University Hospital/Ostra, Göteborg, Sweden (Prof A Rosengren MD); Baragwanath Hospital, Johannesburg, South Africa (Prof K Sliwa MD); Kuwait University, Kuwait City, Kuwait (M Zubaid MD); CHU, Yaoundé, Cameroon (Prof K N Blackett MD); Al-Jazeera and Central Hospitals, Abu Dhabi, United Arab Emirates (W A Almahmeed MD); Department of Medicine, Chulalongkorn University, Bangkok, Thailand (Prof C Sitthi-amorn MD); and Osaka University Graduate School of Medicine, Suita, Japan (H Sato MD)

Correspondence to:  
Prof Salim Yusuf, Population Health Research Institute, 237 Barton Street East, Hamilton, ON, Canada L8L 2X2  
[yusufs@mcmaster.ca](mailto:yusufs@mcmaster.ca)

## Patients and methods

### Study population

12 461 incident cases of acute myocardial infarction from 262 centres in 52 countries representing all geographic regions, and 14 637 age-matched, sex-matched, and site-matched controls free of clinical heart disease, took part in the study.<sup>16,17</sup> Centres attempted to recruit consecutive patients. Recruitment was from February, 1999, until March, 2003. Patients admitted to the coronary care unit or equivalent cardiology ward of participating centres were screened to identify incident cases of acute myocardial infarction and enrolled within 24 h. Details of criteria used for the definition of acute myocardial infarction are described in the accompanying paper.<sup>16</sup> At least one control was recruited and matched to every case of acute myocardial infarction by age (up to 5 years older or younger) and sex. Eligible control sources were community-based (visitor or relative of a patient from a non-cardiac ward, or an unrelated visitor of a cardiac patient) or hospital-based.<sup>16</sup>

INTERHEART was approved by appropriate regulatory and ethics committees in all participating countries and centres. All participants provided informed consent before taking part in the study.

### Procedures

We obtained data for demographic factors (country of origin, first language), socioeconomic status (education, occupation, income), lifestyle (tobacco use, physical activity, dietary patterns), personal and family patterns of cardiovascular disease, and risk factors.<sup>16</sup> Staff were trained in study procedures with standard manuals, videotapes, and instructions at meetings or at site visits. Trained staff administered the questionnaire before patients left the hospital. A standard yet simple set of questions that inquired about psychosocial conditions during the previous 12 months was included in the interview. Complete data on psychosocial variables were available for 11 119 cases and 13 648 controls.

We assessed psychological stress with two single-item questions relating to stress at work and home. Stress was defined as feeling irritable, filled with anxiety, or as having sleeping difficulties as a result of conditions at work or at home. Patients were specifically asked to respond about their condition before their acute myocardial infarction. For every question, we asked participants to report how often they had felt stress, using the following response options: 1) never; 2) some periods; 3) several periods; or 4) permanent stress. These two questions are an adaptation of a single question that has been used in multiple studies in Göteborg, Sweden since 1970. For example, in a prospective study of Swedish men,<sup>14</sup> permanent stress, according to this question, was associated with an increased risk of acute myocardial infarction, stroke, and death. Because stress at work and at home were highly intercorrelated, and because only 48·8% (5426) of cases and 54·1% (7387) of

controls were currently working, we created a general stress scale that combined stress at work, home, or both and was graded as follows: 1) never experienced stress; 2) experienced some periods at home or at work; 3) experienced several periods at home or at work; 4) experienced permanent stress at home or at work.

We defined level of financial stress as: 1) little or none; 2) moderate; or 3) high or severe. Occurrence of major adverse life events was documented by asking participants whether they had experienced any specified life events in the past year—marital separation or divorce, loss of job or retirement, loss of crop or business failure, violence, major intrafamily conflict, major personal injury or illness, death or major illness of a close family member, death of a spouse, or other major stress.

Generalised locus of control—that is, the perceived ability to control life circumstances—was determined by responses to a questionnaire containing six scale items that have been used extensively in studies in eastern Europe.<sup>18</sup> Responses from controls were used to provide a summary score that was divided into quartiles, of which the first quartile represented the lowest and the fourth quartile the highest score.

We assessed depression by asking whether, during the past 12 months, the participant had felt sad, blue, or depressed for 2 weeks or more in a row, and if yes, graded by a set of seven no-yes questions—lose interest in things, feel tired or low on energy, gain or lose weight, trouble falling asleep, trouble concentrating, think of death, feeling worthless—of which five or more positive responses were defined as clinical depression. This questionnaire is an adaptation of the short form DSM-IV CIDI questionnaire for depression.<sup>19</sup>


Standard physical measurements were done in duplicate, by the same examiner, on every participant: height, weight, and waist and hip circumference. Waist and hip circumferences were measured with a non-stretchable standard tape measure. We recorded relevant items, including history of tobacco use, diabetes, family history, physical activity, and patterns of alcohol and food consumption. In 292 controls, we readministered the questionnaire after a median interval of 409 days. The weighted  $\kappa$  statistic was 0·53 for global stress, 0·66 for financial stress, 0·43 for life events, 0·60 for locus of control, and 0·44 for depression.

### Statistical analysis

Details of statistical analysis are provided in the accompanying paper.<sup>16</sup> Briefly, we accounted for the potential differences in age structure of the populations (subdivided by region or ethnic origin) by direct standardisation of the frequencies to the overall INTERHEART age distribution, using a five-level age stratification factor (<45, 45–55, 56–65, 66–70, >70).<sup>20</sup> We calculated means and medians to summarise continuous effects and compared them with *t* tests, or

See [Articles](#) page 937

See [Articles](#) page 937




**Figure 1: Moderate or severe general stress in cases and controls by geographic region and ethnic origin**

Percentages and controls are age adjusted. Odds ratios are adjusted for geographic region, age, sex, and smoking.

appropriate non-parametric tests when distributional assumptions were in doubt. When testing for associations between stress subgroups, linear regression was used for continuous variables and the Cochran-Armitage trend test for frequencies. The findings presented are for models fitted with unconditional logistic regression, adjusted for age, sex, geographic region, and potential confounders.

Relative risk estimates are reported as odds ratios and accompanying 99% CIs. We produced statistical analyses

and graphics with SAS version 8.2 (SAS, Cary, NC, USA) and S-Plus version 6 (Insightful, Seattle, WA, USA). All statistical tests of hypotheses are two-sided. Population attributable risks (PARs)—ie, the proportion of all cases attributable to the relevant factor if causality were proven—and 99% CIs were calculated for various risk factors in the study, using the methods of Benichou and Gail.<sup>21</sup> The PARs presented are adjusted for confounders in a similar fashion to the corresponding logistic regression models for odds ratio estimates. PAR estimates



**Figure 2: Depression in cases and controls by geographic region and ethnic origin**

Percentages of cases and controls are age adjusted. Odds ratios are adjusted for geographic region, age, sex, and smoking.

were calculated with the Interactive Risk Attributable Program software (US National Cancer Institute, 2002).<sup>22</sup>

#### Role of the funding source

The sponsor of the study had no role in study design, data collection, data analysis, data interpretation, or writing of the report. The corresponding author had full access to all data and had final responsibility for the decision to submit for publication.

#### Results

The mean age of cases was 58.2 years (SD 12.2) and of controls 57.1 years (12.1); 24.2% (2686) of cases and 26.5% (3619) of controls were women. Figure 1 outlines the distribution of cases and controls by region and ethnic origin and provides the odds ratios associated with high general stress. We defined high general stress as several periods of or permanent stress, at work, home, or both. Figure 2 shows corresponding data for

depressive mood, which we defined as feeling sad, blue, or depressed for 2 weeks or more in a row during the past 12 months.

Table 1 presents analyses among controls by general stress score. Individuals with high general stress were younger, heavier, and more usually smokers, had a slightly lower waist/hip ratio, and were less likely to have low income and low education. Strong associations were noted between perceived general stress and financial stress, having had two or more life events, depression, and low control. Inter-relations between the different psychosocial variables were very similar among cases and controls.

Half the study population (5426 cases [48.8%] and 7387 controls [54.1%]) were working outside the home. Higher ratings for work stress were more usual in cases than controls (table 2). For example, during the previous year, just over a third more cases experienced several periods of work stress compared with controls (odds ratio 1.38 [99% CI 1.19–1.61]), and permanent work stress was experienced by twice as many cases than controls (2.14 [1.73–2.64]), after adjustment for age, sex, geographic region, and smoking. The PAR among those working was 9% (99% CI 1–18). Further adjustment for education and income, hypertension, diabetes, level of physical activity, waist/hip ratio, dietary patterns, alcohol, or raised plasma lipids did not alter these results to a significant degree. Therefore, we report results from the most parsimonious model.

Compared with controls, cases also reported more frequent periods of stress at home during the previous 12 months (table 2). For example, more cases than controls experienced several periods of stress (odds ratio 1.52 [1.34–1.72]), whereas twice as many cases reported permanent stress at home compared with controls (2.12 [1.68–2.65]). The PAR for stress at home was 8% (4–12). Reports of severe financial stress were also more typical in cases than controls (adjusted odds ratio 1.33 [1.19–1.48]; table 2). The PAR for financial stress was 11% (7–14). Two or more stressful life events were reported by about a half more cases compared with controls (odds ratio 1.48 [1.33–1.64]; PAR 10% [8–13]; table 2). Of these events, business failure (1.60 [1.42–1.80]) and major intra-family conflict (1.55 [1.41–1.70]) had the highest risks, whereas job loss (1.36 [1.22–1.52]), death of spouse (1.37 [1.11–1.69]) and violence (1.31 [1.13–1.53]), although still significant, were associated with lower risk. Divorce, injury, and death of other family members were similar among cases and controls.

High locus of control was a significant protective factor. After adjustment for age, sex, geographic region and smoking, the odds ratio was 0.72 (99% CI 0.65–0.79) for the second highest quartile of the population relative to those in the lowest quartile for locus of control, and 0.68 (0.61–0.76) for the highest

	Never (n=3688)	Some (n=7193)	Several (n=2183)	Permanent (n=584)	p*
<b>Risk factor</b>					
Age (years)	60.7 (12.01)	56.7 (12.0)	53.6 (11.4)	52.5 (10.6)	<0.0001
Body-mass index (kg/m <sup>2</sup> )	25.7 (4.0)	25.8 (4.1)	26.0 (4.3)	26.6 (4.5)	<0.0001
Waist/hip ratio	0.918 (0.082)	0.906 (0.084)	0.906 (0.082)	0.912 (0.085)	<0.0001
Systolic blood pressure (mm Hg)	131 (18)	129 (17)	128 (17)	128 (18)	<0.0001
Serum cholesterol (mmol/L)	5.03 (1.31)	5.04 (1.22)	5.22 (1.22)	5.32 (1.17)	<0.0001
Serum triglyceride (mmol/L)	1.90 (1.29)	1.96 (1.35)	2.05 (1.65)	2.03 (1.32)	0.0003
Serum HDL (mmol/L)	1.08 (0.38)	1.07 (0.38)	1.08 (0.41)	1.11 (0.40)	0.67
ApoB/ApoA1 ratio	0.790 (0.374)	0.791 (0.326)	0.806 (0.327)	0.800 (0.243)	0.16
Smoking	25.6% (943)	27.2% (1952)	30.2% (658)	34.0% (197)	<0.0001
Sedentary leisure time physical activity	52.3% (1927)	55.1% (3963)	51.8% (1131)	49.9% (291)	0.48
Diabetes	7.9% (291)	7.0% (503)	7.2% (157)	7.2% (42)	0.28
Hypertension	24.0% (885)	21.7% (1559)	20.1% (438)	23.1% (135)	0.006
<b>Income</b>					
Lowest two-fifths	52.3% (1901)	49.2% (3494)	46.7% (1001)	44.1% (254)	<0.0001
<b>Education</b>					
Fewer than 8 years	44.5% (1639)	37.5% (2693)	30.5% (666)	26.7% (156)	
College or university	29.6% (1093)	36.2% (2605)	48.2% (1052)	49.5% (289)	<0.0001
<b>Other psychosocial variables</b>					
High or severe financial stress	4.8% (177)	11.6% (832)	21.7% (473)	30.3% (177)	<0.0001
Two or more events	7.2% (266)	10.7% (771)	25.0% (546)	32.2% (188)	<0.0001
Depression	10.1% (374)	15.7% (1132)	29.4% (642)	43.8% (256)	<0.0001
Low locus of control	16.0% (590)	18.7% (1343)	22.7% (496)	32.5% (190)	<0.0001

Data are mean (SD) or percentage of controls (n). Missing data for smoking in 46 controls, diabetes in nine, hypertension in eight, physical activity in eight, income in 192, and education in five. Spearman correlation coefficients (cases/controls) between global stress and financial stress 30.0/25.7, two or more life events 26.2/22.2, depression 22.7/19.5, locus of control -12.1/-5.6. \*Test for trend.

**Table 1: Cardiovascular risk factors, income, education, and other psychosocial variables in controls (both working and non-working) by general stress score**

relative to the lowest quartile (table 2). However, after full adjustment for all risk factors, the apparent effect was substantially attenuated, with an estimated odds ratio of 0.75 (0.65–0.86) in the fourth quartile relative to the first. The PAR for low locus of control was 16% (10–22).

More cases than controls reported feeling sad, blue, or depressed for more than 2 weeks or more in a row (odds ratio 1.55 [1.42–1.69]; table 2), and this difference did not change substantially after adjustment for other factors. No relation was reported between number of items with positive responses for the depression question and risk of acute myocardial infarction. The PAR associated with sadness and depression was 9% (7–10).

The general stress measure—combining work and home stress—was associated with an odds ratio of 1.45 (1.30–1.61) for several periods and 2.17 (1.84–2.55) for permanent stress (table 2). Adding financial stress, locus of control, life events, and reports of feeling sad, either as single elements or in combination with this scale, did not improve its discriminatory power. For general stress, the PAR was 12% (7–17); however, combining any exposure to general stress, financial stress, stressful life events, depression, and low locus of control, a PAR of 29% (22–35) was estimated after adjustment for age, sex, geographic region, and smoking. Further adjustment for all risk factors as above changed the PAR estimate to 33% (25–41).

	Number of cases (%)	Number of controls (%)	Odds ratio (99% CI)	PAR (99% CI)
<b>Stress at work (n=12 813)</b>				
Never	1138 (21.0%)	1768 (23.9%)	1	
Some of the time	2499 (46.1%)	3923 (53.1%)	0.95 (0.84–1.08)	
Several periods	1249 (23.0%)	1324 (17.9%)	1.38 (1.19–1.61)	
Permanent	540 (10.0%)	372 (5.0%)	2.14 (1.73–2.64)	9% (1–18)
<b>Stress at home (n=24 767)</b>				
Never	4086 (36.8%)	5343 (39.2%)	1	
Some of the time	5361 (48.2%)	6873 (50.4%)	1.05 (0.97–1.13)	
Several periods	1288 (11.6%)	1179 (8.6%)	1.52 (1.34–1.72)	
Permanent	384 (3.5%)	253 (1.9%)	2.12 (1.68–2.65)	8% (4–12)
<b>General stress* (n=24 767)</b>				
Never	2777 (25.0%)	3688 (27.0%)	1	
Some period, home or work	5352 (48.1%)	7193 (52.7%)	1.05 (0.96–1.14)	
Several periods, home or work	2139 (19.2%)	2183 (16.0%)	1.45 (1.30–1.61)	
Permanent, home or work	851 (7.7%)	584 (4.3%)	2.17 (1.84–2.55)	12% (7–17)
<b>Financial stress (n=24 767)</b>				
Little or none	4872 (43.8%)	6628 (48.6%)	1	
Moderate	4625 (41.6%)	5361 (39.3%)	1.19 (1.11–1.29)	
Severe	1622 (14.6%)	1659 (12.2%)	1.33 (1.19–1.48)	11% (7–14)
<b>Stressful life events (n=24 767)</b>				
None	6425 (57.8%)	8528 (62.5%)	1	
1	2904 (26.1%)	3349 (24.5%)	1.23 (1.13–1.34)	
2 or more	1790 (16.1%)	1771 (13.0%)	1.48 (1.33–1.64)	10% (8–13)
<b>Locus of control (n=24 767)</b>				
Q1	2620 (23.6%)	2619 (19.2%)	1	
Q2	2938 (26.4%)	3265 (23.9%)	0.89 (0.80–0.98)	
Q3	3614 (32.5%)	4839 (35.5%)	0.72 (0.65–0.79)	
Q4	1947 (17.5%)	2925 (21.4%)	0.68 (0.61–0.76)	16% (10–22)
<b>Feeling depressed† (n=24 767)</b>				
No	8446 (76.0%)	11244 (82.4%)	1	
Yes	2673 (24.0%)	2404 (17.6%)	1.55 (1.42–1.69)	9% (7–10)
<b>Depression (n=24 767)</b>				
Not depressed	8446 (76.0%)	11244 (82.4%)	1	
0–1 items	346 (3.1%)	298 (2.2%)	1.50 (1.21–1.86)	
2–4 items	1369 (12.3%)	1145 (8.4%)	1.65 (1.47–1.85)	
5 or more items	958 (8.6%)	961 (7.0%)	1.44 (1.27–1.65)	

All associations are significant at  $p < 0.0001$ . Q=quartile, where Q1 is lowest locus. \*Includes both working and non-working participants. †Felt sad, blue, or depressed for more than 2 consecutive weeks in past year.

Table 2: Psychosocial risk factors in cases and controls

Separate analyses in men and women showed that, by contrast to men, work stress did not seem to be associated with acute myocardial infarction in women ( $p=0.006$ , for interaction; table 3). However, this apparent interaction should be interpreted cautiously because several subgroup analyses were done. For all other factors, effects among women were similar to those seen among men, with no significant heterogeneity. A separate analysis in the 435 patients who died while still in hospital showed the same relation with stress as the total population of patients with acute myocardial infarction (not shown).

Table 4 presents results of the effect of general stress in various subgroups stratified by age, smoking, income, and education. The association with acute myocardial infarction did not differ greatly by any of these factors. Permanent stress was less prevalent in older versus younger participants, but no significant interaction was recorded between age and stress ( $p=0.13$ ). The effect of general stress was similar across different strata defined by age, income, or education and among smokers and

non-smokers. Table 5 describes the combined effects of general stress and depression, and of general stress and locus of control.

The prevalence of moderate or severe general stress ranged from 7.7% among controls in China to 35.3% in North America (figure 1). Despite this difference, stress was more common among cases than controls in all geographic regions, with odds ratios varying between 1.3 and 2.1, with the exception of central and eastern Europe, where the difference in rates of stress between cases and controls seemed to be only minor. Almost all ethnic groups were characterised by more general stress in cases than controls (figure 1). Although the prevalence of sadness and depression also varied between regions and various ethnic groups (figure 2), it was related to acute myocardial infarction in various subgroups examined.

## Discussion

Our study shows that several elements reflecting psychosocial stress are associated with increased risk of acute myocardial infarction. These factors include those that are subjective and perceived by the patients, such as stress, defined as tension or anxiety due to external influences. Some of these measures—eg, locus of control or depression—are not generally perceived by lay people to be stressors. Further, discrete external events (eg, major life events), which are less subjective and less likely to be subject to any biases, were also more frequent in cases than controls. The effect of stress is independent of socioeconomic status and smoking, and is by and large consistent across geographic regions, in different age groups, and in men and women. The excess risk of acute myocardial infarction associated with high levels of stress was still significant after adjusting for other cardiovascular risk factors.

During the past two decades, considerable evidence has accumulated with respect to the association of markers of stress and other psychosocial factors with coronary disease.<sup>4,23</sup> However, compared with many other biological and lifestyle risk factors, stress is a more difficult construct in that no consensus exists with respect to either definition or measurement. Further, stress is inevitably a subjective measurement, and hence is potentially open to biases and confounding. Nevertheless, the strong relation between self-reported stress and other more objective markers, such as life events or depression, and constructs such as locus of control (which is not associated by lay people as being related to stress) suggests face validity for the measures used in the study.

To date, most studies have dealt with stress at work, with stress outside the workplace receiving less attention. Both cross-sectional and prospective studies have shown a positive association between level of work stress and disease.<sup>8,10,13</sup> Number of work stressors has been associated with increased cardiovascular mortality

	Men			Women			
	Cases (%)	Controls (%)	Odds ratio (99% CI)	Cases (%)	Controls (%)	Odds ratio (99% CI)	p*
Stress at work							
Never	993 (20.3)	1504 (24.1)	1	145 (26.7)	264 (23.3)	1	0.006
Some of the time	2265 (46.4)	3315 (53.0)	1.00 (0.88–1.15)	234 (43.0)	608 (53.6)	0.68 (0.48–0.96)	
Several periods	1125 (23.0)	1117 (17.9)	1.45 (1.23–1.70)	124 (22.8)	207 (18.2)	1.02 (0.68–1.54)	
Permanent	499 (10.2)	316 (5.1)	2.34 (1.86–2.93)	41 (7.5)	56 (4.9)	1.11 (0.60–2.06)	
Stress at home							
Never	3314 (39.3)	4171 (41.6)	1	772 (28.7)	1172 (32.4)	1	0.12
Some of the time	4008 (47.5)	5005 (49.9)	1.01 (0.93–1.10)	1353 (50.4)	1868 (51.6)	1.16 (0.99–1.35)	
Several periods	862 (10.2)	718 (7.2)	1.53 (1.32–1.78)	426 (15.9)	461 (12.7)	1.53 (1.23–1.90)	
Permanent	249 (3.0)	135 (1.4)	2.36 (1.75–3.17)	135 (5.0)	118 (3.3)	1.88 (1.31–2.69)	
General stress†							
Never	2074 (24.6)	2682 (26.7)	1	703 (26.2)	1006 (27.8)	1	0.091
Some period, home or work	4024 (47.7)	5322 (53.1)	1.02 (0.93–1.13)	1328 (49.4)	1871 (51.7)	1.10 (0.94–1.30)	
Several periods, home or work	1654 (19.6)	1601 (16.0)	1.46 (1.29–1.66)	485 (18.1)	582 (16.1)	1.40 (1.13–1.73)	
Permanent, home or work	681 (8.1)	424 (4.2)	2.32 (1.93–2.80)	170 (6.3)	160 (4.4)	1.74 (1.25–2.40)	
Financial stress							
Little or none	3707 (44.0)	4921 (49.1)	1	1165 (43.4)	1707 (47.2)	1	0.74
Moderate	3495 (41.4)	3918 (39.1)	1.19 (1.09–1.29)	1130 (42.1)	1443 (39.9)	1.22 (1.05–1.41)	
Severe	1231 (14.6)	1190 (11.9)	1.33 (1.17–1.50)	391 (14.6)	469 (13.0)	1.33 (1.08–1.64)	
Stressful life events							
None	4857 (57.6)	6242 (62.2)	1	1568 (58.4)	2286 (63.2)	1	0.17
1	2186 (25.9)	2483 (24.8)	1.21 (1.10–1.33)	718 (26.7)	866 (23.9)	1.30 (1.11–1.52)	
2 or more	1390 (16.5)	1304 (13.0)	1.51 (1.34–1.70)	400 (14.9)	467 (12.9)	1.37 (1.12–1.68)	
Locus of control							
Q1	1843 (21.9)	1780 (17.8)	1	777 (28.9)	839 (23.2)	1	0.019
Q2	2130 (25.3)	2304 (23.0)	0.89 (0.79–1.00)	808 (30.1)	961 (26.6)	0.89 (0.74–1.07)	
Q3	2835 (33.6)	3668 (36.6)	0.73 (0.65–0.81)	779 (29.0)	1171 (32.4)	0.71 (0.60–0.85)	
Q4	1625 (19.3)	2277 (22.7)	0.72 (0.63–0.82)	322 (12.0)	648 (17.9)	0.55 (0.44–0.69)	
Feeling depressed in past 2 weeks							
No	6532 (77.5)	8387 (83.6)	1	1914 (71.3)	2857 (78.9)	1	0.17
Yes	1901 (22.5)	1642 (16.4)	1.53 (1.38–1.69)	772 (28.7)	762 (21.1)	1.60 (1.37–1.88)	

\*Interaction with sex. †Includes both working and non-working participants.

**Table 3: Psychosocial risk factors in cases and controls by sex**

\*Interaction with sex. †Includes both working and non-working participants.

**Table 3: Psychosocial risk factors in cases and controls by sex**

in the Multiple Risk Factor Intervention Trial.<sup>8</sup> During a 9-year follow-up period of 12 336 men, those with three or more work stressors had an increased risk of cardiovascular death (relative risk 1.26 [95% CI 1.07–1.48]), and the experience of divorce also increased risk (1.37 [1.09–1.72]) relative to those who remained married or single). In one prospective study of women with coronary heart disease,<sup>24</sup> level of marital stress, according to the Stockholm Marital Stress Scale, was associated with a higher risk of recurrent events. By contrast, data from a large North American study of nurses<sup>25</sup> showed no association between a measure of work strain and subsequent coronary heart disease. However, this study had low power (only 146 events) and in view of the homogeneity of the population studied (all nurses) there might have been little variation in the distribution of stress levels.

Other general measures of stress have also been used. In a prospective survey of middle-aged Swedish men,<sup>14</sup> self-reported permanent stress over a defined period in the recent past (as measured by a single-item question similar to that used in INTERHEART) was associated with an increased risk of incident coronary heart disease (odds ratio 1.5 [95% CI 1.2–1.9], after adjustment for conventional coronary risk factors) during the 12-year

follow-up. Similar results were noted in a large prospective study involving 281 cases in 73 424 Japanese men and women,<sup>9</sup> which reported an association between perceived mental stress and coronary heart disease mortality. The consistency of results that has been reported across multiple studies using different study designs and approaches provide a body of evidence that supports an association between various types of stress and coronary disease. In view of the large number of cases in the study, INTERHEART provides robust results, even in subgroups. Furthermore, the effect of stress on acute myocardial infarction in the present study was by and large consistent across different geographic regions, in different ethnic groups, in men and women, and at all ages.

Depression has been associated with an increased risk of coronary heart disease in both men and women.<sup>1–3</sup> In the present study, we showed that feeling sad, blue, or depressed for 2 weeks or more in a row was associated with acute myocardial infarction across different populations and across groups of people with different ethnic origins. A meta-analysis of 11 studies concluded that depression predicts the development of coronary heart disease in initially healthy people (odds ratio 1.64 [95% CI 1.29–2.08]).<sup>26</sup> Sensitivity analysis showed that

	Cases (%)	Controls (%)	Odds ratio (99% CI)	Cases (%)	Controls (%)	Odds ratio (99% CI)	Cases (%)	Controls (%)	Odds ratio (99% CI)
<b>Age (years)</b>	<b>&lt;56</b>			<b>56-64</b>			<b>&gt;64</b>		
General stress									
Never	782 (16.2)	1179 (18.7)	1	794 (26.5)	1106 (29.7)	1	1201 (36.4)	1403 (38.8)	1
Some period, home or work	2224 (46.2)	3436 (54.5)	0.98 (0.85-1.13)	1502 (50.0)	1956 (52.6)	1.11 (0.95-1.30)	1626 (49.3)	1801 (49.8)	1.07 (0.94-1.23)
Several periods, home or work	1263 (26.2)	1301 (20.6)	1.49 (1.26-1.76)	500 (16.7)	537 (14.4)	1.37 (1.11-1.69)	376 (11.4)	345 (9.5)	1.37 (1.10-1.71)
Permanent, home or work	548 (11.4)	391 (6.2)	2.10 (1.69-2.62)	206 (6.9)	122 (3.3)	2.50 (1.80-3.47)	97 (2.9)	71 (2.0)	1.82 (1.19-2.78)
<b>Smoking</b>	<b>Never</b>			<b>Former</b>			<b>Current</b>		
General stress									
Never	1062 (28.0)	1919 (27.5)	1	641 (28.9)	816 (28.4)	1	1047 (21.0)	897 (24.9)	1
Some period, home or work	1909 (50.3)	3789 (54.3)	1.01 (0.89-1.14)	1017 (45.9)	1429 (49.7)	1.01 (0.84-1.20)	2369 (47.6)	1888 (52.4)	1.12 (0.97-1.30)
Several periods, home or work	618 (16.3)	1017 (14.6)	1.41 (1.19-1.67)	391 (17.6)	499 (17.4)	1.25 (0.99-1.57)	1099 (22.1)	629 (17.5)	1.65 (1.37-1.98)
Permanent, home or work	208 (5.5)	251 (3.6)	2.07 (1.59-2.71)	169 (7.6)	132 (4.6)	2.17 (1.54-3.04)	465 (9.3)	189 (5.3)	2.33 (1.79-3.02)
<b>Income</b>	<b>1-2 (low)</b>			<b>3</b>			<b>4-5 (high)</b>		
General stress									
Never	1424 (25.0)	1901 (28.6)	1	541 (22.7)	729 (26.3)	1	779 (27.0)	1007 (25.0)	1
Some period, home or work	2802 (49.1)	3494 (52.5)	1.13 (1.00-1.27)	1248 (52.3)	1562 (56.4)	1.09 (0.91-1.31)	1248 (43.2)	2043 (50.6)	0.88 (0.75-1.04)
Several periods, home or work	1071 (18.8)	1001 (15.1)	1.52 (1.30-1.77)	444 (18.6)	384 (13.9)	1.74 (1.36-2.21)	586 (20.3)	759 (18.8)	1.21 (1.00-1.48)
Permanent, home or work	410 (7.2)	254 (3.8)	2.40 (1.90-3.04)	154 (6.5)	97 (3.5)	2.23 (1.52-3.28)	274 (9.5)	225 (5.6)	1.91 (1.45-2.52)
<b>Education</b>	<b>&lt;8 years</b>			<b>9-12 years</b>			<b>Trade/college/university</b>		
General stress									
Never	1387 (28.1)	1639 (31.8)	1	711 (24.2)	955 (27.7)	1	679 (20.9)	1093 (21.7)	1
Some period homework	2483 (50.4)	2693 (52.3)	1.11 (1.01-1.22)	1435 (48.9)	1891 (54.8)	1.07 (0.94-1.21)	1432 (44.0)	2605 (51.7)	0.94 (0.83-1.06)
Several periods	798 (16.2)	666 (12.9)	1.53 (1.34-1.75)	548 (18.7)	465 (13.5)	1.72 (1.46-2.03)	793 (24.4)	1052 (20.9)	1.28 (1.11-1.47)
Permanent home or work	262 (5.3)	156 (3.0)	2.23 (1.79-2.79)	240 (8.2)	139 (4.0)	2.53 (1.99-3.22)	348 (10.7)	289 (5.7)	2.02 (1.67-2.45)

Table 4: General stress by age, smoking, income, and education in cases and controls

clinical depression was a stronger predictor than depressive mood. Contrary to the findings in this meta-analysis, we did not find a so-called dose-response relation, because the risk of a myocardial infarction was similarly increased irrespective of the number of items in the depression scale that were positive. The prevalence of clinical depression among the controls in our study (7.0%) was comparable with estimates of mood disorders worldwide,<sup>27</sup> but lower than that for depressive mood (17.6%). This difference between studies suggests that those who were classified as being depressed in our study include some individuals with true clinical depression and some with a less specific reaction to stressors, which could account for the similarity of findings for depressive mood and stress.

Few studies have investigated the effect of external influences like financial stress or life events on risk of coronary disease. Findings of a case-control study<sup>13</sup>

showed that experience of one life event or more during the year preceding an acute myocardial infarction, and dissatisfaction with one's financial situation, was twice as common in cases than controls among men, but no significant relation was found among women. However, the lack of effect in women in the above study<sup>13</sup> might have been because it contained few women. Our study, which includes 2686 female cases and 3619 female controls, shows consistent relations in both men and women. In a Danish registry-based study,<sup>7</sup> an extreme external stressor, such as the death of a child, was shown to be associated with increased risk of future acute myocardial infarction; this finding is consistent with the present study, in which business failure, major intrafamily conflict, job loss, death of spouse, and violence were associated with increased risk.

Our questionnaires were derived from items that were previously shown in longitudinal studies to predict

General stress category												
	Never			Some periods, at home or work			Several periods, at home or work			Permanent, at home or work		
	Cases	Controls	Odds ratio (99% CI)	Cases	Controls	Odds ratio (99% CI)	Cases	Controls	Odds ratio (99% CI)	Cases	Controls	Odds ratio (99% CI)
<b>Depression</b>												
No	2359	3314	1.0	4319	6061	0.95 (0.80-1.14)	1333	1541	1.19 (0.97-1.45)	435	328	1.71 (1.32-2.21)
Yes	418	374	1.41 (1.21-1.64)	1033	1132	1.34 (1.07-1.68)	806	642	1.67 (1.33-2.11)	416	256	2.40 (1.83-3.15)
<b>Locus of control</b>												
Q4	619	917	1.0	867	1481	0.96 (0.80-1.15)	327	427	1.22 (1.00-1.49)	134	100	1.78 (1.38-2.30)
Q3	900	1186	1.03 (0.87-1.21)	1896	2693	0.99 (0.78-1.25)	605	780	1.26 (0.98-1.61)	213	180	1.84 (1.36-2.47)
Q2	781	995	1.34 (1.12-1.62)	1434	1676	1.29 (1.00-1.66)	535	480	1.64 (1.26-2.13)	188	114	2.40 (1.76-3.26)
Q1	477	590	1.55 (1.28-1.89)	1155	1343	1.49 (1.15-1.93)	672	496	1.90 (1.46-2.47)	316	190	2.77 (2.05-3.75)

Odds ratios adjusted for age, sex, smoking status, and geographic region.

Table 5: Odds ratios for combined effect of general stress and depression or locus of control on risk of acute myocardial infarction

cardiovascular events.<sup>14</sup> Because INTERHEART intended to recruit a large number of cases and controls from 52 countries, our questionnaires had to be simple and brief. The fact that such simple questions are informative across such diverse settings is a particular strength of our study.

Levels of stress, depression, and locus of control reported by INTERHEART controls varied substantially across regions. The differences in rates of these factors could be attributable to variations in interpretation of the questions in different cultures and the extent of social desirability with respect to responses, but this difference would not be expected to affect the comparisons between cases and controls, because both groups of participants were recruited within the same centre in every country. The validity of our conclusions is supported by the excess risk associated with most of the measures we used, and it is consistent in various regions. The odds ratios in various subgroups are essentially in the same direction, and the CIs generally overlap the overall odds ratio for every construct examined. Although we cannot exclude that self-reported stress might be subject to biases by cases spuriously reporting stress more often than controls, this is less likely to be a problem with measures such as low locus of control, life events, or perhaps depression, which are not typically judged to be associated with stress by lay individuals. The coherence of the results across both types of questions increases the plausibility of our findings, and the consistency across various geographic regions adds robustness and internal replication. Although quantification of any potential biases is difficult because of the case-control design we used, the similarity of our overall findings with those from several cohort studies is reassuring.

The present study has two additional potential limitations. First, we did not enrol patients who died before they could be interviewed. However, a separate analysis of the 435 patients who died in INTERHEART showed the same relation with stress as the total population of patients with acute myocardial infarction. Second, the experience of the acute myocardial infarction could theoretically alter a patient's perceptions about recent stress and mood. Even so, previous studies have shown that although perception of risk factor status may change, confirmatory search after the discovery of disease only moderately influences the recollection of symptoms in the month preceding disease.<sup>28</sup> A retrospective design has the advantage of assessing stress during the recent past, whereas longitudinal studies can underestimate the effects of a past period of intense stress owing to the long interval between an interview and the occurrence of an event, nor can they explore the relation of stress in the period just before the myocardial infarction. Also, true levels of stress might vary over time in the same individual. In 292 individuals, repeat measures obtained at an interval of greater than a year indicated a moderate correlation of 0.5 (compared

with, for example, 0.7 for apolipoprotein B and 0.9 for smoking). This finding suggests that the effect of an analysis adjusting for variability (eg, regression-dilution bias) could indicate an even stronger relation than that described in the present article. However, adjustment for regression-dilution bias simultaneously across multiple variables is complex, and we are not aware of any study that has done that.

Not all prospective studies have reported a significant relation between psychological stress and ischaemic heart disease.<sup>29–31</sup> The lack of an association could be attributable to low power because of few events<sup>29</sup> in several studies and potential waning of an effect when events arise after measurement of stress.<sup>29–31</sup> Residual confounding of low socioeconomic status has been proposed as a potential explanation.<sup>30</sup> However, in INTERHEART, all associations recorded between psychosocial stress factors and myocardial infarction were unchanged after adjustment for income and education.

The mechanism by which psychosocial factors increase the risk of myocardial infarction is complex. In experimental studies, worsened coronary atherosclerosis<sup>32</sup> and endothelial dysfunction<sup>33</sup> happen in response to social disruption. Several studies have shown links between psychosocial variables and vascular function,<sup>34,35</sup> inflammation,<sup>36</sup> increased blood clotting, and decreased fibrinolysis.<sup>37,38</sup> The exact pathophysiological nature of the influence of psychosocial factors remains to be determined, as does the temporal sequence of events.

In conclusion, our study indicates that psychosocial stressors are related to increased risk of acute myocardial infarction. For severe global stress, the size of the effect was less than that for smoking but comparable with hypertension and abdominal obesity.<sup>16</sup> Our study is unique in having evaluated simultaneously multiple elements of stress, in inclusion of perception of stress and life events, and in objective constructs such as locus of control. In view of the large number of participants and inclusion of multiple populations and ethnic groups, we have shown that the effects of stress on acute myocardial infarction are similar in men and women, in people of various ages, and in all geographic regions of the world that we studied. The PAR for each of the measures ranged from 8% to 16%, and collectively adds up to 33% for all variables. If this effect is truly causal, the importance of psychosocial factors is much more important than commonly recognised, and might contribute to a substantial proportion of acute myocardial infarction.

#### Contributors

S Yusuf initiated the INTERHEART study and supervised the project, data analysis, and writing of the report. A Rosengren had main responsibility for writing the report. S Hawken did statistical analyses. S Ounpuu coordinated the project. All other investigators facilitated and supervised the study in their own country and commented on drafts of the report.

**Conflict of interest statement**

We declare that we have no conflict of interest.

**Acknowledgments**

The INTERHEART study was funded by the Canadian Institutes of Health Research, the Heart and Stroke Foundation of Ontario, the International Clinical Epidemiology Network (INCLIN), and through unrestricted grants from several pharmaceutical companies (with major contributions from AstraZeneca, Novartis, Hoechst Marion Roussel [now Aventis], Knoll Pharmaceuticals [now Abbott], Bristol Myers Squibb, King Pharma, and Sanofi-Syhelabo), and by various national bodies in different countries (see webappendix2; <http://image.thelancet.com/extras/04art8001webappendix2.pdf>).

**References**

- Anda R, Williamson D, Jones D, et al. Depressed affect, hopelessness, and the risk of ischemic heart disease in a cohort of US adults. *Epidemiology* 1993; **4**: 285–94.
- Barefoot JC, Schroll M. Symptoms of depression, acute myocardial infarction, and total mortality in a community sample. *Circulation* 1996; **93**: 1976–80.
- Ferketich AK, Schwartzbaum JA, Frid DJ, Moeschberger ML. Depression as an antecedent to heart disease among women and men in the NHANES I study. *Arch Intern Med* 2000; **160**: 1261–68.
- Hemingway H, Marmot M. Evidence based cardiology: psychosocial factors in the aetiology and prognosis of coronary heart disease: systematic review of prospective cohort studies. *BMJ* 1999; **318**: 1460–67.
- Yoshimasu K. Relation of type A behavior pattern and job-related psychosocial factors to nonfatal myocardial infarction: a case-control study of Japanese male workers and women. *Psychosom Med* 2001; **63**: 797–804.
- Sacker A, Bartley MJ, Frith D, Fitzpatrick RM, Marmot MG. The relationship between job strain and coronary heart disease: evidence from an English sample of the working male population. *Psychol Med* 2001; **31**: 279–90.
- Li J, Hansen D, Mortensen PB, Olsen J. Myocardial infarction in parents who lost a child: a nationwide prospective cohort study in Denmark. *Circulation* 2002; **106**: 1634–39.
- Matthews KA, Gump BB. Chronic work stress and marital dissolution increase risk of posttrial mortality in men from the Multiple Risk Factor Intervention Trial. *Arch Intern Med* 2002; **162**: 309–15.
- Iso H, Date C, Yamamoto A, et al. Perceived mental stress and mortality from cardiovascular disease among Japanese men and women: the Japan Collaborative Cohort Study for Evaluation of Cancer Risk Sponsored by Monbusho (JACC Study). *Circulation* 2002; **106**: 1229–36.
- Kivimäki M, Leino-Arjas P, Luukkonen R, Riihimäki H, Vahtera J, Kirjonen J. Work stress and risk of cardiovascular mortality: prospective cohort study of industrial employees. *BMJ* 2002; **325**: 857–61.
- Stansfeld SA, Fuhrer R, Shipley MJ, Marmot MG. Psychological distress as a risk factor for coronary heart disease in the Whitehall II Study. *Int J Epidemiol* 2002; **31**: 248–55.
- Welin C, Rosengren A, Wedel H, Wiklund I, Wilhelmsen L. Psychological characteristics in patients with myocardial infarction: a case-control study. *Cardiovasc Risk Factors* 1994; **4**: 154–61.
- Welin C, Rosengren A, Wedel H, Wilhelmsen L. Myocardial infarction in relation to work, family and life events. *Cardiovasc Risk Factors* 1995; **5**: 30–38.
- Rosengren A, Tibblin G, Wilhelmsen L. Self-perceived psychological stress and incidence of coronary artery disease in middle-aged men. *Am J Cardiol* 1991; **68**: 1171–75.
- Prescott E, Holst C, Gronbaek M, Schnohr P, Jensen G, Barefoot J. Vital exhaustion as a risk factor for ischaemic heart disease and all-cause mortality in a community sample: a prospective study of 4084 men and 5479 women in the Copenhagen City Heart Study. *Int J Epidemiol* 2003; **32**: 990–97.
- Yusuf S, Hawken S, Öunpuu S, et al. Effect of potentially modifiable risk factors associated with myocardial infarction in 52 countries (the INTERHEART study): case-control study. *Lancet* 2004; **364**: 937–52.
- Öunpuu S, Negassa A, Yusuf S. INTERHEART: a global study of risk factors for acute myocardial infarction. *Am Heart J* 2001; **141**: 711–21.
- Bobak M, Pikhart H, Rose R, Hertzman C, Marmot M. Socioeconomic factors, material inequalities, and perceived control in self-rated health: cross-sectional data from seven post-communist countries. *Soc Sci Med* 2000; **51**: 1343–50.
- Patten SB. Performance of the Composite International Diagnostic Interview Short Form for major depression in community and clinical samples. *Chronic Dis Can* 1997; **18**: 109–12.
- Breslow N, Day N. Statistical methods in cancer research, vol 1: the analysis of case-control studies. Lyon: IARC Scientific Publications, 1980.
- Benichou J, Gail M. Variance calculations and confidence intervals for estimates of the attributable risk based on logistic models. *Biometrics* 1990; **46**: 991–1003.
- Engel L, Chow W, Vaughan T. Population attributable risks of esophageal and gastric cancers. *J Natl Cancer Inst* 2003; **95**: 1404–13.
- Marmot M, Stansfeld S. Stress and the heart: psychosocial pathways to coronary heart disease. London: BMJ books, 2001.
- Orth-Gomer K, Wamala SP, Horsten M, Schenck-Gustafsson K, Schneiderman N, Mittleman MA. Marital stress worsens prognosis in women with coronary heart disease: the Stockholm Female Coronary Risk Study. *JAMA* 2000; **284**: 3008–14.
- Lee S, Colditz G, Berkman L, Kawachi I. A prospective study of job strain and coronary heart disease in US women. *Int J Epidemiol* 2002; **31**: 1147–53.
- Rugulies R. Depression as a predictor for coronary heart disease: a review and meta-analysis. *Am J Prev Med* 2002; **23**: 51–61.
- Demyttenaere K, Bruffaerts R, Posada-Villa J, et al. Prevalence, severity, and unmet need for treatment of mental disorders in the World Health Organization World Mental Health Surveys. *JAMA* 2004; **291**: 2581–90.
- Croyle RT, Sande GN. Denial and confirming search: paradoxical consequences of medical diagnosis. *J Appl Soc Psychol* 1988; **18**: 473–90.
- Moore L, Meyer F, Perusse M, et al. Psychological stress and incidence of ischaemic heart disease. *Int J Epidemiol* 1999; **28**: 652–58.
- MacLeod J, Smith GD, Heslop P, Metcalfe C, Carroll D, Hart C. Are the effects of psychosocial exposures attributable to confounding? Evidence from a prospective observational study on psychological stress and mortality. *J Epidemiol Community Health* 2001; **55**: 878–84.
- Eaker ED, Sullivan LM, Kelly-Hayes M, D'Agostino RB Sr, Benjamin EJ. Does job strain increase the risk for coronary heart disease or death in men and women? The Framingham Offspring Study. *Am J Epidemiol* 2004; **159**: 950–8.
- Kaplan JR, Pettersson K, Manuck SB, Olsson G. Role of sympathoadrenal medullary activation in the initiation and progression of atherosclerosis. *Circulation* 1991; **84**: V123–32.
- Strawn WB, Bondjers G, Kaplan JR, et al. Endothelial dysfunction in response to psychosocial stress in monkeys. *Circ Res* 1991; **68**: 1270–79.
- Ghiadoni L, Donald AE, Cropley M, et al. Mental stress induces transient endothelial dysfunction in humans. *Circulation* 2000; **102**: 2473–78.
- Kop WJ, Krantz DS, Howell RH, et al. Effects of mental stress on coronary epicardial vasomotion and flow velocity in coronary artery disease: relationship with hemodynamic stress responses. *J Am Coll Cardiol* 2001; **37**: 1359–66.
- Lewthwaite J, Owen N, Coates A, Henderson B, Steptoe A. Circulating human heat shock protein 60 in the plasma of British civil servants: relationship to physiological and psychosocial stress. *Circulation* 2002; **106**: 196–201.
- Brunner E, Davey-Smith G, Marmot M, Canner R, Beksinska M, O'Brien J. Childhood social circumstances and psychosocial and behavioural factors as determinants of plasma fibrinogen. *Lancet* 1996; **347**: 1008–13.
- von Kanel R, Mills PJ, Fainman C, Dimsdale JE. Effects of psychological stress and psychiatric disorders on blood coagulation and fibrinolysis: a biobehavioral pathway to coronary artery disease? *Psychosom Med* 2001; **63**: 531–44.

# Effect of potentially modifiable risk factors associated with myocardial infarction in 52 countries (the INTERHEART study): case-control study

Salim Yusuf, Steven Hawken, Stephanie Ôunpuu, Tony Dans, Alvaro Avezum, Fernando Lanas, Matthew McQueen, Andrzej Budaj, Prem Pais, John Varigos, Liu Lisheng, on behalf of the INTERHEART Study Investigators\*



Lancet 2004; 364: 937–52

Published online  
September 3, 2004  
<http://image.thelancet.com/extras/04art8001web.pdf>

See [Comment](#) page 912

\*Listed at end of report.

Population Health Research  
Institute, Hamilton General  
Hospital, 237 Barton Street  
East, Hamilton, Ontario,  
Canada L8L 2X2  
(Prof S Yusuf DPhil,  
S Ôunpuu PhD, S Hawken MSc,  
T Dans MD, A Avezum MD,  
F Lanas MD, M McQueen FRCP,  
A Budaj MD, P Pais MD,  
J Varigos BSc, L Lisheng MD)

Correspondence to:  
Prof Salim Yusuf  
[yusufs@mcmaster.ca](mailto:yusufs@mcmaster.ca)

## Summary

**Background** Although more than 80% of the global burden of cardiovascular disease occurs in low-income and middle-income countries, knowledge of the importance of risk factors is largely derived from developed countries. Therefore, the effect of such factors on risk of coronary heart disease in most regions of the world is unknown.

**Methods** We established a standardised case-control study of acute myocardial infarction in 52 countries, representing every inhabited continent. 15 152 cases and 14 820 controls were enrolled. The relation of smoking, history of hypertension or diabetes, waist/hip ratio, dietary patterns, physical activity, consumption of alcohol, blood apolipoproteins (Apo), and psychosocial factors to myocardial infarction are reported here. Odds ratios and their 99% CIs for the association of risk factors to myocardial infarction and their population attributable risks (PAR) were calculated.

**Findings** Smoking (odds ratio 2·87 for current vs never, PAR 35·7% for current and former vs never), raised ApoB/ApoA1 ratio (3·25 for top vs lowest quintile, PAR 49·2% for top four quintiles vs lowest quintile), history of hypertension (1·91, PAR 17·9%), diabetes (2·37, PAR 9·9%), abdominal obesity (1·12 for top vs lowest tertile and 1·62 for middle vs lowest tertile, PAR 20·1% for top two tertiles vs lowest tertile), psychosocial factors (2·67, PAR 32·5%), daily consumption of fruits and vegetables (0·70, PAR 13·7% for lack of daily consumption), regular alcohol consumption (0·91, PAR 6·7%), and regular physical activity (0·86, PAR 12·2%), were all significantly related to acute myocardial infarction ( $p < 0·0001$  for all risk factors and  $p = 0·03$  for alcohol). These associations were noted in men and women, old and young, and in all regions of the world. Collectively, these nine risk factors accounted for 90% of the PAR in men and 94% in women.

**Interpretation** Abnormal lipids, smoking, hypertension, diabetes, abdominal obesity, psychosocial factors, consumption of fruits, vegetables, and alcohol, and regular physical activity account for most of the risk of myocardial infarction worldwide in both sexes and at all ages in all regions. This finding suggests that approaches to prevention can be based on similar principles worldwide and have the potential to prevent most premature cases of myocardial infarction.

## Introduction

Worldwide, cardiovascular disease is estimated to be the leading cause of death and loss of disability-adjusted life years.<sup>1</sup> Although age-adjusted cardiovascular death rates have declined in several developed countries in past decades, rates of cardiovascular disease have risen greatly in low-income and middle-income countries,<sup>1,2</sup> with about 80% of the burden now occurring in these countries. Effective prevention needs a global strategy based on knowledge of the importance of risk factors for cardiovascular disease in different geographic regions and among various ethnic groups.

Current knowledge about prevention of coronary heart disease and cardiovascular disease is mainly derived from studies done in populations of European origin.<sup>2</sup> Researchers are unsure to what extent these findings apply worldwide. Some data suggest that risk factors for coronary heart disease vary between populations—eg, lipids are not associated with this disorder in south Asians,<sup>3</sup> and increases in blood pressure might be more important in Chinese people.<sup>4</sup> Even if the association of a

risk factor with coronary heart disease is similar across populations, prevalence of this factor might vary, resulting in different population attributable risks (PAR)—eg, serum cholesterol might be lower in Chinese populations.<sup>4</sup> On the other hand, these apparent variations between ethnic populations could be attributable to differences between studies in their design and analysis, information obtained, and small sample sizes.

To clarify whether the effects of risk factors vary in different countries or ethnic groups, a large study undertaken in many countries—representing different regions and ethnic groups and using standardised methods—is needed, with the aim to investigate the relation between risk factors and coronary heart disease. Such a study could also estimate the importance of known risk factors on the PAR for acute myocardial infarction. This aim, however, needs either very large cohort trials or case-control studies with many events—eg, several thousands of cases of myocardial infarction in whom all (or most) currently

known risk factors are measured. We judged the latter most practical.

INTERHEART is a large, international, standardised, case-control study, designed as an initial step to assess the importance of risk factors for coronary heart disease worldwide (slides available at <http://www.phri.ca/interheart>).<sup>5</sup> We aimed to include about 15 000 cases and a similar number of controls from 52 countries, representing all inhabited continents. Specific objectives are to determine the strength of association between various risk factors and acute myocardial infarction in the overall study population and to ascertain if this association varies by geographic region, ethnic origin, sex, or age. A key secondary objective is to estimate the PAR for risk factors and their combinations in the overall population and in various subgroups. This report focuses on the association of nine easily measured protective or risk factors (smoking, lipids, self-reported hypertension or diabetes, obesity, diet, physical activity, alcohol consumption, and psychosocial factors) to first myocardial infarction.

## Methods

### Participants

Study participants were recruited from 262 centres from 52 countries in Asia, Europe, the Middle East, Africa, Australia, North America, and South America (web-table 1; <http://image.thelancet.com/extras/04art8001webtable1.pdf>). The national coordinator selected centres within every country on the basis of feasibility. To identify first cases of acute myocardial infarction, all patients (irrespective of age) admitted to the coronary care unit or equivalent cardiology ward, presenting within 24 h of symptom onset, were screened. Cases were eligible if they had characteristic symptoms plus electrocardiogram changes indicative of a new myocardial infarction (webappendix 1; <http://image.thelancet.com/extras/04art8001webappendix1.pdf>).

At least one age-matched (up to 5 years older or younger) and sex-matched control was recruited per case, using specific criteria. Exclusion criteria for controls were identical to those described for cases, with the additional criterion that controls had no previous diagnosis of heart disease or history of exertional chest pain. The overall median interval from recruitment of cases to inclusion of controls was 1·5 months. Hospital-based controls (58%) were individuals who had a wide range of disorders unrelated to known or potential risk factors for acute myocardial infarction and were admitted to the same hospital as the matching case. Community-based controls (36%) were attendants or relatives of a patient from a non-cardiac ward or an unrelated (not first-degree relative) attendant of a cardiac patient. In the remaining controls, 3% were from an undocumented source and 3% were recruited through the WHO MONICA study.<sup>6</sup>

### Procedures

Structured questionnaires were administered and physical examinations were undertaken in the same manner in cases and controls. Information about demographic factors, socioeconomic status (education, income), lifestyle (smoking, leisure time, physical activity, and dietary patterns), personal and family history of cardiovascular disease, and risk factors (hypertension, diabetes mellitus) was obtained. Psychosocial factors (depression, locus of control, perceived stress, and life events) were systematically recorded and integrated into one score: details are provided in the accompanying paper.<sup>7</sup> Height, weight, waist and hip circumferences, and heart rate were determined by a standardised protocol. Waist and hip circumferences were measured with a non-stretchable standard tape measure: waist measurements were obtained over the unclothed abdomen at the narrowest point between the costal margin and iliac crest, and hip circumferences over light clothing at the level of the widest diameter around the buttocks. Although blood pressure at the time of examination was recorded in both cases and controls, the levels in cases would be systematically affected by the myocardial infarction and treatments—eg,  $\beta$  blockers, nitrates, and angiotensin-converting-enzyme inhibitors—that could lower blood pressure. Therefore, only self-reported history of hypertension is used in the analysis.

Non-fasting blood samples (20 mL) were drawn from every individual and centrifuged within 2 h of admission, separated into six equal volumes, and frozen immediately at  $-20^{\circ}\text{C}$  or  $-70^{\circ}\text{C}$  after processing. Centres were instructed to draw blood from cases within 24 h of symptom onset. However, because of delays in patient presentation, especially in some low-income countries, blood samples could only be obtained within 24 h in two-thirds of cases. Samples were shipped in nitrogen vapour tanks by courier from every site to a blood storage site, where they were stored at  $-160^{\circ}\text{C}$  in liquid nitrogen (Hamilton, Canada) or at  $-70^{\circ}\text{C}$  (India and China). Blood samples from all countries other than China were analysed in Hamilton for total cholesterol, HDL cholesterol, and apolipoproteins B (ApoB) and A1 (ApoA1).

Immunoturbidimetric assays were used to measure apolipoprotein concentrations (Roche/Hitachi 917 analyser with Tina-quant ApoB version 2 and ApoA1 version 2 kits; Roche Diagnostics, Mannheim, Germany). The ApoB method was standardised against the IFCC SP3-07 reference standard<sup>8</sup> and the ApoA1 method against the IFCC SP1-01 reference preparation.<sup>9</sup> The same measurement kits and a Roche/Hitachi 911 analyser were used in Beijing, China. Both laboratories measured the same lot numbers of Precinorm and Precipath controls (Roche Diagnostics) in every run, and in every patient sample analysis run in China, two study patients and two serum reference pool samples (pool A and B) were measured that had previously been analysed

See **Articles** page 953

in the central core laboratory in Canada. Because apolipoprotein concentrations are not affected by the fasting status of the individual (unlike calculated LDL), we used the ApoB/ApoA1 ratio as an index of abnormal lipids in the current analysis.<sup>10</sup> Moreover, this ratio was predictive of myocardial infarction in subsets of patients (<12 h, 12–24 h, and >24 h after symptoms) in the present study (data not shown). Detailed information on lipoprotein fractions will be reported separately.

All data were transferred to the Population Health Research Institute, McMaster University, and Hamilton Health Sciences, Canada, where quality-control checks and statistical analyses were done. Data on smoking were missing in 1·1% of participants, hypertension in 0·6%, diabetes in 0·7%, psychosocial variables in 11%, physical activity in 1·1%, diet in 2·1%, and waist and hip measurements in 3·5%. Blood samples were available in 21 508 (79%) of 27 098 cases and controls.

INTERHEART was approved by appropriate regulatory and ethics committees in all participating countries and centres. All participants provided informed consent before taking part in the study.

We defined current smokers as individuals who smoked any tobacco in the previous 12 months and included those who had quit within the past year. Former smokers were defined as those who had quit more than a year earlier. For waist/hip ratio, tertiles were calculated separately for men and women based on the overall control data. The cutoffs used were 0·90 and 0·95 in men and 0·83 and 0·90 in women, to divide participants into thirds. Cutoffs for ApoB/ApoA1 ratios (deciles and quintiles) were derived from all controls (men and women). Region-specific cutoffs did not alter the results. Individuals were judged to be physically active if they were regularly involved in moderate (walking, cycling, or gardening) or strenuous exercise (jogging, football, and vigorous swimming) for 4 h or more a week. Regular alcohol use was defined as consumption three or more times a week. The combined psychosocial index was devised with a combination of the parameter estimates from the completely adjusted multivariate logistic regression model. The score was based on a combination of depression versus none, stress at work or at home (general stress variable) versus none, moderate or severe financial stress versus minimal or none, one or more life events versus none, and a locus of control score in the lower three quartiles versus the top quartile of the distribution.

### Statistical analysis

Simple associations were assessed with frequency tables and Pearson's  $\chi^2$  tests for two independent proportions. For comparison of prevalence across distinct subgroups—eg, by region, country, or ethnic group—potential differences in age structure of the populations were accounted for by direct standardisation of the frequencies to the overall INTERHEART age distribution with a five-level age-stratification factor

(<45, 46–55, 56–65, 66–70, >70).<sup>11</sup> Means and medians were calculated to summarise continuous effects and were compared by *t* tests or appropriate non-parametric tests when distributional assumptions were in doubt. When data have been categorised by tertiles, quintiles, or deciles, these were based on the overall control data. For waist/hip ratio, sex-specific cutoffs were used. For protective factors (exercise, diet, and alcohol), the PAR is calculated for the group without the exposure.

The findings presented are for models fitted with unconditional logistic regression, adjusted for the matching criteria, for two reasons. First, unmatched analyses were used because for 14% (1763/12461) of cases of myocardial infarction and 5% (738/14637) of controls, perfect matching was not possible. Undertaking a strict matched analysis would mean relevant loss of information because of the exclusion of these participants. Moreover, when data on a risk factor were missing in a case or control, the entire pair would be excluded from all analyses. Therefore, we widened the age-matching criteria and used frequency matching of cases and controls, using age and sex strata. Second, there was general agreement for key results among the many methods compared (conditional logistic regression, mixed models, and unconditional logistic regression, with adjustment for matching criteria). Estimated odds ratios and CIs calculated with the different methods were within 5% of each other, with a slight attenuation of effect estimates in the unconditional versus conditional models (webtable 2; <http://image.thelancet.com/extras/04art8001webtable2.pdf>).<sup>11</sup> Hence, findings presented are adjusted for age, sex, geographic region, and potential confounders, and should be interpreted as providing a slight underestimation of effect sizes for most comparisons.

Adjusted odds ratios for combinations of risk factors can be derived from their respective model coefficients in the multivariate logistic model. By summation of model coefficients and taking the antilog (panel) the combined effect of combinations of exposures can be estimated. Estimates of odds ratios and accompanying 99% CIs are presented for every risk factor and their combinations. Statistical analyses and graphics were produced with SAS version 8.2 (SAS, Cary, NC, USA) and S-Plus version 6 (Insightful, Seattle, WA, USA). All statistical tests of hypotheses are two-sided. PARs and 99% CIs were calculated for various risk factors in the study by a method based on unconditional logistic regression.<sup>12</sup> The PARs presented are adjusted for confounders in a similar manner to the corresponding logistic regression models for odds ratio estimates and, where indicated, are stratified by subgroups of interest. PAR estimates were calculated by the interactive risk attributable program software (US National Cancer Institute, 2002).<sup>13</sup>

For a simple dichotomous exposure and disease, and no adjustment for confounding, the usual formula for PAR was used (panel).<sup>12</sup> PAR adjusted for confounding

**Panel: Formulae**

$$\text{Antilog } e^{\Delta \ln R}$$

$$\text{PAR} = \frac{\text{Pr}(E)(R-1)}{\text{Pr}(E)(R-1)+1}$$

where Pr(E) is probability of exposure to the risk factor and R is the relative risk of the disease in exposed versus unexposed individuals.

PAR adjusted for confounding

$$1 - \sum_{i=2}^I \sum_{j=2}^J \rho_{ij} R_{ij}^{-1}$$

where

$$R_{ij} = \frac{\text{Pr}(D=1|X=x_i, C=c_j)}{\text{Pr}(D=1|X=x_i, C=c_1)}$$

and

$$\rho_{ij} = \text{Pr}(X=x_i, C=c_j|D=1)$$

X is the exposure level, C is the confounder level, and D is disease status (D=0, disease is absent; D=1, disease is present).

is also shown in the panel. For variance estimates, the reader is referred to Benichou and Gail<sup>15</sup> since the derivations and formulae are complex. CI calculations were based on this method using a logit transformation approach, apart from when PAR estimates were negative, in which case conventional Wald type CIs were used.

**Role of the funding source**

The sponsors of the study had no role in study design, data collection, data analysis, data interpretation, or writing of the report. The corresponding author had full access to all the data in the study and had final responsibility for the decision to submit for publication.

**Results**

Between February, 1999, and March, 2003, 15 152 cases and 14 820 controls were enrolled. 1531 cases were diagnosed as having unstable angina, 260 had insufficient data, 205 did not have coronary artery disease, and 695 had a previous myocardial infarction. For 74 controls data were missing and 109 had previous coronary heart disease. Therefore, 12 461 cases and 14 637 controls are included in the analysis. Table 1 shows the distribution of participants by region and ethnic origin. 9459 cases (76%) and 10 851 controls (74%) were male.

Table 2 shows the median age of presentation of cases. The overall median age of cases with first acute myocardial infarction is about 9 years lower in men than in women in all regions of the world. However, the proportion of male cases was highest in countries with a younger age of presentation of acute myocardial infarction—eg, 85% of cases in south Asia and 86% in the Middle East were male compared with 74% in western Europe, 68% in central and eastern Europe, and 70% in

	Cases (n=12 461)	Controls (n=14 637)
<b>Geographic region</b>		
Western Europe	664	767
Central and eastern Europe	1727	1927
Middle East	1639	1786
Africa	578	789
South Asia	1732	2204
China and Hong Kong	3030	3056
Southeast Asia and Japan	969	1199
Australia and New Zealand	589	681
South America and Mexico	1237	1888
North America	296	340
<b>Ethnic origin</b>		
European	3314	3710
Chinese	3130	3167
South Asian	2171	2573
Other Asian	871	1073
Arab	1306	1479
Latin American	1141	1834
Black African	157	369
Coloured African	311	339
Other	60	93

**Table 1: Distribution of study population**

China. Among regions, striking variations were noted in the age of first presentation of acute myocardial infarction, with the youngest patients in south Asia (median age 53 years) and the Middle East (51 years), and the oldest patients in western Europe, China, and Hong Kong (63 years). The highest proportion of cases with first acute myocardial infarction at age 40 years or younger was in men from the Middle East (12·6%), Africa (10·9%), and south Asia (9·7%) and the lowest proportion was in women from China and Hong Kong (1·2%), South America (1·0%), and central and eastern Europe (0·9%).

**Overall effect of risk factors**

Table 3 provides the overall odds ratios for individual risk factors adjusted for age, sex, smoking status, and region and by multivariate adjustment for all risk factors. All risk factors were significantly ( $p < 0.0001$ ) related to acute myocardial infarction, except alcohol, which had a weaker association ( $p = 0.03$ ). After multivariate analysis, current smoking and raised ApoB/ApoA1 ratio (top vs lowest quintile) were the two strongest risk factors, followed by history of diabetes, hypertension, and psychosocial factors (table 3). Body-mass index was related to risk of myocardial infarction, but this relation was weaker than that of abdominal obesity (waist/hip ratio), with body-mass index becoming non-significant with the inclusion of waist/hip ratio in the multivariate model (data not shown). Before multivariate adjustment, abdominal obesity (top vs lowest tertile) doubled the risk of acute myocardial infarction, but the effects were substantially diminished after adjustment for other risk factors, especially apolipoproteins. Daily consumption of fruits or vegetables, moderate or strenuous physical exercise, and consumption of alcohol three or more times per week, were protective (table 3).

	Overall			Men			Women		
	Number	Median age (IQR)	% <40 years (n)	Number	Median age (IQR)	% <40 years (n)	Number	Median age (IQR)	% <40 years (n)
<b>Geographic region</b>									
Western Europe	664	63 (54–72)	2.7 (18)	493	61 (53–70)	2.8 (14)	171	68 (59–76)	2.3 (4)
Central and eastern Europe	1727	62 (52–70)	2.9 (51)	1173	59 (50–68)	3.9 (46)	554	68 (59–74)	0.9 (5)
North America	296	59 (50–71)	4.0 (12)	210	58 (49–68)	3.3 (7)	86	64 (52–75)	5.8 (5)
South America and Mexico	1237	60 (51–70)	3.4 (42)	926	59 (50–68)	4.2 (39)	311	65 (56–73)	1.0 (3)
Australia and New Zealand	589	60 (51–69)	5.3 (31)	464	58 (50–67)	5.6 (26)	125	66 (59–74)	4.0 (5)
Middle East	1639	51 (45–59)	11.2 (184)	1410	50 (44–57)	12.6 (177)	229	57 (50–65)	3.1 (7)
Africa	578	54 (47–62)	9.7 (56)	385	52 (46–61)	10.9 (42)	193	56 (49–65)	7.3 (14)
South Asia	1732	53 (46–61)	8.9 (54)	1480	52 (45–60)	9.7 (143)	252	60 (50–66)	4.4 (11)
China and Hong Kong	3030	63 (53–70)	4.5 (135)	2131	60 (50–68)	5.8 (124)	899	67 (62–72)	1.2 (11)
Southeast Asia and Japan	969	57 (49–65)	7.0 (68)	787	55 (47–64)	8.3 (65)	182	63 (56–68)	1.7 (3)
<b>Ethnic origin</b>									
European	3314	62 (52–71)	3.2 (107)	2371	59 (51–69)	3.8 (89)	943	68 (58–75)	1.9 (18)
Chinese	3130	63 (53–70)	4.4 (139)	2217	60 (50–68)	5.8 (128)	913	67 (61–72)	1.2 (11)
South Asian	2171	52 (45–60)	10.6 (231)	1889	50 (45–60)	11.7 (220)	282	60 (51–66)	3.9 (11)
Other Asian	871	57 (48–65)	7.0 (61)	705	55 (47–64)	8.2 (58)	166	63 (56–68)	1.8 (3)
Arab	1306	53 (46–60)	9.0 (118)	1083	52 (45–59)	10.3 (111)	223	57 (50–65)	3.1 (7)
Latin American	1141	60 (51–69)	3.7 (42)	854	58 (50–67)	4.5 (38)	287	64 (55–72)	1.4 (4)
Black African	157	52 (46–61)	14.0 (22)	98	52 (46–59)	17.4 (17)	59	54 (48–67)	8.5 (5)
Coloured African	311	54 (47–63)	8.7 (27)	196	52 (46–62)	9.7 (19)	115	58 (49–65)	7.0 (8)
Other	60	57 (48–64)	6.7 (4)	46	53 (48–62)	6.5 (3)	14	63 (59–73)	7.1 (1)
<b>Overall</b>	<b>12 461</b>	<b>58 (49–67)</b>	<b>6.0 (751)</b>	<b>9459</b>	<b>56 (48–65)</b>	<b>7.2 (683)</b>	<b>3002</b>	<b>65 (56–72)</b>	<b>2.3 (68)</b>

Table 2: Median age (years) of presentation of cases

A strong and graded relation was noted between numbers smoked and risk of myocardial infarction, with the risk increasing at every increment, so that individuals smoking greater than 40 cigarettes per day had an odds ratio of 9.16 (99% CI 6.18–13.58; figure 1). The ApoB/ApoA1 ratio also showed a graded relation with myocardial infarction risk, with no evidence of a threshold, with an odds ratio of 4.73 (99% CI 3.93–5.69) for the top versus the lowest decile of ApoB/ApoA1 ratio (figure 1).


### Cumulative effect of risk factors

Figure 2 shows the effect of multiple risk factors on increased risk of myocardial infarction. Together, current smoking, hypertension, and diabetes increased the odds ratio for acute myocardial infarction to 13.01 (99% CI 10.69–15.83) compared to those without these risk factors, and they accounted for 53% of the PAR of acute myocardial infarction. Addition of ApoB/ApoA1 ratio (top vs lowest quintile) increased the odds ratio to

Risk factor	Prevalence		Odds ratio (99% CI) adjusted for age, sex, and smoking (OR 1)	PAR (99% CI)	Odds ratio (99% CI) adjusted additionally for all other risk factors (OR 2)	PAR 2 (99% CI)
	Controls (%)	Cases (%)				
Current smoking*	26.76	45.17	2.95 (2.72–3.20)	–	2.87 (2.58–3.19)	–
Current and former smoking*	48.12	65.19	2.27 (2.11–2.44)	36.4% (33.9–39.0)	2.04 (1.86–2.25)	35.7% (32.5–39.1)
Diabetes	7.52	18.45	3.08 (2.77–3.42)	12.3% (11.2–13.5)	2.37 (2.07–2.71)	9.9% (8.5–11.5)
Hypertension	21.91	39.02	2.48 (2.30–2.68)	23.4% (21.7–25.1)	1.91 (1.74–2.10)	17.9% (15.7–20.4)
Abdominal obesity (2 vs 1)†	33.40	30.21	1.36 (1.24–1.48)	–	1.12 (1.01–1.25)	–
Abdominal obesity (3 vs 1)†	33.32	46.31	2.24 (2.06–2.45)	33.7% (30.2–37.4)	1.62 (1.45–1.80)	20.1% (15.3–26.0)
All psychosocial‡	–	–	2.51 (2.15–2.93)	28.8% (22.6–35.8)	2.67 (2.21–3.22)	32.5% (25.1–40.8)
Vegetables and fruit daily*	42.36	35.79	0.70 (0.64–0.77)	12.9% (10.0–16.6)	0.70 (0.62–0.79)	13.7% (9.9–18.6)
Exercise*	19.28	14.27	0.72 (0.65–0.79)	25.5% (20.1–31.8)	0.86 (0.76–0.97)	12.2% (5.5–25.1)
Alcohol intake*	24.45	24.01	0.79 (0.73–0.86)	13.9% (9.3–20.2)	0.91 (0.82–1.02)	6.7% (2.0–20.2)
ApoB/ApoA1 ratio (2 vs 1)§	19.99	14.26	1.47 (1.28–1.68)	–	1.42 (1.22–1.65)	–
ApoB/ApoA1 ratio (3 vs 1)§	20.02	18.05	2.00 (1.74–2.29)	–	1.84 (1.58–2.13)	–
ApoB/ApoA1 ratio (4 vs 1)§	19.99	24.22	2.72 (2.38–3.10)	–	2.41 (2.09–2.79)	–
ApoB/ApoA1 ratio (5 vs 1)§	20.00	33.49	3.87 (3.39–4.42)	54.1% (49.6–58.6)	3.25 (2.81–3.76)	49.2% (43.8–54.5)
All above risk factors combined¶	–	–	129.20 (90.24–184.99)	90.4% (88.1–92.4)	129.20 (90.24–184.99)	90.4% (88.1–92.4)

The median waist/hip ratio was 0.93 in cases and 0.91 in controls ( $p < 0.0001$ ), and the median ApoB/ApoA1 ratio was 0.85 in cases and 0.80 in controls ( $p < 0.0001$ ). Percentage of controls with four or five factors positive is 22.2% compared with 29.2% in cases. \*PARs for smoking, abdominal obesity, and ApoB/ApoA1 ratio are based on a comparison of all smokers vs never, top two tertiles vs lowest tertile, and top four quintiles vs lowest quintile. For protective factors (diet, exercise, and alcohol), PARs are provided for the group without these factors. †Top two tertiles vs lowest tertile. ‡A model-dependent index combining positive exposure to depression, perceived stress at home or work (general stress), low locus of control, and major life events, all referenced against non-exposure for all five factors. §Second, third, fourth, or fifth quintiles vs lowest quintile. ¶The model is saturated, so adjusted and unadjusted estimates are identical for all risk factors. The odds ratio of 129.20 is derived from combining all risk factors together, including current and former smoking vs never smoking, top two tertiles vs lowest tertile of abdominal obesity, and top four quintiles vs lowest quintile of ApoB/ApoA1. If, however, the model includes only current smoking vs never smoking, the top vs lowest tertile for abdominal obesity, and the top vs lowest quintile for ApoB/ApoA1, the odds ratio for the combined risk factors increases to 333.7 (99% CI 230.2–483.9).

Table 3: Risk of acute myocardial infarction associated with risk factors in the overall population



**Figure 1: Odds of myocardial infarction according to number of cigarettes smoked and ApoB/ApoA1 ratio**  
Note the doubling scale on the y axis for both figures.

42.3 (33.2–54.0), and the PAR for these four risk factors together (top four quintiles of ApoB/ApoA1 ratio vs lowest quintile) was 75.8% (99% CI 72.7–78.6). Addition of abdominal obesity (top two tertiles vs lowest tertile) further increased the PAR to 80.2% (77.5–82.7).

Figure 3 shows the effects of multiple risk factors on reduced risk of acute myocardial infarction associated with healthy lifestyles. Daily consumption of fruit and vegetables and regular physical activity conferred an odds ratio of 0.60 (99% CI 0.51–0.71). Further, if an individual avoided smoking, the odds ratio would be 0.21 (0.17–0.25; figure 3), suggesting that modification of these aspects of lifestyle could potentially reduce the risk of an acute myocardial infarction by more than three-quarters compared with a smoker with a poor lifestyle.

Incorporation of all nine independent risk factors (current or former smoking, history of diabetes or

hypertension, abdominal obesity, combined psychosocial stressors, irregular consumption of fruits and vegetables, no alcohol intake, avoidance of any regular exercise, and raised plasma lipids) indicates an odds ratio of 129.20 (99% CI 90.24–184.99; table 3), compared with not having any of these risk factors. Substituting the odds ratios for current smoking, the extremes of abdominal obesity (top vs lowest tertile) and ApoB/ApoA1 ratio (top vs lowest quintile) increases the combined effect of all nine risk factors to 333.7 (99% CI 230.2–483.9; figure 2). This represents a PAR of 90.4% (99% CI 88.1–92.4), suggesting that these risk factors account for most of the risk of acute myocardial infarction in our study population. In view of the overlap in the effect of the nine risk factors, most of the PAR could be accounted for by a combination of various risk factors, as long as they included smoking and the ApoB/ApoA1 ratio (PAR for their combination is 66.8% [99% CI 62.8–70.6]). The estimate of the combined effect of all nine risk factors is derived from a model, since very few individuals had zero risk factors or all nine risk factors. However, confidence that the majority of risk is indeed accounted for by these risk factors is lent support by the fact that of the 18 708 individuals with complete data on all risk factors, 43 controls and 24 cases had no risk factors and 49 cases and 11 controls had eight or more. Also, just five risk factors (smoking, lipids, hypertension, diabetes, and obesity), which a large proportion of individuals had, accounted for about 80% of the PAR.

#### Risk in men and women

Figure 4 presents odds ratios and PARs for risk of acute myocardial infarction in men and women. Similar odds ratios were recorded in women and men for the association of acute myocardial infarction with smoking, raised lipids, abdominal obesity, composite of psychosocial variables, and vegetable and fruit consumption. However, the increased risk associated with hypertension and diabetes, and the protective effect of exercise and alcohol, seemed to be greater in women than in men (figure 4).

Table 4 also shows PARs by sex for the various risk factors, adjusted for age and region only and the fully adjusted model. In men, smoking was associated with 42.7% of the PAR for acute myocardial infarction compared with 14.8% in women in the fully adjusted model. Abnormal lipids had the highest PAR in both men (49.5%) and women (47.1%), with high contributions from psychosocial risk factors (28.8% vs 45.2%) and abdominal obesity (19.7% vs 18.7%). Hypertension contributed to PAR in women to a greater extent (29.0%) compared with men (14.9%), partly because of a higher prevalence of hypertension in women who were about a decade older. Collectively, all nine risk factors accounted for 90% of the PAR in men and 94% in women (table 4).

### Risk by age


Smoking, adverse lipid profile, hypertension, and diabetes had a greater relative effect on risk of acute myocardial infarction in younger than older individuals (table 5). Overall, abnormal lipids was the most important risk factor with respect to PAR in both young and old individuals (table 5). Collectively, the nine risk factors accounted for a significantly greater ( $p<0.0001$ ) PAR in younger than older individuals; these patterns were consistent in males and females.

### Regional and ethnic variations in importance of risk factors

When the odds ratio (adjusted for age, sex, smoking, and geographic region) for association of acute myocardial infarction with a risk factor is around 2 or more, eg, for smoking, lipids, hypertension, diabetes, abdominal obesity, and the combined psychosocial index, subgroup analyses are likely to be fairly robust. We recorded a clear, significant, and consistent excess risk of acute myocardial infarction associated with these risk factors in most regions of the world and in every ethnic group (figures 5–10). By contrast, when odds ratios were weaker (0.70–1.50; alcohol consumption, exercise, or diet), greater variability was noted across regions (data not shown). This apparent variability could be attributable to chance, because subgroup analyses are likely to be less reliable when smaller overall differences are subdivided across multiple subsets of the populations. Similar results were noted for analyses across various subgroups defined by ethnic origin, with consistent and clear excess risks being reported for tobacco use, abnormal lipids, history of hypertension, diabetes, and abdominal obesity (data not shown).

### Population attributable risk by geographic region


Table 4 also presents overall PARs and values by sex across different geographic regions. In all regions, the nine risk factors account for between three-quarters and virtually all the PAR for acute myocardial infarction. The relative importance of every risk factor varied, and was largely related to its prevalence. However, raised lipids, smoking, and psychosocial factors were the most important risk factors in all regions in the world. It is noteworthy that in western Europe, North America, and Australia and New Zealand (representing high-income countries) and southeast Asia (mostly middle-income countries), abdominal obesity was associated with a PAR greater than that associated with smoking. A similar pattern was seen for Africa, but most of our data are drawn from South Africa, which is a middle-income country. However, obesity was less important in other parts of the world, where it is less prevalent. For example, obesity accounted for only 5.5% of the PAR in China compared with 35.8% for smoking (where 41% of



**Figure 2: Risk of acute myocardial infarction associated with exposure to multiple risk factors**


Smk=smoking. DM=diabetes mellitus. HTN=hypertension. Obes=abdominal obesity. PS=psychosocial. RF=risk factors. Note the doubling scale on the y axis. The odds ratios are based on current vs never smoking, top vs lowest tertile for abdominal obesity, and top vs lowest quintile for ApoB/ApoA1. If these three are substituted by current and former smoking, top two tertiles for abdominal obesity and top four quintiles for ApoB/ApoA1, then the odds ratio for the combined risk factor is 129.20 (99% CI 90.24–184.99).

male and 4% of female controls smoked). Subdividing the population by ethnic origin, these nine risk factors accounted for a very high proportion of the PAR in every ethnic group (Europeans, 86%; Chinese, 90%; south Asians, 92%; black Africans, 92%; Arabs, 93%; and Latin Americans, 90%).



**Figure 3: Reduced risk of acute myocardial infarction associated with various risk factors**

Smk=smoking. Fr/vg=fruits and vegetables. Exer=exercise. Alc=alcohol. Note the doubling scale on the y axis. Odds ratios are adjusted for all risk factors.



**Figure 4:** Association of risk factors with acute myocardial infarction in men and women after adjustment for age, sex, and geographic region

For this and subsequent figures, the odds ratios are plotted on a doubling scale. Prevalence cannot be calculated for psychosocial factors because it is derived from a model.

### Consistency of results

Subgroup analyses with both types of controls (hospital-based and community-based) showed consistent odds ratios for current smoking (hospital-based 3.1 vs community-based 2.8), for the top quintile versus lowest quintile of lipids (4.2 vs 3.9), for diabetes (2.7 vs 3.4), for hypertension (2.1 vs 3.0), for abdominal obesity (1.7 vs 1.9), for psychosocial factors (1.6 vs 1.5), for consumption of fruits (0.78 vs 0.93) and vegetables (0.78 vs 0.83), for regular physical activity (0.79 vs 0.79), and for alcohol use (0.79 vs 0.86).

583 cases of acute myocardial infarction subsequently died in hospital. Odds ratios for fatal myocardial infarction associated with various risk factors were similar to those overall—smoking (2.1 for fatal myocardial infarction vs 3.0 overall), diabetes (4.0 vs 3.1), hypertension (2.4 vs 2.5), abdominal obesity (1.5 vs 2.2), and lipids (2.6 vs 3.9).

### Family history

Family history of coronary heart disease was associated with an odds ratio of 1.55 (99% CI 1.44–1.67), adjusted for age, sex, smoking, and geographic region. Adjustments for the nine previously described risk factors slightly reduced the odds ratio to 1.45 (1.31–1.60). The PAR was 12.0% (99% CI

9.2%–15.1%), which fell to 9.8% (7.6–12.5) after full adjustment. However, when family history is added to the information from other nine risk factors, the overall PAR rose from 90.4% to only 91.4%, indicating that although family history is an independent risk factor for myocardial infarction, most of the associated risk burden can be accounted for through the other risk factors studied. Family history seemed to be slightly more important in young (PAR 14.8% [11.7–18.5]) compared with old individuals (10.4% [8.3–13.0]).

### Repeat measures

Repeat measures of risk factors were made in 279 controls at a median interval of 409 days. The agreement rates for smoking (Cohen's kappa<sup>16</sup>  $\kappa=0.94$ ), history of diabetes ( $\kappa=0.90$ ), ApoB/ApoA1 (intraclass correlation=0.74), hypertension ( $\kappa=0.82$ ), depression ( $\kappa=0.44$ ), abdominal obesity (intraclass correlation=0.68), regular physical activity ( $\kappa=0.56$ ), and consumption of fruits ( $\kappa=0.66$ ), vegetables ( $\kappa=0.52$ ), and alcohol ( $\kappa=0.52$ ) were high to moderate. These data suggest that the association of myocardial infarction with smoking and diabetes is closer to the real effect, whereas the association of other risk factors measured with greater variability are probably underestimates due to regression-dilution bias.<sup>17</sup>

Region	Lifestyle factors					Other risk factors					
	Smoking (%)	Fruits and vegetables (%)	Exercise (%)	Alcohol (%)	All lifestyles (%)	Hypertension (%)	Diabetes (%)	Abdominal obesity (%)	All psychosocial (%)	Lipids (%)	All nine risk factors (%)
<b>Men</b>											
Western Europe	39.0	13.3	37.7	14.1	69.6	20.5	12.8	68.6	23.7	36.7	92.0
Central and eastern Europe	40.4	7.6	-0.4	10.4	48.9	15.9	5.8	31.7	-0.9	38.7	71.9
Middle East	51.4	5.8	1.9	-2.7	50.7	5.8	13.1	23.9	37.2	72.7	94.8
Africa	45.2	-4.4	15.9	24.1	63.7	26.8	11.6	60.4	33.8	73.7	97.9
South Asia	42.0	16.0	25.5	-5.7	58.1	17.8	10.5	36.0	13.9	60.2	88.4
China	45.3	15.1	16.6	4.2	63.7	19.9	7.9	4.9	32.0	41.3	88.8
Southeast Asia and Japan	39.2	8.5	31.4	24.6	69.6	34.3	19.1	57.9	26.9	68.7	93.7
Australia and New Zealand	46.1	8.0	20.6	11.2	61.0	18.3	5.6	49.5	31.6	48.7	87.5
South America	42.4	7.1	27.6	-7.4	57.7	28.1	9.7	35.2	36.1	41.6	86.1
North America	30.9	22.4	24.7	6.6	53.9	13.9	6.1	64.7	63.7	60.0	100
Overall 1	44.0	10.3	22.9	10.5	63.8	19.5	10.1	32.1	25.3	53.8	89.8*
Overall 2	42.7	11.7	9.3	5.1	56.5	14.9	8.0	19.7	28.8	49.5	89.8*
<b>Women</b>											
West Europe	11.1	8.4	38.3	34.2	65.2	25.9	21.0	50.6	67.1	47.9	97.1
Central and eastern Europe	13.1	12.8	42.7	29.9	65.4	42.7	15.7	20.0	15.0	26.8	86.1
Middle East	8.1	15.9	39.1	59.0	80.3	30.1	30.3	38.9	77.4	63.3	99.4
Africa	27.6	21.0	-37.9	28.8	61.2	35.1	27.5	54.6	54.9	74.6	93.3
South Asia	7.1	30.6	45.0	26.0	59.8	28.9	20.5	48.7	29.2	52.1	99.3
China	12.5	23.6	33.5	35.8	78.6	27.6	15.0	6.3	43.2	48.3	93.6
Southeast Asia and Japan	14.8	19.9	32.8	69.5	84.5	56.3	29.2	58.0	27.0	64.5	96.5
Australia and New Zealand	40.7	15.8	33.6	47.4	80.0	37.0	11.7	67.2	17.2	14.9	†
South America	25.8	5.9	27.4	44.1	71.8	47.9	22.2	63.0	37.8	59.3	96.1
North America	25.3	12.8	27.2	73.3	86.9	30.2	12.4	44.5	32.7	32.2	†
Overall 1	15.8	17.8	37.3	46.9	75.0	35.8	19.1	35.9	40.0	52.1	94.1*
Overall 2	14.8	19.1	27.1	22.1	60.6	29.0	16.1	18.7	45.2	47.1	94.1*
<b>Men and women</b>											
West Europe	29.3	12.4	38.4	18.7	67.6	21.9	15.0	63.4	38.9	44.6	93.9
Central and eastern Europe	30.2	10.2	11.3	12.9	49.6	24.5	9.1	28.0	4.9	35.0	72.5
Middle East	45.5	7.3	4.2	-1.0	47.6	9.2	15.5	25.9	41.6	70.5	95.0
Africa	38.9	4.8	10.1	26.6	63.4	29.6	16.7	58.4	40.0	74.1	97.4
South Asia	37.4	18.3	27.1	-5.5	56.6	19.3	11.8	37.7	15.9	58.7	89.4
China	35.9	18.0	20.3	5.7	62.3	22.1	10.0	5.5	35.4	43.8	89.9
Southeast Asia and Japan	36.2	11.2	31.4	27.9	69.9	38.4	21.0	58.0	26.7	67.7	93.7
Australia and New Zealand	44.8	11.1	23.8	18.6	66.0	22.6	7.2	61.3	28.9	43.4	89.5
South America	38.3	6.6	27.6	-3.7	56.6	32.7	12.7	45.5	35.6	47.6	89.4
North America	26.1	19.8	25.6	25.5	59.9	19.0	8.0	59.5	51.4	50.5	98.7
Overall 1	36.4	12.9	25.5	13.9	62.9	23.4	12.3	33.7	28.8	54.1	90.4*
Overall 2	35.7	13.7	12.2	6.7	54.6	17.9	9.9	20.1	32.5	49.2	90.4*

PAR estimates in women in some countries are based on small numbers and so they are less reliable. Overall 1= adjusted for age, sex, and smoking, Overall 2=adjusted for all risk factors. An extended version of this table with 99% CIs is shown in webtable 3 (<http://image.thelancet.com/extras/04art8001webtable3.pdf>). \*Saturated model, no difference between adjusted and unadjusted models. †Non-estimable.

**Table 4: PARs associated with nine risk factors in men and women by geographic region**

## Discussion

Our study shows that nine easily measured and potentially modifiable risk factors account for an overwhelmingly large (over 90%) proportion of the risk of an initial acute myocardial infarction. The effect of these risk factors is consistent in men and women, across different geographic regions, and by ethnic group, making the study applicable worldwide. The effect of the risk factors is particularly striking in young men (PAR about 93%) and women (about 96%), indicating that most premature myocardial infarction is preventable. Worldwide, the two most important risk factors are smoking and abnormal lipids. Together they account for about two-thirds of the PAR of an acute myocardial infarction. Psychosocial factors, abdominal obesity, diabetes, and hypertension were the next most important risk factors in men and women, but their

relative effect varied in different regions of the world. The usual measure of obesity (body-mass index) showed a modest relation with acute myocardial infarction but was not significant when abdominal obesity was included in the analysis.

Both smoking and apolipoproteins showed a graded relation with the odds of a myocardial infarction, without either a threshold or a plateau in the dose response. In particular, smoking even five cigarettes per day increased risk. This finding suggests that there is no safe level of smoking and that if quitting is not possible, the risk of myocardial infarction associated with smoking could be significantly reduced by a reduction in the numbers smoked. The graded relation between ApoB/ApoA1 ratio across the deciles is consistent with findings of a Swedish study<sup>10</sup> and shows that most populations in the world (at least

	Both sexes		Men		Women	
	Young	Old	≤55 years	>55 years	≤65 years	> 65 years
<b>Odds ratios for relative effect of risk factors (99% CI)</b>						
<b>Lifestyle factors</b>						
Smoking	3.33 (2.86–3.87)	2.44* (2.10–2.84)	3.33 (2.80–3.95)	2.52 (2.15–2.96)	4.49 (3.11–6.47)	2.14 (1.35–3.39)
Fruit and vegetables	0.69 (0.58–0.81)	0.72 (0.61–0.85)	0.72 (0.59–0.88)	0.77 (0.64–0.93)	0.62 (0.44–0.87)	0.55 (0.38–0.80)
Exercise	0.95 (0.79–1.14)	0.79 (0.66–0.94)	1.02 (0.83–1.25)	0.79 (0.66–0.96)	0.74 (0.49–1.10)	0.75 (0.46–1.22)
Alcohol	1.00 (0.85–1.17)	0.85 (0.73–1.00)	1.03 (0.87–1.23)	0.86 (0.73–1.01)	0.74 (0.41–1.31)	0.83 (0.49–1.42)
All four lifestyle factors	0.20 (0.14–0.27)	0.20† (0.15–0.27)	0.23 (0.16–0.33)	0.21 (0.15–0.29)	0.07 (0.03–0.18)	0.16 (0.06–0.41)
Hypertension	2.24 (1.93–2.60)	1.72 (1.52–1.95)	1.99 (1.66–2.39)	1.72 (1.49–1.98)	2.94 (2.25–3.85)	1.82 (1.39–2.38)
Diabetes	2.96 (2.40–3.64)	2.05* (1.71–2.45)	2.66 (2.04–3.46)	1.93 (1.58–2.37)	3.53 (2.49–5.01)	2.59 (1.78–3.78)
Abdominal obesity	1.79 (1.52–2.09)	1.50 (1.29–1.74)	1.83 (1.52–2.20)	1.54 (1.30–1.83)	1.58 (1.14–2.20)	1.22 (0.88–1.70)
Psychosocial	2.87 (2.19–3.77)	2.43 (1.86–3.18)	2.62 (1.91–3.60)	2.45 (1.82–3.29)	3.92 (2.26–6.79)	2.31 (1.22–4.39)
High ApoB/ApoA1 ratio	4.35 (3.49–5.42)	2.50* (2.05–3.05)	4.16 (3.19–5.42)	2.51 (2.00–3.15)	4.83 (3.19–7.32)	2.48 (1.60–3.83)
All risk factors other than smoking	101.86 (61.22–169.46)	43.24* (26.96–69.37)	59.06 (32.25–108.14)	38.88 (22.95–65.86)	473.43 (158.34–1415.5)	67.49 (21.39–212.90)
All nine risk factors including smoking‡	216.47 (126.67–369.94)	81.99* (50.02–134.40)	129.19 (68.60–243.28)	76.25 (44.07–131.93)	1100.6§ (342.72–3534.2)	111.45 (32.59–381.12)
<b>Population attributable risks (99% CI)</b>						
<b>Lifestyle factors</b>						
Smoking	40.7% (35.9 to 45.7)	33.1% (28.9 to 37.6)	52.0% (44.9 to 59.0)	39.0% (34.0 to 44.1)	20.8% (15.7 to 26.9)	8.2% (4.1 to 15.7)
Fruit and vegetables	16.9% (10.8 to 25.3)	11.9% (7.4 to 18.4)	15.7% (8.3 to 27.8)	10.1% (5.3 to 18.2)	18.4% (10.0 to 31.5)	18.7% (10.0 to 32.1)
Exercise	7.5% (0.7 to 46.9)	13.4% (5.4 to 29.7)	0.1% (0.0 to 100.0)	12.5% (4.4 to 30.6)	24.6% (6.8 to 59.2)	23.6% (4.3 to 67.8)
Alcohol	–4.1% (–19.8 to 11.6)	11.1% (4.7 to 23.9)	–9.1% (–25.1 to 6.9)	10.5% (4.3 to 23.6)	24.9% (3.3 to 76.3)	14.6% (0.5 to 84.6)
All four lifestyle factors	52.1% (39.5 to 64.4)	54.8% (46.2 to 63.1)	55.8% (42.1 to 68.7)	57.1% (48.4 to 65.4)	63.3% (36.8 to 83.6)	51.5% (21.7 to 80.3)
Hypertension	19.2% (16.0 to 22.8)	17.0% (14.0 to 20.5)	12.8% (9.4 to 17.1)	15.7% (12.7 to 19.4)	31.9% (25.7 to 38.6)	25.4% (17.1 to 35.8)
Diabetes	12.4% (10.3 to 14.9)	8.6% (6.9 to 10.7)	8.7% (6.6 to 11.5)	7.8% (6.0 to 10.1)	19.3% (15.1 to 24.5)	13.0% (8.9 to 18.5)
Abdominal obesity	24.8% (17.2 to 34.5)	18.1% (12.2 to 26.0)	23.4% (14.4 to 35.7)	18.3% (11.9 to 27.0)	24.9% (12.4 to 43.7)	11.8% (2.1 to 46.1)
Psychosocial	43.5% (32.2 to 55.6)	25.2% (16.0 to 37.2)	39.7% (25.4 to 56.0)	23.7% (13.9 to 37.4)	53.0% (35.4 to 69.9)	30.6% (10.6 to 62.1)
High ApoB/ApoA-1 ratio	58.9% (50.9 to 66.5)	43.6% (36.6 to 50.8)	59.7% (48.6 to 70.0)	45.3% (37.5 to 53.3)	56.1% (43.7 to 67.7)	36.3% (21.8 to 53.8)
All risk factors other than smoking	89.4% (84.7 to 92.7)	81.7% (76.4 to 86.1)	85.6% (77.7 to 91.0)	80.8% (74.8 to 85.7)	95.5% (90.0 to 98.0)	86.4% (70.8 to 94.3)
All nine risk factors including smoking	93.8% (90.9 to 95.8)	87.9% (84.1 to 90.8)	93.1% (88.9 to 95.8)	88.3% (84.4 to 91.4)	96.5% (92.0 to 98.5)	87.7% (73.1 to 94.9)

\*p<0.001 are only provided for the overall comparison. †These values differ slightly but appear similar because of rounding. ‡Based on combining current and former smokers vs never smokers, top two tertiles vs lowest tertile for abdominal obesity, and top four quintiles vs lowest quintile for ApoB/ApoA1 ratio. If, however, extreme exposures (current vs never, top vs lowest tertile for abdominal obesity, and top vs lowest quintile for ApoB/ApoA1 ratio) were included, the odds ratios for all risk factors for the young group increases to 756.0 and in the old group to 160.8. §Unstable estimate, should be interpreted cautiously.

Table 5: Importance of risk factors in young and old individuals

urban) have lipid abnormalities, which increase the risk of myocardial infarction. Since ApoB/ApoA1 ratio was the most important risk factor in all geographic regions in our study, a substantial modification of its population distribution is important for worldwide

reduction of myocardial infarction. This act will probably need a concerted effort, including both population-based strategies to shift the distribution and treatments targeted at people with the greatest abnormalities.





Figure 5: Risk of acute myocardial infarction associated with current or former smoking, overall and by region after adjustment for age and sex


W Eur=western Europe. CE Eur=central and eastern Europe. MEC=Middle East Crescent. Afr=Africa. S=South. HK=Hong Kong. SE=southeast. ANZ=Australia and New Zealand. N=North. Am=America.



**Figure 6:** Risk of acute myocardial infarction associated with ApoB/ApoA1 ratio (top vs lowest quintile), overall and by region after adjustment for age, sex, and smoking  
PAR is for the top four quintiles versus the lowest quintile.

Our data show that risks associated with the major risk factors (odds ratio of about 2 or greater on univariate analyses, such as smoking, abnormal lipids, psychosocial factors, hypertension, diabetes, and abdominal obesity) were consistently adverse in all regions of the world and in all ethnic groups. In particular, the odds ratios for these risk factors were qualitatively similar (although some quantitative differences were apparent), despite variations in prevalence for every risk factor in controls derived from different subpopulations. However, as expected, the PAR is affected both by the prevalence of the risk factor and the odds ratio. We are unaware of any other large study that has assessed whether risk factors have a similar or differing effect in many ethnic groups.

Our finding that most risk factors have directionally similar odds ratios in ethnic groups and countries differs from inferences reached by comparison of results of different studies, which used other methods.<sup>3,4</sup> Some of these researchers suggested that the effects of the major risk factors could vary qualitatively in different regions and ethnic groups, possibly because of inconsistent methodologies, differences in criteria used to recruit participants, variations in information obtained, and a fairly modest number of events in each study, thereby leading to imprecise estimates of risk that could have been exaggerated or diluted by the play of chance. Since we had more than 800 cases of acute myocardial infarction within every major ethnic group (other than black or coloured Africans), our results within most



**Figure 7:** Risk of acute myocardial infarction associated with self-reported hypertension, overall and by region after adjustment for age, sex, and smoking




Figure 8: Risk of acute myocardial infarction associated with self-reported diabetes, overall and by region after adjusting for age, sex, and smoking

ethnic groups are statistically robust. The number of cases of myocardial infarction in this study within every region or ethnic group is larger than in most previous studies, especially in those of non-European origin.

The prevalence of several risk factors varied substantially, especially when subdivided by sex. For example, smoking in female controls worldwide has a prevalence of only 9.25% compared with 33% in male controls. As a result, despite similar odds ratios in women and men, the PAR attributable to smoking varied greatly (16% in women and 44% in men). These data suggest that the overall approach to prevention of

coronary heart disease could be similar worldwide, but with varying emphasis in different subgroups (eg, sex and geographic region) on the basis of the prevalence of individual risk factors and economic and cultural factors. The above data also suggest that smoking cessation is very important in most male populations worldwide and in women in North and South America, western Europe, and Australia and New Zealand. By contrast, quitting smoking is currently less important for reducing acute myocardial infarction in women in most other geographic regions. However, if women in these countries start smoking they are likely to have a




Figure 9: Risk of acute myocardial infarction associated with abdominal obesity measured as waist/hip ratio (upper tertile vs lowest tertile), overall and by region after adjusting for age, sex, and smoking  
PARs are for top two tertiles vs lowest tertile.




Figure 10: Risk of acute myocardial infarction associated with the composite psychosocial index, overall and by region

substantial increase in rates of acute myocardial infarction attributable to smoking.

Hypertension and diabetes were associated with a greater odds ratio and PAR in women compared with men, but women with these factors were about a decade older than men. Further, the protective effects of exercise and alcohol consumption also seemed greater in women than in men. While the amplified effect of diabetes in women has been reported before,<sup>18</sup> we are not aware of similar data about the other three factors. Thus, even though significant interactions were noted between these risk factors and sex for the odds of myocardial infarction, it would be prudent to seek independent confirmation.

Known risk factors (generally smoking, hypertension, raised lipids, and diabetes) have sometimes been claimed to account for only about half the risk of a myocardial infarction. The origins of this claim are unclear.<sup>19</sup> Our analysis, which is based on traditional and some newly described risk factors, suggests that more than 90% of the risk of an acute myocardial infarction in a population can be predicted by the risk factors included in our study. Findings of several previous studies—in which fewer risk factors were measured (most large studies have not included apolipoproteins, psychosocial factors or abdominal obesity)—lend support to our observations. Stamler and colleagues<sup>20</sup> studied five US cohorts and categorised individuals on the basis of the presence of five factors (abnormal electrocardiogram, diabetes, smoking, cholesterol, and blood pressure). Those without any of these risk factors were judged to be in the low-risk category and had an 80–90% lower risk of coronary heart disease in every cohort compared with the rest of the population. Similar results were also reported in an analysis of the Göteborg population, in which individuals with low blood pressure and a low amount of cholesterol, who were also non-smokers, had an age-adjusted relative risk of 0.09,

which was much lower than that for the average population (relative risk 1.0) in the study.<sup>21</sup>

The importance of modifying risk factors is lent support by data from randomised trials—eg, blood-pressure lowering,<sup>22</sup> lipid lowering,<sup>23</sup> dietary modification<sup>24</sup>—or persuasive evidence of causality from observational studies<sup>25</sup> (eg, smoking cessation).<sup>26</sup> Some investigators have suggested that a pill that combines a statin, antihypertensive drugs, and aspirin, together with avoidance of smoking, could potentially reduce the risk of myocardial infarction by more than 80% to 90%.<sup>27</sup> These studies, along with INTERHEART, suggest that one of the major emphases in research should be to understand why currently known risk factors develop in some individuals and populations, and to identify approaches to prevent their development or reduce them. For example, understanding the mechanisms by which societal factors affect development of risk factors (urbanisation, food and tobacco policies, shifts in occupation from energy expending jobs to sedentary ones, and urban structure, etc) could lead to new approaches to prevent development of risk factors (primordial prevention),<sup>4</sup> which in turn could reduce coronary heart disease substantially.

Although the odds ratio for an acute myocardial infarction in people with a family history was about 1.5, the PAR rose from 90% with the nine potentially modifiable risk factors to 91% with the addition of family history. This finding suggests that a large part of the effect of family history might be mediated through known risk factors, which could be affected by both shared lifestyles and genetic factors rather than through independent pathways. Therefore, the main challenge in the next few decades will be a combination of discovering more effective strategies to substantially alter or prevent development of known risk factors by understanding the societal, environmental, and biological causes of the development of these factors.

One of the most important risk factors for acute myocardial infarction in our study was smoking, which accounts for about 36% of the PAR of acute myocardial infarction worldwide (and about 44% in men). Regular consumption of fruits and vegetables was associated with a 30% relative risk reduction. Thus, eating fruit and vegetables, taking exercise, and avoiding smoking could lead to about 80% lower relative risk for myocardial infarction. Our results are similar to the findings of the US Nurses Health Study,<sup>28</sup> which also indicated that lifestyle modification could potentially avoid more than three-quarters of the risks of coronary heart disease and strokes in women. These conclusions are also lent support by the results of the Lyon Heart Study,<sup>24</sup> which suggested that dietary modification by itself reduced the risk of coronary heart disease by about half in patients with coronary disease. Our data suggest that lifestyle modification is of substantial importance in both men and women, at all ages, in individuals from all geographic regions of the world, and in those belonging to all major ethnic groups. Therefore, smoking avoidance, increased consumption of fruits and vegetables, and moderate activity (along with lipid lowering) should be the cornerstone of prevention of coronary heart disease in all populations worldwide.

We also recorded an additional protective effect of moderate alcohol consumption (PAR 7%). The effect seemed to be surprisingly large in women, in whom absence of regular alcohol consumption accounted for about 22% of PAR, but with wide confidence limits (−4.9 to 60.8). This finding suggests that the best estimate of PAR attributable to alcohol consumption in women is probably closer to the overall estimate of 7%. Promotion of the consumption of moderate alcohol to prevent myocardial infarction might also not be acceptable to many populations, for cultural or religious reasons, and might increase the proportion of heavy drinkers and thereby enhance the risk of other diseases such as strokes, some cancers, cirrhosis of the liver, or injuries. The overall PAR without alcohol included in the model is 89.7%; adding alcohol increases it by less than 1% because of the substantial overlap in contributions of other risk factors. Therefore, advice about alcohol use could be best customised to individuals depending on their social, cultural, and religious backgrounds and the overall effect on their health.

Our study has several potential limitations. First, a case-control design is potentially open to confounding if there is differential ascertainment of risk factors between cases and controls. We minimised this factor by using standardised methods for data collection in both cases and controls. The inclusion of incident (first) acute myocardial infarction cases reduces the possibility that individuals with previous cardiovascular disease might have substantially altered lifestyles or risk factor levels before this event. Further, the odds ratios associated with all major risk factors—eg, smoking, lipids, diabetes, and

hypertension—in INTERHEART is similar to that reported in other cohort studies in western populations. We attempted to minimise biases in the selection of controls by excluding individuals in whom the risk factors that we were interested in studying were implicated as being protective or harmful. Reanalysis of our data by the two types of controls—hospital-based and community-based—did not alter our results. Our results are qualitatively similar for most risk factors in all regions of the world, providing internal replication. Any selection biases are unlikely to have been similarly prevalent across a large number of centres in 52 countries. Therefore, we think that there is little material bias in our results because of the use of a case-control study design.

Second, whereas some of the risk factors were ascertained or measured with high accuracy (eg, smoking), others (eg, history of diabetes or hypertension) were based on history and therefore ascertained with some error. The actual blood pressure value after a myocardial infarction is potentially confounded because it might have fallen in some patients because of the infarction itself or as a result of the drugs used in the management of the acute phase. Similarly, glucose concentrations rise with acute myocardial infarction (stress hyperglycaemia) and are therefore not an indication of earlier levels. We obtained blood samples for HbA1c but these are yet to be analysed. Therefore, our approach to diagnosis of hypertension or diabetes might have led to misclassification in some individuals with respect to their risk-factor status. These misclassifications would tend to underestimate the real relation between these risk factors and outcomes. Analysis of our control group data indicates a relation between the reported prevalence of hypertension in every centre and measured blood pressure in controls (data not shown), suggesting that there is some validity in using self-reports of hypertension as a surrogate for measured blood pressure. However, the absence of available blood pressure and glucose values could have underestimated their importance.

Third, the correlations between repeated measures of several variables (eg, diet or physical activity) many months apart is only moderate. Methods to correct for measurement error and regression dilution bias for one risk factor have been described;<sup>17</sup> however, we are not aware of methods that adjust for several risk factors simultaneously. However, if correction for regression-dilution bias could have been made it could further increase the odds ratios for most risk factors, which in turn would increase the overall PAR accounted for by the nine risk factors that we measured. This outcome means that the nine risk factors measured in this study probably account for virtually all the PAR for myocardial infarction in the population included in this study.

Fourth, our data are based on hospital-based patients with acute myocardial infarction and matched controls (mainly from urban areas) and are therefore unlikely to

reflect the population prevalence of risk factors in an entire country or region. This fact could potentially have an effect on our estimates of PAR. However, the key to ensuring internal validity of the study is to recruit cases and controls from the same population, which has been our emphasis. Therefore, our estimates of PAR should be regarded as providing reliable information about the specific population enrolled into our study. Nevertheless, when data are available from several countries (eg, for smoking), the rates in controls in INTERHEART closely match published reports for similar age-groups and sexes. As a result, our overall conclusions that the risk factors measured in this study account for most of the risk of acute myocardial infarction is probably broadly applicable. In view of the consistency of our data, the odds ratios from the present study could be applied to other populations and their PAR can then be derived by using population-specific prevalence rates of specific risk factors.

Fifth, although the effects of individual risk factors and combinations of four or five of them are reasonably robust, our estimates of the effect of all nine is model-dependent because very few individuals have eight or nine risk factors or, conversely, none. However, crude examination of the extremes of risk-factor distribution, and the fact that just five risk factors (smoking, lipids, hypertension, diabetes, and obesity) for which we have a sizeable number of individuals predicts about 80% of the PAR, suggests that our model-based estimates are reasonably valid.

Our study has several strengths. First, the case-control study has several advantages over other designs, especially a cohort study. It allows efficient enrolment of large numbers of cases and hence greater statistical power, rapid and cost-effective study conduct, and enhances the ability to recruit a large number of cases occurring at young ages, in whom disease association might be stronger. Second, our study included several risk factors that have previously not been assessed with conventional risk factors, including apolipoproteins (ApoB/A1 ratio), which might be the best marker of the balance of atherogenic and antiatherogenic particles,<sup>10</sup> psychosocial factors,<sup>7</sup> and measures of abdominal obesity, all of which have added substantial information to the other commonly studied risk factors. Third, the large size of the study provides high power and precision in estimates both overall and in subgroups. Fourth, the inclusion of large numbers of individuals from all regions of the world and multiple ethnic groups makes our study results broadly applicable.

In conclusion, our study has shown that nine easily measured risk factors are associated with more than 90% of the risk of an acute myocardial infarction in this large global case-control study. These results are consistent across all geographic regions and ethnic groups of the world, men and women, and young and old. Although priorities can differ between geographic regions because of variations in prevalence of risk

factors and disease and economic circumstances, our results suggest that approaches to prevention of coronary artery disease can be based on similar principles throughout the world. Therefore, modification of currently known risk factors has the potential to prevent most premature cases of myocardial infarction worldwide.

#### Contributors

S Yusuf initiated the INTERHEART study, supervised its conduct and data analysis and had primary responsibility for writing this paper. S Ounpuu coordinated the worldwide study and reviewed and commented on drafts. S Hawken did all data analyses and reviewed and commented on drafts. All other authors coordinated the study in their respective countries and provided comments on drafts of the manuscript.

#### INTERHEART project office staff (Population Health Research Institute, Hamilton Health Sciences and McMaster University, Hamilton, ON, Canada)

**Coordination and Data Management**—S Yusuf (principal investigator), S Ounpuu (project officer), P Montague, J Keys, S Rangarajan (study coordinators), S Hawken, A Negassa (statisticians), C Wright, R Mayhew, L Westfall, P Mackie, C Cuvay, D Cunningham, K Picken, S Anand, H Gerstein.

**Core Laboratory**—M McQueen, K Hall, J Keys.

#### INTERHEART national coordinators and investigators

**Argentina**—R Diaz\*, E Paolasso\*, M Ciruzzi\*, R Nordaby, C H Sosa, H L Lucardi, E M Marzetti, J A Piasentin, M Amuchastegui, C Cuneo, G Zapata, J O Bono, E M Marxetti, A Caccavo, R A A Guerrero, R Nordaby, C Dizeo; **Australia and New Zealand**—D Hunt\*, J Varigos\*, M Sallaberger, G Aroney, P Hicks, D Campbell, J Amerena, W Walsh, G Nelson, D Careless, J Durech, K Gunawardane, A Thomson, D Owensby, A Buncle, B Watson, H Ikram, M Audeau, M Hills, D Rosen, J Rankin, A Tunesi, J Sampson, K Roberts, A Hamer, L Roberts, B Singh, C Singh, R Hendriks, G Tulloch, A Hill, D Rees, C Hall; **Bahrain**—M A Halim\*, M I Amin; **Bangladesh**—S Haque\*, G M Faruque, M Hossain; **Benin Republic**—H Agboton\*; **Botswana**—C Onen\*; **Brazil**—A Avezum\*, L Piegas\*, G Reis, J P Esteves, L C Bodanese, A R Junior, J A M Neto, R F Ramos, E S Silva, A Labrunie, A C C Carvalho, I P Filho, D C D Albuquerque, J A M Abrantes, O C E Gebara, P E Leaes, S S Xavier, R L Marino, O Dutra; **Cameroon**—W Muna\*, K N Blackett\*, **Canada**—K Teo\*, S Yusuf\*, S, Ounpuu, Y K Chan, C Joyner, A D Kitching, A Morris, A Panju, H Lee, I Bata, B May, J G Hiscock, P V Greenwood, W Tymchak, M Natarajan; **Chile**—F Lanas\*, D Standen, A Lanas, C Santibanez, P Soto, S Potthoff, R Soto, E Mercadal; **China**—L S Liu\*, L X Jiang\*, Q Chen, N Sun, J Wang, L Zhang, X Li, H Chen, Z Li, Y Shi, J Meng, D Wang, Z Liu, J Yan, W Zhang, Z Wan, Y Xu, X Li, L Wang, Q Fu, K Zhu, Y Wang, X Wang, F Wang, R Liu, R Zhao, K Sun, S Yue, Y Zhang, X Zhao, L Zhao, C Zhou, X Meng, A Liu, X Li, R Wu, R Xiao, X Zhang, G Wan, X Chen, Z Ge, T Zhou, Y Li, X Zhou, F Zhang; **Colombia**—L E Bautista\*, J P Casas, C A Morillo, E Ardilla, I A Arenas, M Lindarte, D Aristizabal; **Croatia**—Z Rumboldt\*, M Rumboldt, V Carevic; **Czech Republic**—P Widimsky\*, M Branny, P Gregor; **Egypt**—M S Khedr\*, M G Abdel-Aziz, S Abdel-Kader, M A Abdel-Moneim, F El-Demerdash, A El-Said, S Gharib, M Hassanin, F Abdel-Hamid Maklady, I Sadek; **Germany**—H J Rupprecht\*, T Wittlinger\*, P Schuster, A Schmidt, H R Ochs; **Greece**—N Karatzas\*, A Pipilis\*, D Symeonides, K Karidis, T H Tsaknakis, K Kyfinidis; **Guatemala**—M Luna\*, A M Arroyo, G Sotomora, M A Rodas, L Velasquez, A Ovando; **Hong Kong**—J Sanderson\*, W Y Ma, S Chan; **Hungary**—M Keltai\*, F Szaboki, P Karpati, K Toth, S Timar, E Kalo, A Janosi, M Rusznak, D Lehoczy, J Tarjan, E Kiss, E Sitkei, P Valyi, I Edes; **India**—P Pais\*, K S Reddy\*, P P Joshi, D Prabhakaran, D Xavier, A Roy, L Ramakrishnan; **Iran**—M R M Hasani\*, S H Mirkhani\*; **Israel**—D Halon\*, B Lewis\*; **Italy**—M G Franzosi\*, G Tognoni\*, E Gardinale, M Vilella, M Mennuni, F Saiu, G Pettinati, S Ciricugno, S Antonaci, L Carrieri, E Balli, D Bicego, M G DelleDonne, A Ottaviano, L Moretti, C Melloni, G Melandri, E Carbonieri, A Vetrano; **Japan**—M Hori\*, H Sato, K Fujii; **Kenya**—E N Ogola\*, P Wangai; **Kuwait**—M Zubaid\*, W Rashed; **Malaysia**—C C Lang\*, W A W Ahmad, A Kadirvelu, R Zambahari; **Mexico**—M A Ramos-Corralles\*, **Mozambique**—A Damasceno\*, **Nepal**—M R Pandey\*, G Baniya, A Sayami, M Kiratee, B Rawat, M Pandey, J Gurung; **Netherlands**—R Peters\*, D C G Basart, P N A Bronzwaer, J A Heijmeriks, H R Michels, G M A Pop; **Nigeria**—K K Akinroye\*; **Pakistan**—K Kazmi\*, M A Memon, S K Memon, S Nishar, A Badar, M A Mattu, M Yakub, K Soomra, J Khatri, A M A Faruqui, S I Rasool, A Samad; **Philippines**—AL Dans\*, F Q Punzalan, M V C Villarruz, B T Mendoza, C S Recto III, B R Tamesis, V Mendoza, D J Torres, D D Morales, I Ongtengco; **Poland**—A Budaj\*, L Ceremuzynski\*, S Stec, J Gorny, K Prochnewska, T K Urbanek, P Wojewoda, M Pawlowski, K Religa, R Klabisz, H Latocha, M Szpajer, K Cymerman, M Laniec,

H Danielewicz, M Ogorek, D Kopcik, A Baranowska, A Kozlowski, M Blachowicz, A Jedrzejewski, T Waszyrowski, Z Zielinski, K Janik, M Tomzynski, M Mytnik, A Maziarz, G Rembelska, M Piepiorka, K Debkowska, T Krynski, B S Szczeklik, W Krasowski, M Rozwodowska, P Miekus, J Surwilo, S Malinowski, J Hybel, M Michalak, P Achremczyk, J Majchrzak, W S Korombel, M Dobrowolski, J Gessek; *Portugal*—J Morais\*; *Qatar*—A A Gehani\*, H A Hajar\*, M M Almwola, A A Omer; *Russia*—E G Volkova\*, S U Levashov, O M Filatova, U I Evchenko, G S Malkiman, N N Karaulovskaya; *Seychelles*—J Panovsky\*; *Singapore*—B A Johan\*, A Cheng, K S Ng, Y L Lim, K H Neoh, K S Tan, L N Lum, Y Cheah; *South Africa*—P Commerford\*, K Steyn\*, B Brown, F Martiz, P Raubenheimer, A Aboo, J D Marx, E Batiste, K Sliwa, P Sareli, C Zambiakides; *Spain*—V Valentin\*, J Ferriz, A Rovira, A Rodriguez-Llorian, E Cereijo, T Monzo, N Alonso, E Gomez-Martinez, A Lopez-Perez; *Sri Lanka*—S Mendis\*, T Jayalath; *Sultanate of Oman*—A T A Hinai\*, M O Hassan; *Sweden*—A Rosengren\*; *Thailand*—C Sitthi-Amorn\*, S Chaitiraphan, K Bhuripanyo, S Srimahachota, M Anukulwutipong, P Loothavorn, P Sritara, S Tanomsup, J Tantitham, P Tatsanavivat, T Yipinsoi, W Jintapakorn, W Puavilai, B Koanantakul, P Kasemsuwan, C Supanantareauk, C Pearnsonboon, B Saejueng, K Jeamsomboon, A Sukontasup; *United Arab Emirates*—W Almahmeed\*, L O Abdelwareth, A Bokhari, N S Rao, S Bakir, A M Yusufali, E Hatou, Q Zaidi; *UK*—K Fox\*, M Flather\*; *USA*—J Probstfield\*, R Freeman, J Mathew, M Schweiger, W Johnson, E Ofili; *Zimbabwe*—J Chifamba\*, A J G Hakim.

\*National coordinator.

#### Conflict of interest statement

We declare that we have no conflict of interest.

#### Acknowledgments

We thank Gilles Dagenais, Sonia Anand, Hertzel Gerstein, and Janet Wittes for helpful comments on the manuscript; Judy Lindeman for secretarial assistance; and the staff at the core laboratories in Hamilton, Canada, and Beijing, China for their excellent efforts. Further, we thank WHO and the World Heart Federation for their endorsement, and our friends and colleagues for various forms of help that led to the successful completion of this global study. SY is supported by an endowed chair of the Heart and Stroke Foundation of Ontario and a Senior Scientist Award from the Canadian Institutes of Health Research (CIHR). SO held a Heart and Stroke Foundation of Canada Fellowship and a Canadian Institutes of Health Research Senior Research Fellowship during this study. The INTERHEART study was funded by the Canadian Institutes of Health Research, the Heart and Stroke Foundation of Ontario, the International Clinical Epidemiology Network (INCLEN), and through unrestricted grants from several pharmaceutical companies (with major contributions from AstraZeneca, Novartis, Hoechst Marion Roussel [now Aventis], Knoll Pharmaceuticals [now Abbott], Bristol Myers Squibb, King Pharma, and Sanofi-Syhelabo), and by various national bodies in different countries (see webappendix2; <http://image.thelancet.com/extras/04art8001webappendix2.pdf>).

#### References

- Murray CJL, Lopez AD, eds. The global burden of disease: a comprehensive assessment of mortality and disability from diseases, injuries, and risk factors in 1990 and projected to 2020. Boston: Harvard School of Public Health, 1996.
- Yusuf S, Reddy S, Ounpuu S, Anand S. Global burden of cardiovascular diseases, part I: general considerations, the epidemiologic transition, risk factors, and impact of urbanization. *Circulation* 2001; **104**: 2746–53.
- Pais P, Pogue J, Gerstein H, et al. Risk factors for acute myocardial infarction in Indians: a case-control study. *Lancet* 1996; **348**: 358–63.
- Yusuf S, Reddy S, Ounpuu S, Anand S. Global burden of cardiovascular diseases, part II: variations in cardiovascular disease by specific ethnic groups and geographic regions and prevention strategies. *Circulation* 2001; **104**: 2855–64.
- Ounpuu S, Negassa A, Yusuf S, for the INTER-HEART investigators. INTER-HEART: a global study of risk factors for acute myocardial infarction. *Am Heart J* 2001; **141**: 711–21.
- Wilhelmsen L, Rosengren A, Johansson S, Lappas G. Coronary heart disease attack rate, incidence and mortality 1975–1994 in Göteborg, Sweden. *Eur Heart J* 1997; **18**: 572–81.
- Rosengren A, Hawken S, Ounpuu S, et al. Association of psychosocial risk factors with risk of acute myocardial infarction in 11 119 cases and 13 648 controls from 52 countries (the INTERHEART study): case-control study *Lancet* 2004; **364**: 953–62.
- Marcovina SM, Albers JJ, Kennedy H, et al. International Federation of Clinical Chemistry Standardization Project for Measurements of Apolipoproteins A-1 and B: IV comparability of apolipoprotein B values by use of International Reference Material. *Clin Chem* 1994; **40**: 586–92.
- Marcovina SM, Albers JJ, Henderson LO, Hannon WH. International Federation of Clinical Chemistry Standardization Project for Measurements of Apolipoprotein A-1 and B: III comparability of apolipoprotein A-1 values by use of International Reference Material. *Clin Chem* 1993; **39**: 773–81.
- Walldius G, Jungner I, Holme I, Aastveit AH, Kolar W, Steiner E. High apolipoprotein B, low apolipoprotein A-1, and improvement in the prediction of fatal myocardial infarction (AMORIS study): a prospective study. *Lancet* 2001; **358**: 2026–33.
- Breslow N, Day N. Statistical methods in cancer research, vol 1: the analysis of case-control studies. Lyon: IARC Scientific Publications, 1980.
- Walter SD. The distribution of Levin's measure of attributable risk. *Biometrika* 1975; **62**: 371–74.
- Engel LS, Chow WH, Vaughan TL, et al. Population attributable risks of esophageal and gastric cancers. *J Natl Cancer Inst* 2003; **95**: 1404–13.
- Bruzzi P, Green SB, Byar DP, et al. Estimating the population attributable risk for multiple risk factors using case-control data. *Am J Epidemiol* 1985; **122**: 904–14.
- Benichou J, Gail MH. Variance calculations and confidence intervals for estimates of the attributable risk based on logistic models. *Biometrics* 1990; **46**: 991–1003.
- Cohen J. A coefficient of agreement for nominal scales. *Educ Psychol Meas* 1960; **20**: 37–46.
- MacMahon S, Peto R, Cutler J, et al. Blood pressure, stroke, and coronary heart disease: part 1, prolonged differences in blood pressure: prospective observational studies corrected for the regression dilution bias. *Lancet* 1990; **335**: 765–74.
- Barrett-Connor E, Cohn BA, Wingard D, Edelstein SL. Why is diabetes mellitus a stronger risk factor for fatal ischemic heart disease in women than in men? The Rancho Bernardo Study. *JAMA* 1991; **265**: 627–31.
- Canto JG, Iskandrian AE. Major risk factors for cardiovascular disease: debunking the “only 50%” myth. *JAMA* 2003; **290**: 947–49.
- Stamler J, Stamler R, Neaton JD, et al. Low risk-factor profile and long-term cardiovascular and noncardiovascular mortality and life expectancy: findings for 5 large cohorts of young adult and middle-aged men and women. *JAMA* 1999; **282**: 2012–18.
- Rosengren A, Dotevall A, Eriksson H, Wilhelmsen L. Optimal risk factors in the population: prognosis, prevalence, and secular trends. *Eur Heart J* 2001; **22**: 136–44.
- Blood Pressure Lowering Treatment Trialists' Collaboration. Effects of different blood-pressure-lowering regimens on major cardiovascular events: results of prospectively-designed overviews of randomised trials. *Lancet* 2003; **362**: 1527–35.
- Heart Protection Study Collaborative Group. MRC/BHF Heart Protection Study of cholesterol-lowering with simvastatin in 5963 people with diabetes: a randomised placebo-controlled trial. *Lancet* 2003; **361**: 2005–16.
- de Lorgeril M, Salen P, Martin JL, et al. Mediterranean diet, traditional risk factors, and the rate of cardiovascular complications after myocardial infarction: final report of the Lyon Diet Heart Study. *Circulation* 1999; **99**: 779–85.
- Parish S, Collins R, Peto R, et al. Cigarette smoking, tar yields, and non-fatal myocardial infarction: 14000 cases and 32000 controls in the United Kingdom. *BMJ* 1995; **311**: 471–77.
- Doll R, Peto R, Boreham J, Sutherland I. Mortality in relation to smoking: 50 years' observations on male British doctors. *BMJ* 2004; **328**: 1519–28.
- Wald NJ, Law MR. A strategy to reduce cardiovascular disease by more than 80%. *BMJ* 2003; **326**: 1419–23.
- Stampfer MJ, Hu FB, Manson JE, et al. Primary prevention of coronary heart disease in women through diet and lifestyle. *N Engl J Med* 2000; **343**: 16–22.

## Research Article

# Effectiveness of advance directives for the care of terminally ill patients in Chiang Mai University Hospital, Thailand

Sudarat Sittisombut, RN, PhD,<sup>1</sup> Colleen Maxwell, PhD,<sup>2</sup> Edgar J. Love, MD, PhD<sup>3</sup> and Chitr Sitthi-Amorn, MD, PhD<sup>4</sup>

<sup>1</sup>Faculty of Nursing, Chiang Mai University, Chiang Mai, <sup>4</sup>Department of Medicine, Faculty of Medicine, Chulalongkorn University, Bangkok, Thailand, <sup>2</sup>Department of Community Health Sciences and Medicine and <sup>3</sup>Department of Community Health Sciences, Faculty of Medicine, University of Calgary, Calgary, Alberta, Canada

## Abstract

The key hypothesis behind advance directives (ADs) proposes that, if an intervention enhances a person's right to choose, a dying person will not opt for expensive, life-prolonging medical care and an ethically acceptable saving of resources will result. In order to assess the acceptability and effectiveness of ADs in reducing cardiopulmonary resuscitation (CPR) attempts and in-hospital death among terminally ill patients in a tertiary care hospital in northern Thailand, a non-randomized, controlled intervention study using an after-only unequal control group design was conducted. The majority of the subjects and the surrogates preferred to employ ADs in expressing their preferences on CPR and there was a high level of agreement between the subjects and surrogates on the decision. The use of ADs appeared to be effective in reducing futile CPR attempts and the in-hospital mortality rate among subjects during the index hospitalization. Advance directives were accepted well in this study setting.

## Key words

advance directives, cardiopulmonary resuscitation, terminal care, terminally ill patients.

## INTRODUCTION

Cardiopulmonary resuscitation (CPR) is one of the most frequently performed medical interventions. This technique was originally developed to revive victims of sudden cardiac or respiratory arrest (Kowenhoven *et al.*, 1960). In many hospitals, it has been attempted on any patient who has cardiopulmonary arrest regardless of the underlying disease. Such routine application of CPR has contributed to several problems, including high mortality rates after CPR attempts (DeBard, 1981; Bedell *et al.*, 1983; McGrath, 1987; Rozenbaum & Shenkman, 1988; Peberdy *et al.*, 2003), continued physical suffering and worsening of clinical status (Bedell *et al.*, 1983), a vegetative state (Rozenbaum & Shenkman, 1988; FitzGerald *et al.*, 1997), high cost (Detsky *et al.*, 1981), and financial burden (FitzGerald *et al.*, 1997; Desbiens & Wu, 2000; Levenson *et al.*, 2000; Lynn *et al.*, 2000; Somogyi-Zalud *et al.*, 2000; Whitcomb & Blackman, 2007). The high death rate following CPR might reflect the fact that CPR is commonly performed on patients with terminal illnesses who

often experience a poor outcome following CPR (Bedell *et al.*, 1983; Rozenbaum & Shenkman, 1988).

Terminally ill patients usually suffer from the deterioration of their disease processes, such as severe pain (Lynn *et al.*, 2000; Roth *et al.*, 2000; Somogyi-Zalud *et al.*, 2000), dyspnea (Desbiens & Wu, 2000; Levenson *et al.*, 2000; Lynn *et al.*, 2000), depression (Block, 2000; Levenson *et al.*, 2000; Lynn *et al.*, 2000), and confusion (Lynn *et al.*, 2000; Somogyi-Zalud *et al.*, 2000). Following successful CPR, they might go on living, but their lives could be marred by severe and sometimes intolerable suffering. The inappropriate resuscitation of patients can result in an increased cost of care with little tangible medical benefits.

Our previous study of hospital deaths in a tertiary care hospital reveals that 65.7% of terminally ill patients received CPR prior to death (Sittisombut *et al.*, 2001). The decision to perform CPR in this setting was based mainly on the physicians in charge, with minimal input obtained from patients or their relatives. Several studies indicate that most terminally ill patients prefer CPR to be limited in the face of unfavorable outcomes (Steinbrook *et al.*, 1986; Covinsky *et al.*, 2000) and an increased familiarity with CPR usually leads to a decreased desire for this maneuver (Murphy *et al.*, 1994; Desbiens & Wu, 2000; Sittisombut *et al.*, 2005). The excessive use of CPR could be reduced if terminally ill patients

Correspondence address: Sudarat Sittisombut, Faculty of Nursing, Chiang Mai University, 110 Intawaroros Rd, Chiang Mai 50200, Thailand. Email: sudarat@mail.nurse.cmu.ac.th

Received 19 April 2007; accepted 26 July 2007

are informed well in advance and their preferences regarding the use of CPR are taken into consideration.

By providing a means for competent persons to guide their medical care after they have become incompetent, advance directives (ADs) are regarded as an effective mechanism to ensure that patients' wishes about medical treatment at the end of their life are respected (Crane *et al.*, 2005). However, although ADs are widely advocated in Western countries, only a few studies dealing with the use of ADs in Asian countries have been reported (Low *et al.*, 2000; Miura *et al.*, 2001; Akabayashi *et al.*, 2003) and ADs have not yet achieved legal recognition in this region.

Whether it is possible to engage terminally ill patients in discussing and making decisions on such profound issues as CPR and terminal care planning remains unknown in Thailand. This study was intended to determine whether ADs were acceptable to terminally ill inpatients in a tertiary care hospital in northern Thailand and to examine the effectiveness of ADs in reducing the use of CPR, as well as the associated in-hospital deaths among these patients.

## METHODS

This study employed a quasi-experimental design comparing the patients allotted to the intervention group with those in the non-randomized control group. It was conducted after the research proposal received approval from the institutional review board of the Faculty of Medicine, Chiang Mai University (CMU).

Patients who were admitted to the medical wards in the CMU Hospital with at least one of the following five diagnoses, as determined by the physicians in charge, were eligible for the study: non-small cell lung cancer, stages III or IV (NSCLC), multi-organ system failure with sepsis, untraumatic and non-diabetic coma with Glasgow coma score of 3, carcinoma of the colon with liver metastasis, and end-stage liver disease (ESLD). A surrogate, a person who was most qualified to make decisions on behalf of each patient, also was recruited. Inclusion was limited to patients and/or surrogates who were at least 40 years of age, alert, oriented, able to speak Thai, and who provided written informed consent for participation in the study.

During the period from 1 April to 30 November 2001, 217 patients who were eligible to participate in the study were assigned as potential control subjects. In a subsequent period, 1 December 2001 to 31 May 2002, an additional 231 eligible patients were recruited into the intervention group. Among all the eligible patients, only 376 met the inclusion criteria. The final study sample consisted of 188 subjects in each group. Comparisons revealed no significant differences in the number of exclusions from the control group (13.4%, [95% CI: 8.8–17.9]) and the intervention group (18.6%, [95% CI: 13.6–23.6]) with respect to discharge or transfer before the interview was possible, clinical deterioration or death.

During the first of the two chronologically separated phases, the enrollment of and observations on the control group were performed. To identify potentially eligible study subjects, the investigator, two nurses, and a research assistant made rounds and reviewed all hospital admissions in

each of the medical wards from Monday to Friday of the business week. For admissions during the weekend and holidays, the patient records were searched in the same manner on the following business day. For those considered eligible, the team introduced themselves to the patient and/or surrogate, explained the objective of the study, and invited them to participate in the study. The demographic and clinical data were gathered from the medical records and at the initial interview for the consenting subjects. The three outcomes (CPR/do not resuscitate [DNR] event, hospital discharge, and death in hospital) were observed and recorded on a daily basis.

In the second phase, the patients eligible for the study were identified and enrolled as subjects in the intervention group in the same manner as in the first phase. The only difference between the control group and the intervention group was the administration of the AD intervention prior to the observation in the latter. A tool for the AD intervention was developed, based on the relevant literature that reflected the objectives of the study, and it was validated by six experts. To adapt the tool to the culture and beliefs of the subjects, input from 10 terminally ill patients was sought by face-to-face discussion with the patients. Following the appropriate adaptation, the same six experts were invited to review, comment, and provide additional suggestions. Their comments and suggestions were used in the final adjustment of the tool. The inter-rater reliability of the tool after improvement was 8.5, as assessed by using Cohen's kappa method.

In the implementation of ADs, nurses interacted with patients and their surrogates in a step-by-step manner. They: (i) provided assurance to patients regarding their authority and autonomy; (ii) encouraged patients and their surrogates to seek information on diagnosis and prognosis from the attending physicians; (iii) provided information on CPR and its associated outcomes; (iv) assessed the patients' perception of authority and autonomy, diseases, prognosis, and CPR and its outcomes for their specific illnesses; (v) provided psychological support to patients and their families during the entire study period; and (vi) assessed the patients' preference for CPR. The surrogates also were asked whether they wished to have CPR performed on the respective patients and their choices on behalf of the patients were recorded. In order to obtain independent data from the patients and their surrogates, the interviews were performed separately for each patient and surrogate. In every step of the implementation of ADs, adequate time was allowed for discussion, clarification, and questions. Otherwise, an appointment for a return visit was made. The nature and timing of the visits depended on the wishes of the patients.

## Ethical considerations

The implementation of ADs was a sensitive issue as psychological trauma might occur in terminally patients in the intervention group. This problem was alleviated by providing a comprehensive approach and psychological support to the subjects and their families prior to enrollment. Participation was entirely voluntary and was sought only after patients and/or surrogates were given relevant information in at least

one session. Each potential subject and/or surrogate was approached and the research assistant explained the purpose of the study and obtained written consent. Only research assistants with adequate basic knowledge in research, counseling skills, and experience in caring for patients with terminal illness were selected for the study. They were trained in the implementation of ADs at the beginning of the study and then regularly supervised throughout the study. In addition, the study was approved by the institutional review board of the study hospital.

### Statistical analysis

The baseline demographic and clinical characteristics of the patient groups were determined using descriptive statistics. The differences between the control group and the intervention group were tested by using *t*-tests for the continuous variables and  $\chi^2$  tests for the categorical variables. The Mantel-Haenszel (MH)  $\chi^2$  or covariate analyses were performed to test the homogeneity for any significant difference observed.

## RESULTS

### Baseline demographic and clinical characteristics

The majority of the subjects were married males who lived in a rural area with limited formal education and irregular income, while NSCLC and ESLD represented the two most common diagnoses, accounting for 93.3% of these patients. Comparisons of the demographic characteristics and selected clinical parameters revealed that, with two exceptions, there were no significant differences between the intervention group and the control group (Tables 1,2). The clinical parameters that differentiated between the two groups were the proportions of patients with ESLD and high cerebral performance categories (CPC) scores (Table 2). The proportion of patients admitted with ESLD was greater in the control group than in the intervention group (35.1% vs 19.1%). Similarly, the proportion of the control subjects with CPC scores of 3 and 4, representing patients with dependency regarding their daily activities, was almost double that in the intervention group (57.4%, [95% CI: 50.4–64.5] vs 31.4% [95% CI: 24.7–38.0]). Apparently, the control subjects were in relatively poorer clinical conditions than the intervention subjects. However, when the MH  $\chi^2$  test was employed to assess the distribution of patients based on the proportion of patients with ESLD and high CPC scores, the result showed that there was homogeneity of the study subjects.

### Outcomes of the advance directives implementation

When the ADs were implemented for 188 subjects allocated to the intervention group, it was possible to identify only 132 (70.2%) surrogates willing to participate in the study. Among these subject-surrogate pairs, 80 (60.6%) chose to employ ADs, whereas 17 (12.9%) did not (Table 3). There was an agreement between the patients and their surrogates on the decision concerning the use of ADs in 73.5% of the pairs

**Table 1** Demographic characteristics

Parameter	Total ( <i>n</i> = 376) N (%)	Control group ( <i>n</i> = 188) N (%)	Intervention group ( <i>n</i> = 188) N (%)
Gender			
Male	243 (64.6)	123 (65.4)	120 (63.8)
Female	133 (35.4)	65 (34.6)	68 (36.2)
Age (years)			
Mean (SD)	57.5 (11)	57.0 (10)	58.0 (12)
Range	40–96	40–85	40–96
40–49	103 (27.4)	52 (27.7)	51 (27.1)
50–59	103 (27.4)	55 (29.3)	48 (25.5)
≥ 60	170 (45.2)	81 (43.0)	89 (47.4)
Marital status			
Married	312 (83.0)	156 (83.0)	156 (83.0)
Single	15 (4.0)	9 (4.8)	6 (3.2)
Widowed	49 (13.0)	23 (12.2)	26 (13.8)
Education			
≥ Grade 9	107 (28.5)	64 (34.0)	43 (22.9)
≤ Grades 4–6	254 (67.5)	124 (66.0)	130 (69.1)
No formal education	15 (4.0)	–	15 (8.0)
Income			
Regular income	79 (21.0)	46 (24.5)	33 (17.6)
Irregular income	179 (47.6)	84 (44.7)	95 (50.5)
No income	118 (31.4)	58 (30.9)	60 (31.9)
Residence			
Urban	99 (26.3)	53 (28.2)	46 (24.5)
Rural	277 (73.7)	135 (71.8)	142 (75.5)
Religion			
Buddhist	370 (98.4)	185 (98.4)	185 (98.4)
Others	6 (1.6)	3 (1.6)	3 (1.6)

(Table 3). Notably, the patients and their surrogates were generally hesitant to sign an AD document, preferring instead to have their directives communicated orally to one another.

During the index hospitalization, 33 subjects (8.8%) underwent CPR attempts, whereas 342 subjects (90.9%) were hospitalized without CPR. Only one subject gave a DNR order (Table 4). Of the 33 subjects who received CPR, 24 (12.8% [95% CI: 8.0–17.5]) were in the control group and nine (4.8% [95% CI: 1.7–7.9]) were in the intervention group (Table 4).

Of the 376 subjects, 315 (83.8%) left the study hospital alive and 61 (16.2%) died in the hospital (Table 5). Among the latter, 44 (23.4% [95% CI: 17.4–29.5]) were in the control group and 17 (9.0% [95% CI: 4.9–13.1]) were in the intervention group.

## DISCUSSION

This study was designed to assess the acceptability and effectiveness of ADs for terminally ill patients in a tertiary care, teaching hospital in northern Thailand. The patients in the control group and the intervention group were separately enrolled in two phases of different duration. The disparity reflects a major organizational change in the Thai public

**Table 2** Clinical characteristics

Parameter	Total ( <i>n</i> = 376) N (%)	Control group ( <i>n</i> = 188) N (%)	Intervention group ( <i>n</i> = 188) N (%)
Diagnosis			
NSCLC	249 (66.2)	118 (62.8)	131 (69.7)
ESLD	102 (27.1)	66 (35.1)	36 (19.1)
Others	25 (6.7)	4 (2.1)	21 (11.2)
MOSFS	7 (1.9)	–	7 (3.7)
Coma	10 (2.7)	1 (0.5)	9 (4.8)
Ca colon	8 (2.1)	3 (1.6)	5 (2.7)
Presence of comorbidity	121 (32.2)	63 (33.5)	58 (30.9)
CPC score: independent	209 (55.6)	80 (42.6)	129 (68.6)
CPC 1	113 (30.1)	41 (21.8)	72 (38.3)
CPC 2	96 (25.5)	39 (20.8)	57 (30.3)
CPC score: dependent	167 (44.4)	108 (57.4)	59 (31.4)
CPC 3	150 (39.9)	104 (55.3)	46 (24.5)
CPC 4	17 (4.5)	4 (2.1)	13 (6.9)
Mental status			
Oriented	342 (91.0)	171 (91.0)	171 (91.0)
Not oriented	34 (9.0)	17 (9.0)	17 (9.0)
Confused	17 (4.5)	13 (6.9)	4 (2.1)
Coma	17 (4.5)	4 (2.1)	13 (6.9)

CA colon, cancer of the colon with metastases to the liver; CPC, cerebral performance categories; ESLD, end-stage liver disease; MOSFS, multiple organ system failure with sepsis; NSCLC, stages III and IV non-small cell lung cancer.

**Table 3** Agreement on the employment of advance directives (ADs) between patients and surrogates

	Surrogate		Total N (%)
	Yes AD N (%)	No AD N (%)	
Patient			
Yes AD	80 (60.6)	13 (9.8)	93 (70.5)
No AD	22 (16.7)	17 (12.9)	39 (29.5)
Total	102 (77.3)	30 (22.7)	132 (100.0)

health system, which occurred during the first phase of the study. Patients had to pay all costs of medical care before and at the beginning of this study. However, starting in June 2001, the Ministry of Public Health stipulated that patients had to pay only 30 Bahts per hospital visit (Health System Reform Office, 2000). This benefit applied only when patients were treated in specifically designated, primary health care facilities. Based on the size of the registered population, the primary health care facility was allocated a fixed budget to cover expenses for the care of patients in their setting and for all costs after referral. This policy adversely affected recruitment in that, during the early period of this policy, the primary facility was reluctant to refer patients, especially those with terminal illnesses, as it could consume large amounts of the budget. During the first 3 months of the

**Table 4** Cardiopulmonary resuscitation (CPR) attempts

Event	Total ( <i>n</i> = 376) N (%)	Control group ( <i>n</i> = 188) N (%)	Intervention group ( <i>n</i> = 188) N (%)
CPR performed	33 (8.8)	24 (12.8)	9 (4.8)
CPR not performed	342 (90.9)	164 (87.2)	178 (94.7)
DNR order	1 (0.3)	–	1 (0.5)

DNR, do not resuscitate.

**Table 5** In-hospital mortality and outcome

Outcome	Total ( <i>n</i> = 376) N (%)	Control group ( <i>n</i> = 188) N (%)	Intervention group ( <i>n</i> = 188) N (%)
Dead	61 (16.2)	44 (23.4)	17 (9.0)
Alive	315 (83.8)	144 (76.6)	171 (91.0)
Discharge	256 (68.1)	109 (58.0)	147 (78.2)
Self-discharge	51 (13.5)	32 (17.0)	19 (10.1)
Transfer	8 (2.2)	3 (1.6)	5 (2.7)

implementation of the policy, the number of admissions of terminally ill patients in the study hospital, mostly from referral, decreased from about 25–30 subjects per month to = 20 per month. There was, however, a “rebound” in the subsequent period. The fluctuation in the admission of potential subjects necessitated the adjustment of the recruitment time, especially in the first phase, so that an adequate number of patients could be enrolled.

When the subjects in the intervention group in this study were provided with relevant information and the choice of using ADs to express their wishes, the majority of them decided in favor of using ADs to express their preference on CPR. This finding contrasts with the observation that higher education and socioeconomic status are associated with higher rates of AD completion in the USA (Braun *et al.*, 2001), as most of our subjects had relatively low socioeconomic status and minimal formal education. The high level of acceptability of ADs detected in this study might reflect the method of implementation of ADs and a high regard for the health profession, especially the personnel who provide services in the tertiary care facilities, by people of northern Thai culture, or both. In a previous study, 20 housebound elderly people with restricted formal education were resistant to planning in advance for the hypothetical future when they were interviewed in only one session that took up to 2 h to complete (Carrese *et al.*, 2002). An element in the successful implementation of ADs in our study might lie in the method of intervention, in which empathetic nurses conducted multiple interviews that were paced according to the perception and emotional status of the patients.

An important finding in this study is the observation that patients and their surrogates were only willing to express

their preference on CPR by engaging in verbal discussion, not by signing a document. Our finding is similar to that of Gamble *et al.* (1991), who found that no elder person had signed the living will document. Other studies reported only 4–47% of patients with a completed living will document (Sugarman *et al.*, 1992; Miles *et al.*, 1996; Kerridge *et al.*, 1998; Heiman *et al.*, 2004). It has been suggested that a possible explanation for the low rates of AD use by older people is that they trust their families to make decisions for them (Puchalski *et al.*, 2000). In the absence of a written document, an informal expression of a person's need might help reduce the difficulties when the family decides to make a substitute judgment. Proper management of this issue will be crucial in the large-scale implementation of ADs in the future.

The lower CPR and in-hospital mortality rates detected in the intervention group in this study might not entirely reflect the effectiveness of AD implementation. Other factors, including the discrepancy of the CPC score and the proportion of ESLD between the two groups, might have contributed to the differences. It is likely that the discrepancy between the physician's decision and the patient's preference could modify the outcomes that we measured, especially on the CPR attempts. Although most patients who died in a previous study did not want aggressive care, 63% still received life-sustaining treatment (Somogyi-Zalud *et al.*, 2002). However, it should be noted that, despite the major changes in the Thai public health system, the policy toward CPR in terminally ill medical patients remained unchanged in the study hospital throughout the two phases of the study. As a result of this consistency in the hospital policy, the possible influence of the variations in timing and interval of the two enrollment periods on the outcomes would be minimal.

## Limitations

The study was performed in two chronologically separated phases without randomization. Only terminally ill patients in a tertiary care, teaching hospital were studied. The preference for AD use in other groups might be different and it would be important to extend the study to unhospitalized subjects of other age groups. Also, it would be necessary to examine whether the implementation of ADs is possible and effective in other settings.

## ACKNOWLEDGMENTS

This study was supported by the Golden Jubilee Scholarship from Thailand Research Fund.

## REFERENCES

- Akabayashi A, Slingsby BT, Kai I. Perspectives on advance directives in Japanese society: a population-based questionnaire survey. *BMC Med. Ethics* 2003; **4**: E5.
- Bedell SE, Delbanco TL, Cook EF, Epstein FH. Survival after cardiopulmonary resuscitation in the hospital. *N. Engl. J. Med.* 1983; **309**: 569–576.
- Block SD. Assessing and managing depression in the terminally ill patient. *Ann. Intern. Med.* 2000; **132**: 209–218.
- Braun KL, Onaka AT, Horiuchi BY. Advance directive completion rates and end-of-life preferences in Hawaii. *J. Am. Geriatr. Soc.* 2001; **49**: 1708–1713.
- Carrese JA, Mullaney JL, Faden RR, Finucane TE. Planning for death but not serious future illness: qualitative study of house-bound elderly patients. *BMJ* 2002; **325**: 125–127.
- Covinsky KE, Fuller JD, Yaffe K *et al.* Communication and decision-making in seriously ill patients: findings of the SUPPORT project. The Study to Understand Prognoses and Preferences for Outcomes and Risks of Treatment. *J. Am. Geriatr. Soc.* 2000; **48**: S187–S193.
- Crane MK, Wittink M, Doukas DJ. Respecting end-of-life treatment preferences. *Am. Fam. Physician* 2005; **72**: 1263–1268, 1270.
- DeBard ML. Cardiopulmonary resuscitation – analysis of six years' experience and review of the literature. *Ann. Emerg. Med.* 1981; **10**: 408–416.
- Desbiens NA, Wu AW. Pain and suffering in seriously ill hospitalized patients. *J. Am. Geriatr. Soc.* 2000; **48**: S183–S186.
- Detsky AS, Stricker SC, Mulley AG, Thibault GE. Prognosis, survival, and the expenditure of hospital resources for patients in an intensive-care unit. *N. Engl. J. Med.* 1981; **305**: 667–672.
- FitzGerald JD, Wenger NS, Califf RM *et al.* Functional status among survivors of in-hospital cardiopulmonary resuscitation. SUPPORT investigators study to understand progress and preferences for outcomes and risks of treatment. *Arch. Intern. Med.* 1997; **157**: 72–76.
- Gamble ER, McDonald PJ, Lichstein PR. Knowledge, attitudes, and behavior of elderly persons regarding living wills. *Arch. Intern. Med.* 1991; **151**: 277–280.
- Heiman H, Bates DW, Fairchild D, Shaykevich S, Lehmann LS. Improving completion of advance directives in the primary care setting: a randomized controlled trial. *Am. J. Med.* 2004; **117**: 318–324.
- Kerridge IH, Pearson SA, Rolfe IE, Lowe M. Decision making in CPR: attitudes of hospital patients and healthcare professionals. *Med. J. Aust.* 1998; **169**: 128–131.
- Kowenhoven WB, Jude JR, Knickbocker GG. Closed-chest cardiac massage. *JAMA* 1960; **173**: 1064–1067.
- Levenson JW, McCarthy EP, Lynn J, Davis RB, Phillip RS. The last six months of life for patients with congestive heart failure. *J. Am. Geriatr. Soc.* 2000; **48**: S101–S109.
- Low JA, Ng WC, Yap KB, Chan KM. End-of-life issues: preferences and choices of a group of elderly Chinese subjects attending a day care center in Singapore. *Ann. Acad. Med. Singapore* 2000; **29**: 50–56.
- Lynn J, Ely EW, Zhong Z *et al.* Living and dying with chronic obstructive pulmonary disease. *J. Am. Geriatr. Soc.* 2000; **48**: S91–S100.
- McGrath RB. In-house cardiopulmonary resuscitation – after a quarter of a century. *Ann. Emerg. Med.* 1987; **16**: 1365–1368.
- Miles SH, Koepp R, Weber EP. Advanced end-of life treatment planning: a research review. *Arch. Intern. Med.* 1996; **156**: 1062–1068.
- Miura Y, Asai A, Nagata S *et al.* Dialysis patients' preferences regarding cardiopulmonary resuscitation and withdrawal of dialysis in Japan. *Am. J. Kidney Dis.* 2001; **37**: 1216–1222.
- Murphy DJ, Burrows D, Santilli S *et al.* The influence of the probability of survival on patients' preferences regarding cardiopulmonary resuscitation. *N. Engl. J. Med.* 1994; **330**: 545–549.
- Peberdy MA, Kaye W, Ornato JP *et al.* Cardiopulmonary resuscitation of adults in the hospital: a report of 14 720 cardiac arrests from the National Registry of Cardiopulmonary Resuscitation. *Resuscitation* 2003; **58**: 297–308.

- Puchalski CM, Zhong Z, Jacobs MM *et al.* Patients who want their family and physician to make resuscitation decisions for them: observations from SUPPORT and HELP. Study to Understand Prognoses and Preferences for Outcomes and Risks of Treatment. Hospitalized Elderly Longitudinal Project. *J. Am. Geriatr. Soc.* 2000; **48**: S84–S90.
- Roth K, Lynn J, Zhong Z, Borum M, Dawson NV. Dying with end stage liver disease with cirrhosis: insights from SUPPORT. Study to Understand Prognoses and Preferences for Outcomes and Risks of Treatment. *J. Am. Geriatr. Soc.* 2000; **48**: S122–S130.
- Rozenbaum FA, Shenkman L. Predicting outcome of in-hospital cardiopulmonary resuscitation. *Crit. Care Med.* 1988; **16**: 583–586.
- Sittisombut S, Love EJ, Sitthi-Amorn C. Cardiopulmonary resuscitation performed in patients with terminal illness in Chiang Mai University Hospital, Thailand. *Int. J. Epidemiol.* 2001; **30**: 896–898.
- Sittisombut S, Love EJ, Sitthi-Amorn C. Attitudes toward advance directives and the impact of prognostic information on the preference for cardiopulmonary resuscitation in medical inpatients in Chiang Mai University Hospital, Thailand. *Nurs. Health Sci.* 2005; **7**: 243–250.
- Somogyi-Zalud M, Zhong Z, Hamel MB, Lynn J. The use of life-sustaining treatments in hospitalized persons aged 80 and older. *J. Am. Geriatr. Soc.* 2002; **50**: 930–934.
- Somogyi-Zalud E, Zhong Z, Lynn J, Dawson NV, Hamel MB, Desbiens NA. Dying with acute respiratory failure or multiple organ system failure with sepsis. *J. Am. Geriatr. Soc.* 2000; **48**: S140–S145.
- Steinbrook R, Lo B, Moulton J, Saika G, Hollander H, Volberding PA. Preference of homosexual man with AIDS for life-sustaining treatment. *N. Engl. J. Med.* 1986; **314**: 457–460.
- Sugarman J, Weinberger M, Samsa G. Factors associated with veterans' decisions about living will. *Arch. Intern. Med.* 1992; **152**: 343–347.
- Whitcomb JJ, Blackman VS. Cardiopulmonary resuscitation: how far have we come? *Dimens. Crit. Care Nurs.* 2007; **26**: 1–6.
- Health System Reform Office. *The National Health System Reform B.E.2543*. [Cited 20 Nov 2007.] Available from URL: [http://www.hsro.or.th/en/documents/What\\_is\\_HealthSystem\\_Reform.pdf](http://www.hsro.or.th/en/documents/What_is_HealthSystem_Reform.pdf)

## Research Article

# Attitudes toward advance directives and the impact of prognostic information on the preference for cardiopulmonary resuscitation in medical inpatients in Chiang Mai University Hospital, Thailand

Sudarat Sittisombut, RN, PhD, Assistant Professor,<sup>1</sup> Edgar J. Love, MD, PhD, Professor<sup>2</sup> and Chitr Sitthi-amorn, MD, PhD, Professor<sup>3</sup>

<sup>1</sup>*Department of Surgical Nursing, Faculty of Nursing, Chiang Mai University, Chiang Mai, Thailand,*

<sup>2</sup>*Department of Community Health Sciences, University of Calgary, Calgary, Alberta, Canada,* <sup>3</sup>*College of Public Health, Chulalongkorn University, Bangkok, Thailand*

## Abstract

Our previous study revealed that cardiopulmonary resuscitation (CPR) was performed in 65.7% of 411 terminally ill patients who died in a tertiary-care university hospital in northern Thailand. Advance directives (ADs) are needed to ensure that life-sustaining therapies are used more appropriately. To investigate inpatients' attitudes regarding ADs for CPR and the impact of providing prognostic information on treatment preferences for CPR, we interviewed a randomly selected group of 200 ambulatory medical inpatients in multiple sessions. The results showed that most subjects had a positive attitude towards ADs for CPR. The majority preferred to have CPR when no information was provided on the chance of survival. However, this proportion decreased depending on the prognostic scenarios. Our investigation suggested that the preference of patients for CPR should be assessed individually and gradually, with adequate information given on the chance of survival.

## Key words

advance directives, ambulatory patients, cardiopulmonary resuscitation.

## INTRODUCTION

Cardiopulmonary resuscitation (CPR) is common in hospitals in Thailand, but there have been only a few reports on the use of CPR in English-language journals (Suraseranivongse *et al.*, 1998; Sittisombut *et al.*, 2001) and none in Thai journals. Our previous study revealed that, of 411 terminally ill patients who died in Chiang Mai University Hospital, 65.7% received CPR prior to death, and that most patients receiving CPR had a cerebral performance category score of 3 or 4, indicating that they were in a very poor condition (Sittisombut *et al.*, 2001). In order to reduce the inappropriate use of CPR, counter-balancing measures are needed to ensure that life-sustaining therapies are used more

appropriately. Advance directives (ADs) are widely regarded as the best available mechanism to ensure that patients' wishes about medical treatment at the end of life are respected (Walker *et al.*, 1995; Roter *et al.*, 2000). The ADs provide a means for competent persons to guide their medical care after they have become incompetent (Emanuel & Emanuel, 1989; Daly & Sobal, 1992). Although ADs has been widely advocated in western countries (Danis *et al.*, 1991; Emanuel *et al.*, 1991; Molloy & Guyatt, 1991; Schneiderman *et al.*, 1992b), only two studies dealing with ADs were reported from Asian countries, including Singapore and Japan (Low *et al.*, 2000; Miura *et al.*, 2001), and none from Thailand, as ADs have not yet achieved legal recognition in these Asian countries.

Previous studies indicated that most patients with ADs prefer to limit care (Schneiderman *et al.*, 1992a; Weeks *et al.*, 1994; Covinsky *et al.*, 2000). However, High (1987) noted that discussions of the end-of-life care frequently occurred too late, after the person has

Correspondence address: Sudarat Sittisombut, Department of Surgical Nursing, Faculty of Nursing, Chiang Mai University, 110 Intavara Road, Chiang Mai 50200, Thailand. Email: sudarat@mail.nurse.cmu.ac.th  
Received 11 November 2004; accepted 9 February 2005.

been incapacitated. Therefore, it might be useful if patients could be counseled to plan and express their preferences in advance during a regular admission. By so doing, patient autonomy could be better assured and unnecessary care might be limited. However, such a study has not been tried in Thailand and, specifically, in the northern Thai culture. Whether it would be socially and culturally acceptable to ask patients to discuss so profound an issue as CPR and terminal care-planning remains unknown.

A wide variation in survival following in-hospital CPR episodes has been observed (Cummins *et al.*, 1997). Despite advances in CPR technology, the percentage of patients who are successfully resuscitated in the hospital setting has remained low (7–24%) during the past 30 years in Western countries (DeBard, 1981; Bedell *et al.*, 1983; McGrath, 1987; Woog & Torzillo, 1987; Rozenbaum & Shenkman, 1988; George *et al.*, 1989; Roberts *et al.*, 1990; Berger & Kelley, 1994). For some patient groups, survival can be < 5% or close to zero, depending on the type and severity of their diseases (Bedell *et al.*, 1983; Rozenbaum & Shenkman, 1988; Taffet *et al.*, 1988; George *et al.*, 1989; Moss, 1989; Peterson *et al.*, 1991). Pre-arrest clinical conditions, advanced age (McGrath, 1987; Taffet *et al.*, 1988; George *et al.*, 1989), and CPR that was performed in general hospital wards (Hershey & Fisher, 1982; George *et al.*, 1989) are associated with a poor prognosis. Moreover, the time delay before resuscitation, the quality of CPR efforts, and the duration of the CPR episode all contribute to the initial survival and the discharge survival (Bedell *et al.*, 1983; Woog & Torzillo, 1987; Rozenbaum & Shenkman, 1988).

Among several outcomes of CPR attempts, failure resulting in death is, by necessity, an acceptable risk. Other outcomes include a permanent vegetative state (Rozenbaum & Shenkman, 1988; FitzGerald *et al.*, 1997), and the worsening of clinical status (Detsky *et al.*, 1981; Bedell *et al.*, 1983; FitzGerald *et al.*, 1997). In a study by FitzGerald *et al.* (1997), it was found that almost half (44% of 162 cases) of patients who survived CPR were actually living with a worse level of functioning and were profoundly disabled. Many of the resuscitation efforts are followed by the use of mechanical ventilation and/or admission into an intensive care facility, both of which are quite costly (Detsky *et al.*, 1981).

Our goal was to assess the attitudes of Thai ambulatory patients in general medical wards toward ADs for CPR. We also studied the preferences of this population regarding CPR after providing information on different prognostic scenarios. The results of this study will guide our next step in ADs intervention in terminally ill patients.

## METHODS

After the proposal was approved by the Institutional Review Board of the Faculty of Medicine, Chiang Mai University, an acceptability study of ADs for CPR was conducted. This study was a part of the research project on the use of ADs for the care of terminally ill patients in Chiang Mai University Hospital in northern Thailand. The hospital is a government teaching facility, serving as the tertiary-care hospital for the northern provinces of Thailand. Routine admission is allowed from Monday to Friday (08.00 hours–04.00 hours); only emergency and/or seriously ill patients are admitted during unofficial times (16.00 hours–08.00 hours) and on weekends.

Patients who were admitted to all adult medical wards (three female and four male wards) from 1 November 2000 – 31 December 2000 and who met the inclusion criteria were randomly selected (generally 4–6 cases/day, depending on the number of admissions). Those who were  $\geq 18$  years of age, alert, orientated, able to communicate in Thai, and who agreed to participate by giving informed consent were eligible for the study. When a selected patient's consent could not be obtained, another eligible patient in the same ward was randomly selected. Patients with one or more of the following diagnoses were excluded: non-small cell lung cancer (stages III or IV), non-traumatic and non-diabetic coma, multi-organ system failure with sepsis, carcinoma of the colon with metastasis to the liver, end-stage liver disease, exacerbation of congestive heart failure, exacerbation of chronic obstructive pulmonary disease, and acute respiratory failure. This was because they might need a more specific approach. Similarly, patients with HIV infection or AIDS were excluded. Two-hundred patients were invited to participate in the study. Consent could not be obtained from three patients and replacements were selected accordingly. All subjects were interviewed in person by trained interviewers using a structured questionnaire.

The questionnaire was developed based on related literature that reflected the objective of the study. The questionnaire was then validated by five experts and pretested with 10 patients who had similar characteristics to our study subjects. The questionnaire inquired about the participants' gender, age, marital status, religion, education, personal income, and usual place of residence (rural/urban). After explaining the CPR procedure, the patients were asked about their preferences with respect to CPR in different prognostic scenarios associated with varied post-CPR survival. The scenarios for assessing CPR preferences were as follows: (i) without any information regarding the chance of survival; (ii) with the chance of discharge of 7–24% (as in

acute-onset diseases); (iii) with the chance of survival of 0–5% (as in some specific diseases); and (iv) CPR followed by living permanently on mechanical ventilation and/or coma. The responses to these questions were either “yes CPR”, “no CPR”, “up to physician”, or “up to relative”. Finally, patients were asked to express their feelings regarding the discussion of ADs for CPR.

During the reliability testing, it was found that the questionnaire should not be administered in a single interview as the patients appeared to be tense. The interviews were subsequently performed in multiple sessions. The inter-rater and intra-rater reliability, assessed using Cohen’s Kappa statistic (Munro & Page, 1993), were 0.80 and 0.85, respectively. No adjustment was made in the final questions or in their sequences, other than those previously suggested by the experts.

During the data collection phase, the interviews were conducted in person by the primary investigator and three nurses who had previous research experience, were trained in the use of the structured questionnaire, and who were supervised by the primary investigator. The questionnaire was administered over at least two sessions by empathetic nurse-researchers who “paced” the interview according to when the subject seemed ready for the next group of questions.

In the first interview, the interviewers introduced themselves, explained the objective, the process of the study, and the method of selection of the subjects. After receiving signed consent, the balance of the initial interview focused on general information, including the demographic data. The next interview started again with the general topic and, then, moved forward to the participants’ CPR preferences and their attitude toward advanced planning for CPR. All patients were assured that the confidentiality of their information would be respected.

In the analysis, the demographic characteristics of the patients and the responses of the interviewees were computed using descriptive statistics. Chi-squared statistics were used to test the difference (Polit & Hungler, 1999) in the demographic characteristics by gender. Patients’ preference for CPR (“no CPR” vs “yes CPR” in different survival scenarios) by gender, age, marital status, income level, residential place, and type of illness were assessed using the multiple logistic regression method (Munro, 2001). The odds ratio (OR) for each variable was adjusted for possible confounding by all other variables in the table. The statistically significant level was set at  $P < 0.05$ .

## RESULTS

Of the first 200 patients who fulfilled the inclusion criteria, 197 (98.5%) agreed to participate and only three

were additionally recruited. Among the participants, 129 (64.5%) were male (which was comparable to the gender distribution in the medical inpatients of Chiang Mai University Hospital), 52.0% were  $\geq 45$  years, 71.0% were married, 77.5% were residents of rural areas, and 62.0% were admitted with chronic illnesses. The majority (80.0%) was poor (without regular income, with unstable income, or with yearly personal income of 33 600 Baht, equivalent to  $\leq$  \$US840), with more than half (57.5%) having no regular income (Table 1). All were Buddhists. Seventy-five percent had primary education (grades 4–6). When the demographic data were compared between male and female participants, the only statistically significant difference was that more females were in the lower income category than males (93.0% vs 72.9%).

## Attitudes toward advance directives

Nearly all of the subjects (97.0%) thought that it was a good idea to discuss advanced planning for CPR on a routine basis for all admitted patients. All were interested in knowing the outcomes. However, our subjects expressed a distrust of formal documents and they preferred to provide oral instructions to surrogate(s) for their CPR planning.

## Preferences for cardiopulmonary resuscitation

After receiving the information only on the CPR procedures, most of the subjects (87.5%) preferred to have CPR. When the chances of survival were also provided, the proportion of patients with preference for CPR decreased to 68.5% when the chance of discharge was 7–24%, and to 45.5% when the chance of survival was 0–5%. Only 27.5% of the subjects still preferred to have CPR if it was followed by living permanently on mechanical ventilation and/or in a coma.

## Factors affecting preferences

Subjects’ preferences for or against CPR were affected by the level of prognostic information (Table 2). Gender, age, marital status, personal income, and the type of illness also affected the preferences.

Females were more likely to prefer no CPR, as compared with males, when the prognostic information was not provided (OR = 5.37, 95% confidence interval (CI) = 1.47–19.58), when the survival chance was 0–5% with CPR (OR = 3.10, 95% CI = 1.47–6.54), and when CPR might be followed by mechanical ventilation and/or coma (OR = 7.58, 95% CI = 2.91–19.76). Gender did not affect the preference only when there was a high chance of survival.

**Table 1.** Demographic characteristics

Characteristic	Male ( <i>n</i> = 129) N (%)	Female ( <i>n</i> = 71) N (%)	Total ( <i>n</i> = 200) N (%)	<i>P</i> *
Age (years)				0.246
< 45	58 (45.0)	38 (53.5)	96 (48.0)	
< 30	17 (13.2)	13 (18.3)	30 (15.0)	
30–44	41 (31.8)	25 (35.2)	66 (33.0)	
> 45	71 (55.0)	33 (46.5)	104 (52.0)	
≥ 60	30 (23.2)	15 (21.1)	45 (22.5)	
Marital status				0.646
Married	93 (72.1)	49 (69.0)	142 (71.0)	
Not married	36 (27.9)	22 (31.0)	58 (29.0)	
Single	24 (18.6)	13 (18.3)	37 (18.5)	
Widowed	8 (6.2)	8 (11.3)	16 (8.0)	
Divorced	4 (3.1)	1 (1.4)	5 (2.5)	
Personal income (Baht/year)				0.001**
≤ 33 600	94 (72.9)	66 (93.0)	160 (80.0)	
No regular income	62 (48.1)	53 (74.7)	115 (57.5)	
Unstable income	31 (24.0)	11 (15.5)	42 (21.0)	
≤ 33 600	1 (0.8)	2 (2.8)	3 (1.5)	
> 33 600	35 (27.1)	5 (7.0)	40 (20.0)	
33 601–60 000	17 (13.2)	2 (2.8)	19 (9.5)	
> 60 000	18 (13.9)	3 (4.2)	21 (10.5)	
Place of residence				0.730
Rural	99 (76.7)	56 (78.9)	155 (77.5)	
Urban	30 (23.3)	15 (21.1)	45 (22.5)	
Type of illness				0.756
Acute	48 (37.2)	28 (39.4)	76 (38.0)	
Chronic	81 (62.8)	43 (60.6)	124 (62.0)	

\**P*-value by chi-squared test with 1 degree of freedom; \*\**P* < 0.01.

Similar results also were observed in different age groups. Subjects who were ≥ 45 years (older adults) who were presented with the scenario in which the survival chance with CPR was 7–24% preferred no CPR four times more than the subjects who were < 45 years (younger adults) (OR = 3.96, 95% CI = 1.28–12.26). Meanwhile, when the scenario indicated that the survival chance with CPR was 0–5%, or when the scenario was that CPR might be followed by mechanical ventilation, coma, or both, older adults said they would prefer no CPR approximately two to three times more than younger adults (OR = 2.85, 95% CI = 1.33–6.12 and OR = 2.37, 95% CI = 1.02–5.50, respectively).

Subjects who were not married were more likely to decide in favor of no CPR compared to those who were married. However, the OR of preferences varied between 3.87, 3.40, and 2.34, depending on which of the following scenarios were presented: no prognostic information, a survival chance of 7–24% with CPR, and a survival chance of 0–5% with CPR, respectively.

Subjects who had a low income (≤ 33 600 Baht/year) were more likely to decide that no CPR would be their preference as compared to those who had a higher

income. Increased ORs for the preference of no CPR associated with low income had been observed in two scenarios: when the survival chance was 0–5% with CPR (OR = 3.26, 95% CI = 1.01–10.56), and if CPR might be followed by mechanical ventilation, coma, or both (OR = 7.88, 95% CI = 2.65–23.47).

Finally, the OR of preferring no CPR differed between patients admitted for chronic illness and patients admitted for acute illness. A significant OR was observed only in the scenario in which CPR might be followed by mechanical ventilation, coma, or both. Subjects with a chronic illness were more likely to express a preference for no CPR as compared to those with acute illness (OR = 3.12, 95% CI = 1.40–6.98).

### Deferral of decision

When the information regarding the prognosis was not provided, up to 7.7% of the male subjects decided to defer the decision, compared to none of the female subjects (Table 3). With the three prognostic circumstances, the proportion of the male subjects who deferred the decision increased to 22.5–34.1%,

**Table 2.** Preference for no cardiopulmonary resuscitation (CPR) by demographic variables when scenarios with different prognostic probabilities were provided

Variable	Scenario						CPR followed by mechanical ventilation and/or coma	
	No information		Survival of 7–24%		Survival of 0–5%		OR (95% CI)	P
	OR (95% CI)	P	OR (95% CI)	P	OR (95% CI)	P		
Gender	5.37 (1.47–19.58)	0.011	2.11 (0.70–6.33)	0.183	3.10 (1.47–6.54)	0.003	7.58 (2.91–19.76)	< 0.001
Age	1.85 (0.57–6.05)	0.306	3.96 (1.28–12.26)	0.017	2.85 (1.33–6.12)	0.007	2.37 (1.02–5.50)	0.045
Marital status (not married/married)	3.87 (1.21–12.37)	0.022	3.40 (1.13–10.22)	0.029	2.34 (1.03–5.28)	0.042	0.75 (0.31–1.77)	0.508
Personal income	0.37 (0.08–1.80)	0.217	0.90 (0.17–4.80)	0.904	3.26 (1.01–10.56)	0.049	7.88 (2.65–23.47)	< 0.001
Place of residence (urban/rural)	1.01 (0.25–4.05)	0.988	1.05 (0.31–3.64)	0.933	1.92 (0.83–4.46)	0.129	1.21 (0.46–3.21)	0.701
Type of illness (chronic/acute)	3.44 (0.87–13.52)	0.077	1.29 (0.43–3.86)	0.652	1.49 (0.70–3.19)	0.304	3.12 (1.40–6.98)	0.006

whereas this proportion remained low in the female subjects. All male subjects chose to defer their decision to the physicians but the female subjects preferred the relatives.

## DISCUSSION

This study was designed to survey ambulatory individuals who were admitted to general medical wards of a tertiary-care hospital in northern Thailand to assess their attitude toward ADS, their preferences for or against CPR, and to identify any issue(s) which needed improvement prior to the intervention study. Our results might be different from previous studies performed in more developed countries (Emanuel & Emanuel, 1989; Danis *et al.*, 1991; Emanuel *et al.*, 1991; Molloy & Guyatt, 1991; Daly & Sobal, 1992; Walker *et al.*, 1995). One of the distinguishing features of this study is the low socioeconomic status of the subjects. These patients were generally very poor; the proportion of subjects who were living in poverty was higher than the general Thai population (Tangcharoensathien *et al.*, 2000). The majority of subjects lived in rural areas and were undereducated. (Table 1)

## High response rate

A very high response rate (98.5%) was obtained from patients who were approached to enrol in this study and it was quite unexpected. Previous studies reported lower rates (77%) in other group of subjects (Murphy *et al.*, 1994). Such a high response might reflect the cultural norms of rural residents of northern Thailand and the way they think about health-care workers. However, we feel that our interviewing method, in which the interviewers employed an empathetic manner and took time to ensure that the patients were comfortable with the pace of the interviews, might also have increased the confidence of the interviewees in agreeing to participate and expressing their attitude and preferences. This approach should be useful in studying similar groups of subjects in the future.

The majority of our northern Thai subjects welcomed the discussion on CPR issues. Most agreed that planning for CPR should be discussed in advance. This is in contrast to a previous study in Singapore where only 37.2% of 43 elderly Chinese agreed that making an AD would be necessary (Low *et al.*, 2000). Although our subjects expressed distrust in signing formal documents and preferred to provide only oral instructions to their surrogates, this attitude is not uncommon. In other studies in the USA, Emanuel *et al.* (1993) and Campbell and Frank (1997; p. 200) found that the pro-

**Table 3.** Deferral of decision regarding cardiopulmonary resuscitation (CPR) by gender

Information provided before expression of preference	Decision deferred to physicians		Decision deferred to relatives	
	Male N (%)	Female N (%)	Male N (%)	Female N (%)
None	10 (7.7)	0	0	0
Survival of 7–24%	43 (33.3)	0	0	2 (2.8)
Survival of 0–5%	44 (34.1)	0	0	2 (2.8)
CPR might be followed by the need for permanent mechanical ventilation and/or coma	29 (22.5)	0	0	3 (4.2)

Total males = 129; total females = 71.

portion of patients who had advanced care planning “in a written document” was low. Similarly, Hanson *et al.* (1994) found that only 41% of the decisions to withhold CPR from 311 incompetent patients was guided by the patients’ preference. Although this practice is more acceptable to the patients, the lack of written ADs might pose practical problems for health-care workers when the actual decision for CPR has to be made in the absence of surrogates.

The proper use of ADs requires that patients have access to the necessary information and are allowed an adequate period of time to consider the alternatives. Our patients agreed that the assessment of preference concerning CPR should take place on a routine visit for every admitted patient. Another study has revealed that many ambulatory medical inpatients want to be involved in the decision-making on CPR (Gamble *et al.*, 1991). To this end, clinicians and/or other health-care workers should be encouraged to take a more active role in promoting discussions of the patient’s wishes regarding end-of-life care by informing patients and their families of the patient’s autonomy regarding advance medical planning. As we have found in this study, medical inpatients in our hospital settings are quite receptive to such discussion well before they become unable to participate in the decision process. The high rate of CPR performed on terminally ill patients reported earlier (Sittisombut *et al.*, 2001) indicates that the discussion on end-of-life care and/or ADs did not take place or received low priority. Further studies are needed to determine who among various health-care workers would be best responsible for this valuable work and how to motivate such personnel to initiate discussion with patients and their families.

In the absence of information regarding CPR outcomes, the majority of our subjects favored resuscitation; however, many, especially those who were poor, refused resuscitation when it would be accompanied by mechanical ventilation and/or coma. This change in preference reinforces the role of information in pref-

erence previously documented in many studies (or confirms that prognostic information had resulted in more patients’ refusal of CPR). Walker *et al.* (1995) reported that only a small percentage of retired elderly desired CPR in the scenarios of terminal illness, functional impairment, and cognitive impairment. Other surveys showed that most persons opposed aggressive treatment in the persistent vegetative state (Emanuel *et al.*, 1991), dementia, and terminal situations (Emanuel *et al.*, 1991; Miura *et al.*, 2001). Molloy and Guyatt (1991) found that 88% of 119 elderly persons in Canada requested not to even have CPR performed. The high proportion of patients with preference for CPR in the absence of prognostic scenarios in this study strongly suggests that patients have had limited access to information on the end-of-life care and the possible choices that they can make.

It is intriguing that gender strongly affects the decision and the expressed preference in our subjects. Male subjects chose to defer the decision more than their female counterparts. Moreover, all of the male subjects who decided to defer the decision after receiving each of the four prognostic scenarios preferred that their physician(s) should make the decision. In contrast, all of the females who choose to defer the CPR decision selected their family members as the decision-maker; none chose their physician(s). These results are quite contradictory to the role of the male. Generally, the man is the leader of the family in Asian societies (H. Sethabouppha, unpubl. data, 2002). For a serious medical topic, the man probably believes that if he could not make a decision, no one else in his family could do so. This may explain, in the case of non-deciding males, why the decision on CPR was delegated to the physician. The role of gender in preference was also evident in the finding that female subjects preferred the “no-CPR” decision more commonly than the male counterparts. This might reflect the role of females as the care provider in the family and the thinking that they should not expect help from other family members. It is notable that the OR

between the two genders was highest in the scenario where CPR would be accompanied by mechanical ventilation and/or coma.

There were some limitations in this study. This report concerns only ambulatory subjects at a tertiary-care regional hospital. As the study was conducted at a single site and targeted only northern Thai people with their unique culture, the study was clearly limited in terms of generalizability. The mental status of the subjects was not formally assessed; however, all patients were able to participate in the interview without difficulty.

## CONCLUSIONS

Our findings are different from previous studies in many aspects. We obtained a very high response rate and gender-specific differences that might indicate cultural differences. Most of our subjects were poor, had only primary education and were from rural areas. They were unfamiliar with the questions and specific information on the end-of-life care; therefore, the method of interview had been adjusted accordingly. This technique should be repeated to test its reliability in a similar group of subjects, as well as others. Importantly, our patients distrusted AD documents. These feelings were respected but further investigation is needed to identify the most effective strategy to address the patient's needs in this society.

We confirmed previous findings that the survival chance after CPR and the worst condition postCPR can dramatically change the preference for CPR. However, the preferences of other groups of subjects or patients with various other conditions might be quite different. For future study, it would be of interest to extend the study to non-hospitalized subjects (young adults < 40 years, middle-aged adults aged 40–59 years, and older adults ≥ 60 years), and hospitalized patients with a chronic and/or terminal illness. Also, it would be important to test whether the ADs are effective in reducing unnecessary CPR episodes in terminally ill patients in our health-care settings.

## ACKNOWLEDGMENT

This study was supported by the Golden Jubilee scholarship from Thailand Research Fund.

## REFERENCES

- Bedell SE, Delbanco TL, Cook EF *et al.* Survival after cardiopulmonary resuscitation in the hospital. *N. Engl. J. Med.* 1983; **309**: 569–576.
- Berger R, Kelley M. Survival after in-hospital cardiopulmonary arrest of non-critically ill patients. *Chest* 1994; **106**: 872–879.
- Campbell ML, Frank RB. Experience with end-of-life practice at a university hospital. *Crit. Care Med.* 1997; **25**: 197–201.
- Covinsky KE, Fuller JD, Yaffe K *et al.* Communication and decision-making in seriously ill patients: findings of the SUPPORT project, The Study to Understand Prognoses and Preferences for Outcomes and Risks of Treatments. *J. Am. Geriatr. Soc.* 2000; **48**: S187–S193.
- Cummins RO, Chamberlain D, Hazinski MF *et al.* Recommended guidelines for reviewing, reporting, and conducting research on in-hospital resuscitation: the in-hospital 'Utstein style'. *Circulation* 1997; **95**: 2213–2239.
- Daly MP, Sobal J. Advance directives among patients in a house call program. *J. Am. Board Fam. Pract.* 1992; **5**: 11–15.
- Danis M, Southerland LL, Garrett JM *et al.* A prospective study of advance directives for life-sustaining care. *N. Engl. J. Med.* 1991; **324**: 882–888.
- DeBard ML. Cardiopulmonary resuscitation—analysis of six years' experience and review of the literature. *Ann. Emerg. Med.* 1981; **147**: 37–38.
- Detsky AS, Stricker SC, Mulley AG *et al.* Prognosis, survival, and the expenditure of hospital resources for patients in an intensive-care unit. *N. Engl. J. Med.* 1981; **305**: 667–672.
- Emanuel EJ, Weinberg DS, Gonin R *et al.* How well is the patient self determination act working? An early assessment. *Am. J. Med.* 1993; **95**: 619–628.
- Emanuel LL, Barry MJ, Stoeckle JD *et al.* Advance directives for medical care: a case for greater use. *N. Engl. J. Med.* 1991; **324**: 889–895.
- Emanuel LL, Emanuel EJ. The medical directive: a new comprehensive advance care document. *JAMA* 1989; **261**: 3288–3293.
- FitzGerald JD, Wenger NS, Califf RM *et al.* Functional status among survivors of in-hospital cardiopulmonary resuscitation. SUPPORT Investigators Study to Understand Progress and Preferences for Outcomes and Risks of Treatment. *Arch. Intern. Med.* 1997; **157**: 72–76.
- Gamble ER, McDonald PJ, Lichstein PR. Knowledge, attitudes, and behavior of elderly persons regarding living wills. *Arch. Intern. Med.* 1991; **151**: 277–280.
- George AL Jr, Folk BP, Crecelius PL *et al.* Pre-arrest morbidity and other correlates of survival after in-hospital cardiopulmonary arrest. *Am. J. Med.* 1989; **87**: 28–34.
- Hanson LC, Danis M, Mutran E *et al.* Impact of patient incompetence on decisions to use or withhold life-sustaining treatment. *Am. J. Med.* 1994; **97**: 235–241.
- Hershey CO, Fisher L. Why outcome of cardiopulmonary resuscitation in general wards is poor. *The Lancet* 1982; **1**: 31–34.
- High DM. Planning for decisional incapacity: a neglected area in ethics and aging. *J. Am. Geriatr. Soc.* 1987; **35**: 814–820.
- Low JA, Ng WC, Yap KB *et al.* End-of-life issues: preferences and choices of a group of elderly Chinese subjects

- attending a day care center in Singapore. *Ann. Acad. Med. Sing.* 2000; **29**: 50–56.
- McGrath RB. In-house cardiopulmonary resuscitation—after a quarter of a century. *Ann. Emerg. Med.* 1987; **16**: 1365–1368.
- Miura Y, Asai A, Nagata S. *et al.* Dialysis patients' preferences regarding cardiopulmonary resuscitation and withdrawal of dialysis in Japan. *Am. J. Kidney Dis.* 2001; **37**: 1216–1222.
- Molloy DW, Guyatt GH. Comprehensive health care directives in a home for the aged. *Can. Med. Assoc.* 1991; **145**: 307–311.
- Moss AH. Informing the patient about cardiopulmonary resuscitation: when the risks outweigh the benefits. *J. Gen. Intern. Med.* 1989; **4**: 349–355.
- Munro BH. *Statistical Method in Health Care Research*, 4th edn. Philadelphia, PA: Lippincott, 2001.
- Munro BH, Page EB. *Statistical Method for Health Research*, 2nd edn. Philadelphia, PA: Lippincott, 1993.
- Murphy DJ, Burrows D, Santilli S *et al.* The influence of the probability of survival on patients' preferences regarding cardiopulmonary resuscitation. *N. Engl. J. Med.* 1994; **330**: 545–549.
- Peterson MW, Geist LJ, Schwartz DA *et al.* Outcome after cardiopulmonary resuscitation in a medical intensive care unit. *Chest* 1991; **100**: 168–174.
- Polit DF, Hungler BP. *Nursing Research: Principles and Methods*, 6th edn. Philadelphia, PA: Lippincott, 1999.
- Roberts D, Landofo K, Light RB *et al.* Early predictors of mortality for hospitalized patients suffering cardiopulmonary arrest. *Chest* 1990; **97**: 413–419.
- Roter DL, Larson S, Fischer GS *et al.* Experts practice what they preach: a descriptive study of best and normative practices in end-of-life discussions. *Arch. Intern. Med.* 2000; **160**: 3477–3485.
- Rozenbaum FA, Shenkman L. Predicting outcome of in-hospital cardiopulmonary resuscitation. *Crit. Care Med.* 1988; **16**: 583–586.
- Schneiderman LJ, Kronick R, Kaplan RM *et al.* Effects of offering advance directives on medical treatments and costs. *Ann. Intern. Med.* 1992a; **117**: 599–606.
- Schneiderman LJ, Pearlman RA, Kaplan RM *et al.* Relationship of general advance directive instructions to specific life-sustaining treatment preferences in patients with serious illness. *Arch. Intern. Med.* 1992b; **152**: 2114–2122.
- Sittisombut S, Love EJ, Sitthi-amorn C. Cardiopulmonary resuscitation performed in patients with terminal illness in Chiang Mai University Hospital, Thailand. *Int. J. Epidemiol.* 2001; **30**: 896–898.
- Suraseranivongse S, Somprakit P, Soontranant P *et al.* Factors influencing CPR outcome in Siriraj Hospital. *J. Med. Assoc. Thai.* 1998; **81**: 836–843.
- Taffet GE, Teasdale TA, Luchi RJ. In-hospital cardiopulmonary resuscitation. *JAMA* 1988; **260**: 2069–2072.
- Tangcharoensathien V, Harnvoravongchai P, Pitayarangsarit S *et al.* Health impacts of rapid economic changes in Thailand. *Soc. Sci. Med.* 2000; **51**: 789–807.
- Walker RM, Schonwetter RS, Kramer DR *et al.* Living wills and resuscitation preferences in an elderly population. *Arch. Intern. Med.* 1995; **155**: 171–175.
- Weeks WB, Kofoed LL, Wallace AE *et al.* Advance directives and the cost of terminal hospitalization. *Arch. Intern. Med.* 1994; **154**: 2077–2083.
- Woog RH, Torzillo PJ. In-hospital cardiopulmonary resuscitation: prospective survey of management and outcome. *Anaesth. Intens.* 1987; **15**: 193–198.

## Research Article

# Physicians' attitudes and practices regarding advanced end-of-life care planning for terminally ill patients at Chiang Mai University Hospital, Thailand

Sudarat Sittisombut, RN, PhD,<sup>1</sup> Colleen Maxwell, PhD,<sup>2,3</sup> Edgar J. Love, MD, PhD<sup>2</sup> and Chitr Sitthi-Amorn MD, PhD<sup>4</sup>

<sup>1</sup>Faculty of Nursing, Chiang Mai University, Chiang Mai, <sup>4</sup>Department of Medicine, Faculty of Medicine, Chulalongkorn University, Bangkok, Thailand; and Departments of <sup>2</sup>Community Health Sciences and <sup>3</sup>Medicine, Faculty of Medicine, University of Calgary, Calgary, Alberta, Canada

## Abstract

This study examined physicians' attitudes toward advanced directives and practices for the end-of-life care at Chiang Mai University Hospital, Thailand. The data were collected from 55 physicians (24 instructors and 31 residents) using self-reported questionnaires. The majority of the participants affirmed the usefulness of the advance directive (AD) for cardiopulmonary resuscitation and respected the patients' wish for this directive, although advanced end-of-life care and resuscitation planning with the patients was limited. Mostly, the relatives were consulted regarding ADs. This study suggests that, in traditional Thai culture, physicians and families are more inclined to make decisions for the patient when they feel that it is in the patient's best interest. Further research is needed to investigate how and to what extent such attitudes can affect medical practice for end-of-life care in the context of the rapid development and consequent changes taking place in Thailand.

## Key words

advanced directive, physician's attitude, terminal care, terminally ill patient.

## INTRODUCTION

Advances in medical technology have contributed to extraordinary cures and recovery in patients around the world (Greenwald, 2007). In some circumstances, however, these advances have created problems, particularly for terminally ill patients (Levenson *et al.*, 2000; Whitcomb & Blackman, 2007). Aggressive and costly technology is often used to extend life (Somogyi-Zalud *et al.*, 2002; Grossman, 2005) and, in many cases, prolong the suffering of terminally ill patients (Reynolds *et al.*, 2005). Caring for dying patients appears to be one of the most difficult challenges of modern medicine. Decisions to limit treatment are an increasingly common feature in the clinical management of patients towards the end of life (Holzapfel *et al.*, 2002; Braun *et al.*, 2007; Szalados, 2007).

By providing a means for competent persons to guide their medical care after they have become incompetent, advanced directives (ADs) allow patients to influence treatment decisions by expressing a personal view on the balance between the quality and duration of life. Advanced directives are documents that enable people to decide what will happen to

them or who will make decisions on their behalf if the time comes when they are no longer able to participate in discussions or decisions about their own health care (Ramsey & Mitty, 2003). In the USA, the Patient Self-determination Act of 1991 requires hospitals and nursing homes to inform patients of their right to refuse medical treatment (La Puma *et al.*, 1991). In this circumstance, ADs serve as a legal document that allow competent people to give instructions regarding the health care they would like to receive during a time of crisis when they will not be competent to make their own decisions (Fisher, 2008). The use of ADs constitutes a part of the whole process of assisting patients to understand their condition, options for medical treatment, and the potential complications. It also involves discussing the available choices with patients and their family and reflecting on those choices in light of personal values, goals, and religious or cultural beliefs. Such advance planning is helpful for patients who might lose the ability to speak for themselves because of a progressive and/or serious illness.

The initiation of the discussion on end-of-life care planning is especially important for patients with a terminal illness. These patients seem to be precisely those expected to benefit from earlier discussions of the merits of their terminal care. Unfortunately, these means have been underutilized (Quill, 2000; Gallo *et al.*, 2003; Booth *et al.*, 2004; Dougherty *et al.*, 2007). A recent report in Thailand showed that the use of

Correspondence address: Sudarat Sittisombut, Faculty of Nursing, Chiang Mai University, 110 Intawaroros Street, Chiang Mai 50200, Thailand. Email: sudarat@mail.nurse.cmu.ac.th

Received 30 May 2008; accepted 24 July 2008

ADs appears to be effective in reducing futile cardiopulmonary resuscitation (CPR) attempts and the in-hospital mortality rate among terminally ill patients (Sittisombut *et al.*, 2008).

Although ADs are widely advocated in Western countries, such as the USA, Canada, UK, the Netherlands, and Switzerland, only a few studies dealing with ADs in Asian countries have been reported (Low *et al.*, 2000; Miura *et al.*, 2001; Akabayashi *et al.*, 2003; Sittisombut *et al.*, 2005). Most of these studies focused on the attitudes of patients towards ADs; thus, little is known concerning the attitudes of Asian physicians (Tee *et al.*, 1997; Asai *et al.*, 1998). The purpose of this study was to examine the attitudes and practices of Thai physicians regarding the end-of-life care planning for terminally ill patients in an acute care medical center in northern Thailand.

## METHODS

This was a descriptive study utilizing questionnaires that were developed from the literature that reflected the objectives of the study. The questions were focused on the provision of information about terminal illness, the attitudes towards ADs for CPR, and the experiences of physicians in using ADs in clinical practice. After the content validity of the questionnaires was approved by five local experts, the questionnaires were tested in a pilot study with 10 physicians who had similar characteristics to the study's sample. The inter-rater reliability, as assessed by using Cohen's Kappa statistics, was 0.85.

## Ethical considerations

This study was a part of a larger research project on the use of ADs in the care of terminally ill patients at Chiang Mai University Hospital, Thailand (Sittisombut *et al.*, 2008). The research was approved by the institutional review board of the Faculty of Medicine, Chiang Mai University. Participation in the study was voluntary and anonymous, and all the participants signed an informed consent form before the commencement of the study. The data were adequately protected by utilizing a code system: no names were used and the data were stored in a secure locker, which could be accessed only by the research team.

## Recruitment of the participants

Physicians who were working and caring for terminally ill patients in the Medical Department, Chiang Mai University Hospital, at the time of the study and who agreed to participate by giving informed consent were eligible for the study. There was a total of 106 physicians working in the Medical Department during the study period. Forty-four of these physicians were faculty members (instructors) and 62 were in the residential training program. Eleven faculty members were out of the country during the study period, leaving a total of 95 physicians eligible for participation in the study. The physicians were informed of the objectives and methods and then invited to participate in the study. The questionnaires

were distributed by the investigator. The completed questionnaires were returned by the physicians at their convenience.

## Statistical analysis

The demographic characteristics of the participants and the responses to the questionnaires were analyzed using descriptive statistics. Chi-squared statistics were used to test the differences in the demographic characteristics according to the employment position.

## RESULTS

### Response rate

Of the 95 potential participants, one declined to participate, three had not been involved recently in terminal care, and 36 did not return the questionnaires. Among 55 (57.9%) of the eligible physicians who participated in the study, there were 24 instructors and 31 residents. The difference in the response rate between the instructors and residents was significant at  $P = 0.03$ .

### Demographic characteristics

The participants were predominantly male (61.8%) and in the age range of 23–57 years (Table 1). More than half of the participants were  $\leq 30$  years of age. The majority (67.3%) were single. The participants' educational backgrounds include a Doctor of Medicine (MD) degree (56.4%), and a MD degree with the completion of a Board of Internal Medicine training program (40%). Two of the participants held a PhD degree in addition to a MD degree and the completion of the Board of Internal Medicine training program. Approximately half (54.5%) of the participants had  $\geq 3$  years of clinical experience. Significant differences were noted between the faculty members and the residents in all the demographic variables, including age, gender, marital status, education, and working experience (Table 1).

### Provision of information on terminal illness

In the study hospital, information about terminal illness was normally provided by the physicians to the patients' immediate family members when the patients were incapacitated and unable to communicate. In cases where the patients remained capable of communicating, 41.8% of the participating physicians indicated that they still preferred to inform a relative about the patients' condition (Table 2). The instructors were more likely to prefer telling a relative than the residents (45.8% vs 38.7%). Only 20% of the participants preferred to inform the patients only, whereas 36.4% preferred to inform both the patients and their relatives.

On their care of patients with a terminal illness in the past, 21 participants (38.2%) stated that they had informed the "majority" of their patients about their illness (Table 2). Only 11 participants (20.0%) had told all of their patients about their illness, while six participants (10.9%) stated that they had provided this information in "some cases". In contrast, 45

**Table 1.** Demographic characteristics stratified by the employment position

Characteristic	Total ( <i>n</i> = 55) N (%)	Instructor ( <i>n</i> = 24) N (%)	Resident ( <i>n</i> = 31) N (%)	<i>P</i> -value
Gender				0.020
Male	34 (61.8)	19 (79.2)	15 (48.4)	
Female	21 (38.2)	5 (20.8)	16 (51.6)	
Age (years)				0.001
23–30	31 (56.4)	3 (12.5)	28 (90.3)	
> 30	14 (25.5)	14 (58.3)	–	
Missing data	10 (18.1)	7 (29.2)	3 (9.7)	
Marital status				0.001
Single	37 (67.3)	7 (29.2)	30 (96.8)	
Married	18 (32.7)	17 (70.8)	1 (3.2)	
Education				0.001
MD degree	31 (56.4)	– (0.0)	31 (100.0)	
Higher than MD degree	24 (43.6)	24 (100.0)	– (0.0)	
Work experience				0.001
≤ 3 years	25 (45.5)	3 (12.5)	22 (71.0)	
> 3 years	30 (54.5)	21 (87.5)	9 (29.0)	

MD, Doctor of Medicine.

**Table 2.** Participants' responses to questions informing the patient and/or relative about the diagnosis and prognosis in cases where the patient was capable

Question/response	Total ( <i>n</i> = 55) N (%)	Instructor total ( <i>n</i> = 24) N (%)	Resident total ( <i>n</i> = 31) N (%)	<i>P</i> -value
1. Which person would you prefer to inform about the diagnosis and prognosis?				0.873
(a) Patient	11 (20.0)	5 (20.8)	6 (19.4)	
(b) Relative	23 (41.8)	11 (45.8)	12 (38.7)	
(c) Both the patient and relative	20 (36.4)	8 (33.4)	12 (38.7)	
(d) No answer	1 (1.8)	–	1 (3.2)	
2. How often have you informed the patient about the terminal diagnosis and prognosis?				0.921
(a) Every patient	11 (20.0)	3 (12.5)	8 (25.8)	
(b) The majority of the patients	21 (38.2)	8 (33.4)	13 (41.9)	
(c) Some patients	6 (10.9)	3 (12.5)	3 (9.7)	
(d) None of the patients	4 (7.3)	3 (12.5)	1 (3.2)	
(e) Other	2 (3.6)	2 (8.3)	–	
(f) No answer	11 (20.0)	5 (20.8)	6 (19.4)	
3. How often have you informed the relative about the terminal diagnosis and prognosis of the patient?				0.190
(a) Every case	45 (81.8)	18 (75.0)	27 (87.1)	
(b) The majority of the cases	7 (12.8)	4 (16.7)	3 (9.7)	
(c) Some cases	2 (3.6)	2 (8.3)	–	
(d) No answer	1 (1.8)	–	1 (3.2)	

participants (81.8%) responded that they informed the immediate relatives in every case. There was no significant difference between the faculty members and the residents with regard to providing information about the patients' illness (Table 2).

### Attitudes toward advanced directives for cardiopulmonary resuscitation

Only one-half of the participating physicians (50.9%) thought that it was appropriate to inform all terminally ill

patients about CPR and to allow them to make decisions in advance (Table 3). Fourteen participants (25.5%) preferred to decide on a case-by-case basis by considering the physical and psychological conditions, religion, and the cultural background of each individual. One-fifth (20.0%) did not agree that information on CPR should be discussed with terminally ill patients. No statistically significant difference was observed between the instructors and the residents in the responses to this question.

In their responses to the question, "Within the Thai cultural environment, is it a good idea to provide CPR

**Table 3.** Participants' responses to questions regarding providing patients information about cardiopulmonary resuscitation (CPR)

Question/response	Total ( <i>n</i> = 55) N (%)	Instructor total ( <i>n</i> = 24) N (%)	Resident total ( <i>n</i> = 31) N (%)	<i>P</i> -value
1. Is it appropriate to inform all terminally ill patients about CPR and allow them to decide in advance?				0.215
(a) Yes	28 (50.9)	9 (37.5)	19 (61.3)	
(b) No	11 (20.0)	6 (25.0)	5 (16.2)	
(c) Other	14 (25.5)	8 (33.4)	6 (19.3)	
(d) No answer	2 (3.6)	1 (4.1)	1 (3.2)	
2. In Thai culture, is it a good idea to provide CPR information to all admitted patients and allow them to decide about CPR in advance?				0.005
(a) Yes	19 (34.5)	3 (12.5)	16 (51.6)	
(b) No	17 (31.0)	6 (25.0)	11 (35.5)	
(c) Other	19 (34.5)	15 (62.5)	4 (12.9)	

**Table 4.** Participants' responses to questions regarding their behavior in relation to cardiopulmonary resuscitation (CPR)

Question/response	Total ( <i>n</i> = 55) N (%)	Instructor total ( <i>n</i> = 24) N (%)	Resident total ( <i>n</i> = 31) N (%)	<i>P</i> -value
1. Have you ever asked patients who were terminally ill whether or not they wanted to have CPR performed?				0.292
(a) Yes	19 (34.5)	10 (41.7)	9 (29.0)	
(b) No	34 (61.8)	14 (58.3)	20 (64.5)	
(c) Other	2 (3.6)	–	2 (6.5)	
2. Have you ever asked the relatives of a terminally ill patient whether or not they wanted to have CPR performed for the patient?				0.183
(a) Yes	52 (94.6)	22 (91.7)	30 (96.8)	
(b) No	2 (3.6)	2 (8.3)	–	
(c) Other	1 (1.8)	–	1 (3.2)	
3. If the patients with terminal illnesses request that they do not want to have CPR, would you prefer to follow their wishes?				0.063
(a) Yes	50 (90.9)	24 (100.0)	26 (83.9)	
(b) Other	4 (7.3)	–	4 (12.9)	
(c) No answer	1 (1.8)	–	1 (3.2)	

information to all admitted patients and allow them to decide about CPR in advance?", about one-third (34.5%) of the participants thought it was appropriate to inform all admitted patients regarding CPR and to allow them to make advance decisions (Table 3). Although another one-third (31.0%) responded that it was not appropriate, the remainder (34.5%) stated that their responses to this question depended on the individual patient. The residents (51.6%) were more likely to respond positively to this question than the instructors (12.5%) and there was a statistically significant difference between the two groups ( $P = 0.005$ ).

### Use of advanced directives in medical practice

More than one-half (61.8%) of the participating physicians had never asked their terminally ill patients whether they wanted to have CPR performed or not (Table 4). However,

the majority (94.5%) had discussed this issue with the patients' relative(s). If the patients with a terminal illness were to specify that they did not want to have CPR, the majority of the participants (90.9%) stated that they would follow their wishes. There was no significant difference between the instructors and the residents on this response (Table 4).

### DISCUSSION

Although most physicians accept the ethical proposition that patients are entitled to know their prognosis (Hancock *et al.*, 2007), the provision of such important information appears to be different among physicians in various social and cultural backgrounds. A small proportion of Thai physicians participating in this study preferred to inform the terminally ill patients about their diagnosis and prognosis. Our findings

were in accordance with previous studies conducted in Asian countries on physicians' attitudes. Only 13% (Tinada, 1994) and 31% of Japanese physicians (Mizushima *et al.*, 1990) and no Chinese physicians (Feldman *et al.*, 1999) would inform their patients with cancer of their diagnosis. Similarly, Jiang *et al.* (2006) reported that only 40.5% of Chinese oncologists believe that patients with a terminal illness should know the truth about their diagnosis. A high rate of non-disclosure of a cancer diagnosis appears to be common among physicians in Asia (Back & Huak, 2005). However, this might reflect a general attitude prevalent in their communities. Families are known to play a significant role in ethical decision-making for patients in India (Chattopadhyay & Simon, 2008). There is evidence to indicate that families are strongly resistant to informing patients about their diagnosis (Wang *et al.*, 2004). A study in Japan found that only 4.3% of families consent to the physician notifying patients about their diagnosis (Tazawa *et al.*, 1990). Some studies revealed that Asian patients themselves do not wish to be told the truth (Seo *et al.*, 2000; Elwyn *et al.*, 2002).

Similar to previous reports from the USA (La Puma *et al.*, 1991), the majority of Thai physicians in this study expressed a positive attitude towards ADs and respect for the patient's wishes regarding CPR. Yet, only one-third of them agreed with discussing advanced CPR planning with all admitted patients and one-half supported the offering of an AD for CPR to terminally ill patients. This attitude of Thai physicians does not reflect those of the patients. In the same health-care setting, the majority of ambulatory patients thought that it was a good idea to discuss advance planning for CPR on a routine basis with all admitted patients (Sittisombut *et al.*, 2005). Although they were still able to communicate, terminally ill patients were also in favor of using an AD to determine their preference for CPR (Sittisombut *et al.*, 2008). These studies suggest that, in contrast to the physicians' attitude, Thai patients are receptive to such discussions well before they become unable to participate in the decision-making process.

An interesting finding from this study indicates that, when it comes to ADs, Thai physicians tend to give greater weight to the decisions of family members than to those of the patients. While one-third of the participants reported asking terminally ill patients whether or not they wanted to have CPR performed, almost all of them raised this question with the patients' relatives. Most of the participating physicians felt that discussing this sensitive issue with the patients was time-consuming and might be harmful to certain patients. These findings are similar to those from previous studies of physicians in Japan (Asai *et al.*, 1998) and Singapore (Tee *et al.*, 1997), but they differ from similar studies in the USA and Canada, where physicians are more likely to give patients a greater voice in their own decisions (Bruera *et al.*, 2000; Ruhnke *et al.*, 2000). In a study by Bruera *et al.* (2000), the physicians surveyed in Europe, South America, and Canada agreed that "do not resuscitate" orders should be discussed with the terminally ill patient in all cases. A survey in the USA found that the residents discussed the patient's preferences for CPR with most patients on the day of admission (Smith *et al.*, 2006).

It is not clear yet whether the physicians' preference actually contributes to the high rate of CPR in Chiang Mai University Hospital (Sittisombut *et al.*, 2001). Currently, when terminally ill patients are provided with the information concerning their prognosis and the possibility of CPR during their hospitalization, many of the northern Thai patients do wish to make an advanced decision and the use of the AD appears to be effective in reducing futile CPR attempts (Sittisombut *et al.*, 2008).

## CONCLUSIONS

The data from this study suggest that cultural differences shape physicians' opinions and practices concerning ADs with terminally ill patients. Greater emphasis on individual autonomy might result in more collaboration between the physicians and the patients with respect to ADs and end-of-life decisions, but traditional Thai culture appears to permit the physicians and the families to make decisions on behalf of the patients when they feel it is in the patient's best interest. However, recent reports suggest that Thai patients are willing to participate in the end-of-life decisions. Further research is needed to clarify the effects of culture on end-of-life care, particularly in the context of the rapid development and consequent changes taking place in Thailand.

## ACKNOWLEDGMENT

This study was supported by the Golden Jubilee Scholarship program of the Thailand Research Fund.

## REFERENCES

- Akabayashi A, Slingsby BT, Kai I. Perspectives on advance directives in Japanese society: a population-based questionnaire survey. *BMC Med. Ethics* 2003; **31**: E5.
- Asai A, Miura Y, Tanabe N, Kurihara M, Fukuhara S. Advance directives and other medical decisions concerning the end of life in cancer patients in Japan. *Eur. J. Cancer* 1998; **34**: 1582–1586.
- Back MF, Huak CY. Family centred decision making and non-disclosure of diagnosis in a South East Asian oncology practice. *Psychooncology* 2005; **14**: 1052–1059.
- Booth MG, Doherty P, Fairgrieve R, Kinsella J. Relatives' knowledge of decision making in intensive care. *J. Med. Ethics* 2004; **30**: 459–461.
- Braun UK, Beyth RJ, Ford ME, McCullough LB. Analysis: defining limits in care of terminally ill patients. *BMJ* 2007; **334**: 239–241.
- Bruera E, Neumann CM, Mazzocato C, Stiefel F, Sala R. Attitudes and beliefs of palliative care physicians regarding communication with terminally ill cancer patients. *Palliat. Med.* 2000; **14**: 287–298.
- Chattopadhyay S, Simon A. East meets West: cross-cultural perspective in end-of-life decision making from Indian and German viewpoints. *Med. Health Care Philos.* 2008; **11**: 165–174.
- Dougherty CM, Pyper GP, Au DH, Levy WC, Sullivan MD. Drifting in a shrinking future: living with advanced heart failure. *J. Cardiovasc. Nurs.* 2007; **22**: 480–487.
- Elwyn TS, Fetter MD, Sasaki H, Tsuda T. Responsibility and cancer disclosure in Japan. *Soc. Sci. Med.* 2002; **54**: 281–293.
- Feldman MD, Zhang J, Cummings SR. Chinese and U.S. internists adhere to different ethical standards. *J. Gen. Intern. Med.* 1999; **14**: 469–473.

- Fisher D. *National Empowerment Center – Articles Making Advance Directives Work for You*. Lawrence, MA, USA: National Empowerment Center, 2008.
- Gallo JJ, Straton JB, Klag MJ *et al.* Life-sustaining treatments: what do physicians want and do they express their wishes to others? *J. Am. Geriatr. Soc.* 2003; **51**: 961–969.
- Greenwald P. A favorable view: progress in cancer prevention and screening. *Recent Results Cancer Res.* 2007; **174**: 3–17.
- Grossman T. Latest advances in antiaging medicine. *Keio J. Med.* 2005; **54**: 85–94.
- Hancock K, Clayton JM, Parker SM *et al.* Truth-telling in discussing prognosis in advanced life-limiting illnesses: a systematic review. *Palliat. Med.* 2007; **21**: 507–517.
- Holzapfel L, Demingon G, Piralla B, Biot L, Nallet B. A four-step protocol for limitation of treatment in terminal care: an observational study in 475 intensive care unit patients. *Intensive Care Med.* 2002; **28**: 1309–1315.
- Jiang Y, Li JY, Liu C *et al.* Different attitudes of oncology clinicians toward truth telling of different stages of cancer. *Support. Care Cancer* 2006; **14**: 1119–1125.
- La Puma J, Orentlicher D, Moss RJ. Advance directives on admission: clinical implications and analysis of the Patient Self-determination Act. *JAMA* 1991; **266**: 402–405.
- Levenson JW, McCarthy EP, Lynn J, Davis RB, Phillip RS. The last six months of life for patients with congestive heart failure. *J. Am. Geriatr. Soc.* 2000; **48**: S101–S109.
- Low JA, Ng WC, Yap KB, Chan KM. End-of-life issues: preferences and choices of a group of elderly Chinese subjects attending a day care center in Singapore. *Ann. Acad. Med. Singapore* 2000; **29**: 50–56.
- Miura Y, Asai A, Nagata S *et al.* Dialysis patients' preferences regarding cardiopulmonary resuscitation and withdrawal of dialysis in Japan. *Am. J. Kidney Dis.* 2001; **37**: 1216–1222.
- Mizushima Y, Kashii T, Hoshino K *et al.* A survey regarding the disclosure of the diagnosis of cancer in Toyama prefecture, Japan. *Jpn. J. Med.* 1990; **29**: 146–155.
- Quill TE. Perspectives on care at the close of life. Initiating end-of-life discussions with seriously ill patients: addressing the “elephant in the room”. *JAMA* 2000; **284**: 2502–2507.
- Ramsey G, Mitty E. Advance directives: protecting patient's rights. In: Mezey M, Fulmer T, Abraham I, Zwicker DA (eds). *Geriatric Nursing Protocols for Best Practice* (2nd edn). New York: Springer, 2003; 265–291.
- Reynolds S, Cooper AB, McKneally M. Withdrawing life-sustaining treatment: ethical considerations. *Thorac. Surg. Clin.* 2005; **15**: 469–480.
- Ruhnke GW, Wilson SR, Akamatsu T *et al.* Ethical decision making and patient autonomy: a comparison of physicians and patients in Japan and the United States. *Chest* 2000; **118**: 1172–1182.
- Seo M, Tamura K, Shijo H, Morioka E, Ikegame C, Hirasako K. Telling the diagnosis to cancer patients in Japan: attitude and perception of patients, physicians and nurses. *Palliat. Med.* 2000; **14**: 105–110.
- Sittisombut S, Love EJ, Sitthi-Amorn C. Cardiopulmonary resuscitation performed in patients with terminal illness in Chiang Mai University Hospital, Thailand. *Int. J. Epidemiol.* 2001; **30**: 896–898.
- Sittisombut S, Love EJ, Sitthi-Amorn C. Attitudes toward advance directives and the impact of prognostic information on the preference for cardiopulmonary resuscitation in medical inpatients in Chiang Mai University Hospital, Thailand. *Nurs. Health Sci.* 2005; **7**: 243–250.
- Sittisombut S, Maxwell C, Love EJ, Sitthi-Amorn C. Effectiveness of advance directives for the care of terminally ill patients in Chiang Mai University Hospital, Thailand. *Nurs. Health Sci.* 2008; **10**: 37–42.
- Smith AK, Ries AP, Zhang B, Tulsy JA, Prigerson HG, Block SD. Resident approaches to advance care planning on the day of hospital admission. *Arch. Intern. Med.* 2006; **166**: 1597–1602.
- Somogyi-Zalud M, Zhong Z, Hamel MB, Lynn J. The use of life-sustaining treatments in hospitalized persons aged 80 and older. *J. Am. Geriatr. Soc.* 2002; **50**: 930–934.
- Szalados JE. Discontinuation of mechanical ventilation at end-of-life: the ethical and legal boundaries of physician conduct in termination of life support. *Crit. Care Clin.* 2007; **23**: 317–337.
- Tazawa H, Satoh I, Ishihara J *et al.* Informed consent of the family in the chemotherapy for lung cancer. *Nippon Gan. Chiryo. Gakkai Shi.* 1990; **25**: 1448–1453.
- Tee KH, Seet LT, Tan WC, Choo HW. Advance directive: a study on the knowledge and attitudes among general practitioners in Singapore. *Singapore Med. J.* 1997; **38**: 145–148.
- Tinada N. Japanese attitudes towards truth disclosure in cancer. *Scand. J. Soc. Med.* 1994; **22**: 50–57.
- Wang XS, Di LJ, Reyes-Gibby CC, Guo H, Liu SJ, Cleeland CS. End-of-life care in urban areas of China: a survey of 60 oncology clinicians. *J. Pain Symptom Manage.* 2004; **27**: 125–132.
- Whitcomb JJ, Blackman VS. Cardiopulmonary resuscitation: how far have we come? *Dimens. Crit. Care Nurs.* 2007; **26**: 1–8.

Research article

Open Access

# Comprehensibility, reliability, validity, and responsiveness of the Thai version of the Health Assessment Questionnaire in Thai patients with rheumatoid arthritis

Manathip Osiri<sup>1</sup>, Jeerapat Wongchinsri<sup>2</sup>, Sitthichai Ukritchon<sup>3</sup>, Punchong Hanvivadhanakul<sup>4</sup>, Nuntana Kasitanon<sup>5</sup> and Boonjing Siripaitoon<sup>6</sup>

<sup>1</sup>Division of Rheumatology, Department of Medicine, Faculty of Medicine, Chulalongkorn University, 1873 Rama IV Road, Pathumwan, Bangkok 10330, Thailand

<sup>2</sup>Department of Medicine, Nopparat Rajathani Hospital, 109 Ramindra Km, 12 Kunnayao, Bangkok 10230, Thailand

<sup>3</sup>Division of Rheumatology, Department of Medicine, Srinakharinwirot University, Princess Maha Chakri Sirindhorn Medical Center, 63 Moo 7, Rangsit-Ongkharak Road, Ongkharak, Nakorn-Nayok 26120, Thailand

<sup>4</sup>Division of Rheumatology, Department of Medicine, Faculty of Medicine, Thammasat University, Paholyotin Road, Amphur Klongluang, Pathumthani 12120, Thailand

<sup>5</sup>Division of Rheumatology, Department of Medicine, Faculty of Medicine, Chiang Mai University, 110 Intawaroros Road, Amphur Muang, Chiang Mai 50200, Thailand

<sup>6</sup>Division of Rheumatology, Department of Medicine, Faculty of Medicine, Prince of Songkla University, 15 Karnjanavanit Road, Hat Yai, Songkhla 90110, Thailand

Corresponding author: Manathip Osiri, [manathip.o@chula.ac.th](mailto:manathip.o@chula.ac.th)

Received: 17 Mar 2009 Revisions requested: 15 May 2009 Revisions received: 10 Aug 2009 Accepted: 27 Aug 2009 Published: 27 Aug 2009

*Arthritis Research & Therapy* 2009, **11**:R129 (doi:10.1186/ar2796)

This article is online at: <http://arthritis-research.com/content/11/4/R129>

© 2009 Osiri et al.; licensee BioMed Central Ltd.

This is an open access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

## Abstract

**Introduction** The Health Assessment Questionnaire Disability Index (HAQ-DI) is a commonly used instrument to assess functional status of patients with rheumatoid arthritis (RA). Translations and adaptations of the HAQ-DI have been carried out for use with RA patients in several countries. The objective of this study was to evaluate the psychometric properties of the Thai version of the HAQ-DI (Thai HAQ) in Thai patients with RA.

**Methods** Comprehensibility of the Thai HAQ was assessed by 126 patients with RA from 6 medical centers in Thailand. Another group of 115 patients with active RA was enrolled to test the reliability (internal reliability and 1-week test-retest reliability), construct validity (correlations with other measures of RA disease activity), floor and ceiling effects, and sensitivity to change of the Thai HAQ at 3 months of treatment with disease-modifying antirheumatic drugs.

**Results** More than 98% of the patients regarded the Thai HAQ as comprehensible. The internal consistency of the Thai HAQ was satisfactory with the overall Cronbach alpha of 0.91. The test-retest reliability of the Thai HAQ was acceptable with the intraclass correlation coefficient of 0.89. Moderate correlations between the Thai HAQ and other outcomes of RA disease activity were observed, except erythrocyte sedimentation rate, with the Spearman correlation coefficients ranging from 0.42 to 0.57. The responsiveness of the Thai HAQ was moderate, with a standardized response mean of 0.75 (95% confidence interval 0.56 to 0.94).

**Conclusions** The Thai HAQ is comprehensible, reliable, valid and sensitive to change in the evaluation of functional status of Thai patients with RA. The Thai HAQ is an essential tool to measure treatment effects and progression of disability in RA patients and should be applied in both clinical trials and routine clinical care settings.

ACR: American College of Rheumatology; CI: confidence interval; DMARD: disease-modifying antirheumatic drug; ESR: erythrocyte sedimentation rate; HAQ: Health Assessment Questionnaire; HAQ-DI: Health Assessment Questionnaire Disability Index; ICC: intraclass correlation coefficient; OMERACT: Outcome Measures in Rheumatology Clinical Trials; RA: rheumatoid arthritis; SD: standard deviation; SRM: standardized response mean; Thai HAQ: Thai version of the Health Assessment Questionnaire Disability Index.

## Introduction

The Stanford Health Assessment Questionnaire (HAQ) was originally developed to measure five important health outcomes in patients with chronic diseases [1,2]. These dimensions include premature death, functional disability, pain and discomfort, adverse effects of treatment, and costs. The HAQ Disability Index (HAQ-DI), the HAQ section to evaluate functional capacity, is the most commonly used instrument for assessing disability in patients with rheumatoid arthritis (RA). The HAQ-DI is a predictive factor of future disability and joint damage in patients with RA [3-5]. Because it demonstrated sensitivity to change, the HAQ-DI was chosen by the Outcome Measures in Rheumatology Clinical Trials (OMERACT) and the American College of Rheumatology (ACR) to be incorporated into the core set of outcome measures of RA disease activity [6-8]. The HAQ-DI not only is considered an essential measure of disability in patients with RA in clinical trials, but also is used in clinical practice.

The HAQ-DI has been translated and adapted to suit the activities and cultures in diverse populations from more than 50 countries [9,10]. As with the original HAQ-DI, a number of translations of the HAQ-DI into other languages have been proven to be reliable, valid, and sensitive to change. For the Thai version of the HAQ-DI (Thai HAQ), three items were adapted and two activities were added to the existing items to tailor the questionnaire to the lifestyle and culture of Thai people. The Thai HAQ was back-translated and tested for its validity and responsiveness in a pilot study at a tertiary care hospital in Thailand [11]. However, psychometric validation of the Thai HAQ is still important if it is to be recommended as a standard instrument to measure long-term disability in Thai patients with RA. It is also needed for demonstrating the effectiveness of disease-modifying antirheumatic drug (DMARD) therapy, especially with the expensive biologics, and for use in guidelines to follow the patients over time. Thus, the objective of this study was to evaluate the comprehensiveness, reliability, validity, and responsiveness of the Thai HAQ in Thai-speaking patients with RA from different parts of the country.

## Materials and methods

### Comprehensibility

One hundred twenty-six adult patients who met the ACR 1987 revised criteria for the classification of RA [12] were included in the comprehensibility assessment of the Thai HAQ. These patients were enrolled from six medical centers in Thailand from January to April 2006 regardless of their disease activity. The comprehensibility questionnaire was self-reported by the patients. For older patients and those with poor eyesight, the responses were performed with the assistance of relatives who accompanied them and of rheumatology nurses at the clinics. The patients were asked whether they understood the 20 items from the eight domains of the Thai HAQ. The levels of comprehensibility for each item were categorized on a 4-

point scale (0, not comprehensible; 1, slightly comprehensible; 2, moderately comprehensible; and 3, highly comprehensible). Scores of 2 or more for each item were regarded as comprehensible.

### Reliability, validity, and responsiveness

From the same six institutes, 115 more patients with RA were consecutively selected between January 2006 and July 2007 to be included in this part of study. Each patient had to fulfill all of the following criteria: (a) met the ACR 1987 revised criteria for the classification of RA [12], (b) was at least 18 years of age, and (c) had active disease characterized by (i) at least six tender joints, (ii) at least six swollen joints, (iii) a Westergren erythrocyte sedimentation rate (ESR) of at least 28 mm/hour, and (iv) at study entry, just starting a non-biologic DMARD, an increase in the dose of a non-biologic DMARD, or the addition of another non-biologic DMARD. The patients were excluded if they were pregnant or breastfeeding, receiving prednisolone at a dose of more than 10 mg per day, or did not give informed consent. The mode of Thai HAQ administration was similar to that of the comprehensibility assessment. The assessments of reliability, validity, and responsiveness of the Thai HAQ were conducted in accordance with the OMERACT filter for outcome measures in RA [8]. Both parts of the study protocols were approved by the ethics committees of each institute. This study was conducted in accordance with the Declaration of Helsinki. All patients were required to give written informed consent before entering this study.

### The Thai HAQ

The Thai HAQ included 20 items from eight domains adapted from the original HAQ-DI to suit Thai culture and activities. The ability to perform an activity for each item is rated on a 4-level scale, in which the score ranges from 0 (no difficulty in performing that activity) to 3 (inability to perform that activity). The requirement of a device or physical assistance in any item increases the lower score to 2. To calculate the HAQ-DI score, the maximum scores from each domain were summed and divided by 8 to yield a score that ranged from 0 to 3. The higher score indicated greater disability.

### Statistical analysis

Baseline characteristics of the studied patients were presented as number and percentage for discrete parameters and as mean and standard deviation (SD) for continuous parameters. Comprehensibility assessment of the Thai HAQ was presented as percentage of patients with moderate or high comprehensibility in the Thai HAQ.

### Reliability

Reliability was assessed by test-retest reliability and internal consistency. The test-retest reliability was performed with a 1-week interval. This interval was used because the patients would not be able to remember the first test and the effects of DMARDs added were not expected at 1 week of treatment.

The patients completed the first Thai HAQ at their clinic visits. The second was sent to them by mail. Test-retest reliability of the Thai HAQ was analyzed using intraclass correlation coefficients (ICCs). The ICCs and their 95% confidence intervals (CIs) were calculated using a two-way random-effects model. An ICC value of 0.85 or higher was considered acceptable [13]. Internal consistency among each domain of the Thai HAQ was evaluated by Cronbach alpha using the results from the first administration. The overall Cronbach alpha was calculated from all eight domains of the Thai HAQ. For each domain, Cronbach alpha was obtained by deleting that domain from the questionnaire [14].

### Validity

Construct validity of the Thai HAQ was performed by correlating the baseline eight domains and total scores of the Thai HAQ with the following outcome measures of RA disease activity: number of tender joints (total 68 joints), number of swollen joints (66 joints), patient's assessment of pain, patient global assessment of disease activity, physician global assessment of disease activity, and ESR. Pain score and patient and physician global assessments of disease activity were measured on a 5-level categorical scale, in which the higher score indicated greater pain and worse disease status. The correlation coefficients used in this study were Spearman correlation coefficients. Correlation coefficients of greater than 0.6, of 0.6 to 0.3, and of less than 0.3 were considered strong, moderate, and weak correlations, respectively [15]. The Thai HAQ was also evaluated for floor and ceiling effects. Floor and ceiling effects were considered to be present if at least 15% of the patients scored 0 (the lowest possible score) or 3 (the highest possible score), respectively, on the Thai HAQ [16].

### Responsiveness

Responsiveness of the Thai HAQ and other measures of RA disease activity was calculated from the baseline values and the values at month 3. To assess the responsiveness of the Thai HAQ and other parameters, the differences between baseline and month 3 scores were used for calculating the standardized response mean (SRM) from the formula:

$$\text{SRM} = \text{mean change of the score} / \text{SD change of the score}.$$

The SRM value between 0.6 and 0.8 is considered moderate effect and clinically significant. The value of greater than 0.8 represents large effect [17,18]. The statistical software used in this study was SPSS for Windows, version 11.0 (SPSS Inc., Chicago, IL, USA). The statistically significant level was determined as a *P* value of less than 0.05.

### Results

Demographic data of both groups of studied patients are shown in Table 1. The comprehensibility of the Thai HAQ was assessed by 126 patients, whose domiciles were distributed in all parts of the country. Twenty-six patients (20.6%) were from Bangkok. Among the others, 35 (27.8%) resided in Central Thailand, 23 (18.3%) in the North, 11 (8.7%) in the North-east or East, and 31 (24.6%) in the Southern part of Thailand. The mean (SD) age of the studied patients was 50.5 (13.0) years. One hundred fourteen patients (90.5%) were women. Although 68.3% of the patients (86 patients) completed only primary education (that is, 6 years of formal education), more than 90% of the 126 patients scored each item of the Thai HAQ as moderately or highly comprehensible, as shown in Table 2.

Of the 115 patients with RA enrolled to test the reliability, validity, and responsiveness of the Thai HAQ, 99 (86.1%) were women. The mean (SD) age was 48.9 (11.9) years, and

**Table 1**

#### Demographic characteristics of studied patients

Demographic characteristics	Comprehensibility assessment	Reliability, validity, responsiveness assessment
Number of patients	126	115
Females/males, number (percentage)	114/12 (90.5%/9.5%)	99/16 (86.1%/13.9%)
Age in years, mean $\pm$ SD	50.5 $\pm$ 13.0	48.9 $\pm$ 11.9
Disease duration in months, mean $\pm$ SD	ND	68.9 $\pm$ 71.8
Rheumatoid factor-positive patients, number (percentage)	ND	88 (76.5%)
Formal education, number (percentage) of patients		
$\leq 6$ years	92 (73.0%)	81 (70.4%)
$> 6$ to $\leq 12$ years	26 (20.6%)	24 (20.9%)
$> 12$ years	8 (6.4%)	10 (8.7%)

ND, no data; SD, standard deviation.

**Table 2****Comprehensibility of each item of the Thai HAQ assessed by 126 Thai patients with rheumatoid arthritis**

Thai HAQ domain	Thai HAQ item	Percentage comprehensibility
	Are you able to:	
Dressing and grooming	1. Put on clothes, including buttoning up?	100
	2. Wash your hair?	100
Arising	3. Get up from a chair without armrests?	98.4
	4. Lie down and get up from the bed, or sit in floor-sitting or kneeling position?	100
Eating	5. Slice food with a knife?	99.2
	6. Lift up a glass (filled with water) for drinking?	99.2
	7. Open up food or beverage cans?	99.2
Walking	8. Walk outdoors on level ground?	100
	9. Climb up five steps of stairs?	99.2
Hygiene	10. Apply soap over the body and towel up?	100
	11. Lift up a water bowl to wash yourself?	100
	12. Sit down and get up from a toilet seat?	100
Reach	13. Reach for a 2-kg object from an overhanging cupboard?	98.4
	14. Bend down to pick up an article from the floor?	100
Grip	15. Open a car door?	100
	16. Open containers (such as conserve or Ovaltine jar)?	99.2
	17. Turn on and off a faucet, or wring clothes after washing?	99.2
Activities	18. Go marketing?	99.2
	19. Get on and off a car or a bus?	99.2
	20. Sweep and mop?	98.4

Thai HAQ, Thai version of the Health Assessment Questionnaire Disability Index.

the mean (SD) disease duration was 68.9 (71.8) months. Rheumatoid factor was positive in 77%. The mean (SD) Thai HAQ score at baseline was 1.56 (0.75).

Internal consistency of the Thai HAQ was satisfactory, with the Cronbach alpha of 0.910 among all eight domains. Removal of each domain of the Thai HAQ did not produce a significant change in the Cronbach alpha. The highest alpha was 0.899 when the dressing domain was deleted and the lowest alpha was 0.886 when the reach or activity domain was excluded from the Thai HAQ.

Test-retest reliability of each domain and total Thai HAQ scores was acceptable. The average measure ICC of the Thai HAQ was 0.89 (95% CI 0.84 to 0.92). For each domain of the Thai HAQ, the estimates of ICC ranged from 0.77 to 0.87. The mean and SD of each domain of the Thai HAQ at baseline and day 7 as well as the ICCs and 95% CIs are shown in Table 3.

The Spearman correlation coefficients for construct validity of each domain of the Thai HAQ and total Thai HAQ scores are shown in Table 4. Moderate correlation was observed between the majority of Thai HAQ domains as well as the Thai HAQ and outcomes of RA disease activity. These correlation coefficients ranged from 0.30 to 0.57. The highest correlation coefficient was observed between the Thai HAQ and ACR functional class (correlation coefficient 0.57), whereas the lowest was seen between the Thai HAQ and ESR (correlation coefficient 0.37). ESR correlated weakly with four domains of the Thai HAQ (dressing and grooming, walking, grip, and activity domains). The grip and walking domains of the Thai HAQ also had weak correlations with the number of swollen joints, number of tender joints, and pain level.

Floor and ceiling effects of the Thai HAQ were not observed in the studied patients. At baseline, none (0%) scored 0 or 3

**Table 3****Test-retest reliability of eight domains of the Thai HAQ**

Thai HAQ domain	Mean $\pm$ SD score on day 1	Mean $\pm$ SD score on day 7	ICC (95% CI) <sup>a</sup>
Dressing and grooming	0.90 $\pm$ 0.83	0.82 $\pm$ 0.83	0.83 (0.75-0.88)
Arising	2.05 $\pm$ 1.67	1.85 $\pm$ 0.99	0.84 (0.77-0.89)
Eating	1.60 $\pm$ 1.02	1.47 $\pm$ 1.02	0.82 (0.75-0.88)
Walking	1.27 $\pm$ 0.98	1.12 $\pm$ 1.01	0.87 (0.81-0.91)
Hygiene	1.34 $\pm$ 0.80	1.22 $\pm$ 0.96	0.77 (0.66-0.84)
Reach	1.80 $\pm$ 1.07	1.61 $\pm$ 1.07	0.77 (0.67-0.84)
Grip	1.91 $\pm$ 0.97	1.62 $\pm$ 1.00	0.78 (0.69-0.85)
Activities	1.59 $\pm$ 1.01	1.33 $\pm$ 1.00	0.80 (0.70-0.86)
Thai HAQ	1.56 $\pm$ 0.75	1.38 $\pm$ 0.80	0.89 (0.84-0.92)

<sup>a</sup>Two-way random-effects average measure intraclass correlation coefficient (ICC). CI, confidence interval; SD, standard deviation; Thai HAQ, Thai version of the Health Assessment Questionnaire Disability Index.

on the Thai HAQ. At 3 months of treatment, 6 patients (5.2%) scored 0 and 1 patient (0.9%) scored 3.

Responsiveness of the Thai HAQ and other measures of RA disease activity was moderate as the SRMs were 0.6 or higher, except for ESR. The SRM for the patient global assessment of disease activity was the highest (0.94), whereas that for the Thai HAQ was 0.75. The SRMs for the Thai HAQ compared with the other measures of RA disease activity are shown in Table 5.

**Discussion**

Approximately 90% of the patients enrolled in this study were women. The high proportion of female patients with RA was observed across all six institutes (range 73.3% to 96.8%). This finding was because more women than men are affected by RA and because women with chronic rheumatic diseases in Thailand generally comply with long-term treatment better than men do.

Our study has shown that the Thai HAQ is comprehensible among Thai patients with RA recruited from different parts of Thailand. Despite the finding that more than 70% of the

**Table 4****Spearman correlation coefficients between each domain and the Thai HAQ and outcomes of rheumatoid arthritis disease activity**

Thai HAQ domain	Tender joint count	Swollen joint count	Patient global assessment	Physician global assessment	Pain	ESR	ACR functional class
Dressing	0.31	0.32	0.40	0.34	0.34	0.23 <sub>a</sub>	0.47
Arising	0.42	0.39	0.45	0.42	0.35	0.43	0.35
Eating	0.32	0.33	0.35	0.30	0.31	0.35	0.44
Walking	0.37	0.27 <sub>a</sub>	0.43	0.39	0.35	0.28 <sub>a</sub>	0.53
Hygiene	0.39	0.39	0.44	0.37	0.33	0.40	0.42
Reach	0.39	0.39	0.39	0.35	0.32	0.31	0.42
Grip	0.29 <sub>a</sub>	0.29 <sub>a</sub>	0.35	0.33	0.22 <sub>a</sub>	0.28 <sub>a</sub>	0.38
Activity	0.30	0.35	0.44	0.42	0.41	0.23 <sub>a</sub>	0.51
Thai HAQ	0.46	0.43	0.52	0.48	0.42	0.37	0.57

<sup>a</sup>Correlation coefficient of less than 0.3 represents weak correlation and 0.3 to 0.6 represents moderate correlation between the Thai HAQ (Thai version of the Health Assessment Questionnaire Disability Index) domain and rheumatoid arthritis disease activity.  $P < 0.05$  for all correlation coefficients. ACR, American College of Rheumatology; ESR, erythrocyte sedimentation rate.

**Table 5**

**Differences of treatment effect and standardized response means of rheumatoid arthritis disease activity and function outcomes between baseline and month 3 of treatment**

Rheumatoid arthritis outcome	Mean difference of treatment (95% CI)	SRM (95% CI)
Tender joint count	-3.58 (-2.42 to -4.74)	0.59 (0.40 to 0.78)
Swollen joint count	-2.72 (-1.90 to -3.53)	0.64 (0.45 to 0.83)
Patient global assessment	-0.79 (-0.63 to -0.96)	0.94 (0.74 to 1.13)
Physician global assessment	-0.77 (-0.61 to -0.94)	0.89 (0.70 to 1.09)
Pain level	-0.75 (-0.58 to -0.93)	0.82 (0.62 to 1.01)
ESR	-13.93 (-7.01 to -20.84)	0.41 (0.21 to 0.61)
Thai HAQ	-0.50 (-0.38 to -0.63)	0.75 (0.56 to 0.94)
ACR functional class	-0.44 (-0.29 to -0.60)	0.59 (0.38 to 0.80)

ACR, American College of Rheumatology; CI, confidence interval; ESR, erythrocyte sedimentation rate; SRM, standardized response mean; Thai HAQ, Thai version of the Health Assessment Questionnaire Disability Index.

patients had a limited educational level, there was no significant variation in the comprehensibility of each item of the Thai HAQ. All 126 patients regarded nine items as comprehensible. Only one and two patients, respectively, rated eight and three items of the Thai HAQ as slightly comprehensible. Among the three items with the least comprehensibility, two of these ('get up from a chair without armrests' and 'reach a 2-kg object from an overhanging cupboard') are common activities in many urbanized populations. However, neither of these activities is considered common by older Thai people who get used to living in the traditional Thai style. The other item, sweep and mop, a common activity for Thai people, was rated as slightly comprehensible by two older patients with RA who no longer performed this activity. The comprehensibility of the Thai HAQ was higher than that of the Korean HAQ, which ranged from 76% to 98% [19].

The Thai HAQ scores that included and excluded the use of aids/devices or assistance varied between 0 and 0.25 (data not shown). Use of aids/devices or other people's assistance may enhance the HAQ-DI scores [20]. In this study, most patients who required aids/devices or assistance scored that item 3, instead of 2, as they were unable to do that activity by themselves.

The Thai HAQ has been demonstrated to have satisfactory internal consistency with the Cronbach alpha of 0.910. The Cronbach alpha for each item of the Thai HAQ was also high and varied insignificantly. Our findings were comparable to the results from the HAQ-DI in other Asian countries, including South Korea [19], China [21], Kuwait [22], and Japan [23]. The Cronbach alpha in these studies ranged from 0.86 to 0.95. Test-retest reliability of the Thai HAQ has been shown to be acceptable with the ICC of 0.89. The test-retest reliability of the Thai HAQ was less than those of the Korean and Japanese HAQs but was higher than the Chinese and Arabic

HAQs. For each domain of the Thai HAQ, only the walking domain had an ICC of greater than 0.85. The mean scores of each domain of the Thai HAQ and total scores in the second administration were lower than those in the first visit. This finding was not explained by the effects of DMARDs added but might be caused by the advice from the physicians to rest the inflamed joints or the adjustment of nonsteroidal anti-inflammatory drugs and/or analgesics, which should take effect within 1 week.

The method of assessing test-retest reliability of the Thai HAQ by self-administration on the first visit and mailed response 1 week later was similar to those of the Korean and Japanese HAQ studies. Potential biases incurred from a mailed response, such as the incompleteness of filling out the questionnaire and failure to return the response, were avoided.

The Thai HAQ has been demonstrated to correlate moderately with the other measures of RA disease activity. The correlation coefficients between the Thai HAQ and patient global assessment of disease activity and ACR functional class were higher than those between the Thai HAQ and number of tender joints, number of swollen joints, physician global assessment of disease activity, and pain level. Weak correlations were found between the walking domain and number of swollen joints and between the grip domain and number of tender joints, number of swollen joints, and pain level. Newly designed devices (that is, car doors, faucets, and containers) and footwear and improvement of fabric quality (for the wring cloth item) may help improve the patient's ability to perform the activities in the grip and walking domains.

The lowest correlation coefficient was observed between the Thai HAQ and ESR. This finding corroborated with the results from the Arabic, Chinese, and Korean HAQs as ESR correlates better with RA disease duration, radiographic changes,

and joint deformity than with functional disability [20]. Three measures of RA disease activity, including patient and physician global assessments of disease activity and ACR functional class, had moderate correlation with all domains of the Thai HAQ. These correlations emphasized the importance of function on the overall health status of patients with RA.

Floor and ceiling effects of the Thai HAQ were not detected at either baseline or month 3 of treatment in this study. These effects were not investigated in the Arabic, Chinese, Japanese, or Korean HAQs. At month 3 of DMARD treatment, very few patients reported the lowest and highest possible scores. Six patients who reported the Thai HAQ score of 0 had significant improvement in their function after DMARD treatment. The only patient who scored 3 was an older patient who already had severe disability at baseline and did not respond to the increment of methotrexate dosage.

Responsiveness of the Thai HAQ measured as SRM was regarded as moderate effect and clinically significant. Large response means (SRM >0.8) were observed for the patient and physician global assessments of disease activity and for pain level. Moderate response means were found for the Thai HAQ (0.75) and number of swollen joints (0.64), whereas the number of tender joints and ACR functional class had the SRM closed to moderate effect (0.59). Our study has shown that patient-reported outcomes, including the Thai HAQ, were more efficient than physician-related outcomes, such as the numbers of tender and swollen joints, in detecting treatment effect. As expected, a small response mean was observed for the ESR (0.41).

## Conclusions

This study, together with our previous work, demonstrates that the Thai HAQ has been shown to be comprehensible, reliable, valid, and sensitive to change in detecting disability in Thai patients with RA. Our findings have confirmed the validity of using the Thai HAQ as an instrument to measure functional status of RA patients after treatment with DMARDs or biologic agents or both in clinical trials and daily practice.

## Competing interests

The authors declare that they have no competing interests.

## Authors' contributions

MO initiated the concept and design of the study and collected, analyzed, and interpreted the data and prepared the manuscript and finalized it in accordance with the recommendations. JW, SU, PH, NK, and BS collected the data. All authors read and approved the final manuscript.

## Acknowledgements

The authors thank Chitr Sitthi-amorn, Thailand Research Fund Senior Research Scholar, for providing funding for this study and Peter Tugwell for reviewing the manuscript.

## References

- Fries JF, Spitz PW, Kraines RG, Holman HR: **Measurement of patient outcome in arthritis.** *Arthritis Rheum* 1980, **23**:137-145.
- Fries JF, Spitz PW, Young DY: **The dimensions of health outcomes: the health assessment questionnaire, disability and pain scales.** *J Rheumatol* 1982, **9**:789-793.
- Cohen JD, Dougados M, Goupille P, Cantagrel A, Meyer O, Sibilia J, Daurès JP, Combe B: **Health assessment questionnaire score is the best predictor of 5-year quality of life in early rheumatoid arthritis.** *J Rheumatol* 2006, **33**:1936-1941.
- Lindqvist E, Saxne T, Geborek P, Eberhardt K: **Ten year outcome in a cohort of patients with early rheumatoid arthritis: health status, disease process, and damage.** *Ann Rheum Dis* 2002, **61**:1055-1059.
- Drossaers-Bakker KW, Zwiderman AH, Vliet Vlieland TP, Van Zeben D, Vos K, Breedveld FC, Hazes JM: **Long-term outcome in rheumatoid arthritis: a simple algorithm of baseline parameters can predict radiographic damage, disability, and disease course at 12-year followup.** *Arthritis Rheum* 2002, **47**:383-390.
- Felson DT, Anderson JJ, Boers M, Bombardier C, Chernoff M, Fried B, Furst D, Goldsmith C, Kieszak S, Lightfoot R, Paulus H, Tugwell P, Weinblatt M, Widmark R, Williams HJ, Wolfe F: **The American College of Rheumatology preliminary core set of disease activity measures for rheumatoid arthritis clinical trials.** *Arthritis Rheum* 1993, **36**:729-740.
- Boers M, Tugwell P, Felson DT, van Riel PL, Kirwan JR, Edmonds JP, Smolen JS, Khaltsev N, Muirden KD: **World Health Organization and International League of Associations for Rheumatology core endpoints for symptom modifying antirheumatic drugs in rheumatoid arthritis clinical trials.** *J Rheumatol Suppl* 1994, **41**:86-89.
- Boers M, Brooks P, Strand CV, Tugwell P: **The OMERACT filter for Outcome Measures in Rheumatology.** *J Rheumatol* 1998, **25**:198-199.
- Bruce B, Fries JF: **The Stanford Health Assessment Questionnaire: a review of its history, issues, progress, and documentation.** *J Rheumatol* 2003, **30**:167-178.
- Ramey D, Raynaud J, Fries J: **The Health Assessment Questionnaire 1992: status and review.** *Arthritis Care Res* 1992, **5**:119-129.
- Osiri M, Deesomchok U, Tugwell P: **Evaluation of functional ability of Thai patients with rheumatoid arthritis by the use of a Thai version of the Health Assessment Questionnaire.** *Rheumatology* 2001, **40**:555-558.
- Arnett FC, Edworthy SM, Bloch DA, McShane DJ, Fries JF, Cooper NS, Healey LA, Kaplan SR, Liang MH, Luthra HS, Medsger TA Jr, Mitchell DM, Neustadt DH, Pinals RS, Schaller JG, Sharp JT, Wilder RL, Hunder GG: **The American Rheumatism Association 1987 revised criteria for the classification of rheumatoid arthritis.** *Arthritis Rheum* 1988, **31**:315-324.
- Streiner DL, Norman GR: **Health Measurement Scales: A Practical Guide to Their Development and Use** 3rd edition. New York: Oxford University Press; 2005.
- Cronbach LJ: **Coefficient  $\alpha$  and the internal structure of tests.** *Psychometrika* 1951, **16**:297-334.
- Hinkle DE, Wiersma W, Jurs SG: **Applied Statistics for the Behavioral Sciences** 5th edition. Boston: Houghton Mifflin; 1998.
- McHorney CA, Tarlov AR: **Individual-patient monitoring in clinical practice: are available health status surveys adequate?** *Qual Life Res* 1995, **4**:293-307.
- Wells G, Li T, Maxwell L, Maclean R, Tugwell P: **Responsiveness of patient reported outcomes including fatigue, sleep quality, activity limitation, and quality of life following treatment with abatacept for rheumatoid arthritis.** *Ann Rheum Dis* 2008, **67**:260-265.
- Meiorin S, Pistorio A, Ravelli A, Iusan SM, Filocamo G, Trail L, Oliveira S, Cuttica R, Espada R, Alessio M, Mihaylova D, Cortis E, Martini A, Ruperto N, Paediatric Rheumatology International Trials Organisation: **Validation of the Childhood Health Assessment Questionnaire in active juvenile systemic lupus erythematosus.** *Arthritis Rheum* 2008, **59**:1112-1119.
- Bae SC, Cook EF, Kim SY: **Psychometric evaluation of a Korean Health Assessment Questionnaire for clinical research.** *J Rheumatol* 1998, **25**:1975-1979.

20. Pincus T, Sokka T: **Quantitative measures and indices to assess rheumatoid arthritis in clinical trials and clinical care.** *Rheum Dis Clin North Am* 2004, **30**:725-751.
21. Koh ET, Seow A, Pong LY, Koh WH, Chan L, Howe HS, Lim TH, Low CK: **Cross cultural adaptation and validation of the Chinese Health Assessment Questionnaire for use in rheumatoid arthritis.** *J Rheumatol* 1998, **25**:1705-1708.
22. Shebab D, al-Jarallah K, Moussa MA: **Validation of the Arabic version of the Health Assessment Questionnaire (HAQ) in patients with rheumatoid arthritis.** *Rev Rhum Engl Ed* 1998, **65**:387-392.
23. Matsuda Y, Singh G, Yamanaka H, Tanaka E, Urano W, Taniguchi A, Saito T, Hara M, Tomatsu T, Kamatani N: **Validation of a Japanese version of the Stanford Health Assessment Questionnaire in 3,763 patients with rheumatoid arthritis.** *Arthritis Rheum* 2003, **49**:784-788.

## A Study of 225 Malayan Pit Viper Bites in Thailand

**Guarantor:** Nualnong Wongtongkam, RN MSc

**Contributors:** Nualnong Wongtongkam, RN MSc\*; Henry Wilde, MD FACP+‡; Chitr Sitthi-Amorn, MD PhD‡; Kavi Ratanabanangkoon, PhD§

This study evaluated factors affecting the severity of bite site necrosis and systemic symptoms resulting from envenomation among patients bitten by Malayan pit vipers (*Calloselasma rhodostoma*) in Thailand. We studied 145 victims prospectively. An additional 80 medical records were obtained for a retrospective study. Collected data included gender of the victims, anatomic locations of bites, where attacks took place, and predisposing factors and how they might have affected the clinical course. Most patients presented with minimal to moderate symptoms. Only eight patients required surgical wound debridement. None required amputation. However, 27 subjects developed permanently swollen limbs, presumably because of lymphatic or vascular injury. Significant coagulopathies were common (52.48%), and the only two deaths were attributable to intracranial hemorrhage. The improved clinical outcomes in the prospective and retrospective groups, compared with older series, might have been attributable to better public education, improved road infrastructure, and more health care facilities. Less reliance on nonprofessional healers and fewer applications of tourniquets also might have influenced clinical outcomes. Malayan pit viper antivenin, manufactured in Thailand, appears to be effective in reversing dangerous coagulopathies.

### Introduction

Snake envenomation from Malayan pit viper (MPV) (*Calloselasma rhodostoma*, formerly known as *Angkistrodon rhodostoma*) (Fig. 1) bites can result in serious disability and death. Previous studies reported mortality rates of 1 to 2%.<sup>1,2</sup> Clinical manifestations of MPV bites may be local, systemic, or both. Hemorrhagic effects attributable to coagulopathy are often seen with MPV bites. The MPV is a dangerous and abundant species in southeast Asia and is known to bite without warning. It has been reported in Thailand, Cambodia, Java, Peninsular Malaysia, Myanmar, Sumatra, and Vietnam. Snakes of the Crotalinae subfamily, which includes the MPV, are responsible for the majority of reported snakebites in Thailand.<sup>3-5</sup> Clinical manifestations resulting from coagulopathy include petechiae, epistaxis, hematuria, hemoptysis, uterine, gastrointestinal, and central nervous system hemorrhage, disseminated intravascular coagulation, and shock. In MPV venom, as with other viperid venoms, hemorrhagic metalloproteinases are responsible for local hemorrhage, as a result of degradation of collagen of the vascular basement membrane, followed by total disintegration

of the vascular structure.<sup>6,7</sup> The major hemorrhagin of this venom is a glycoprotein termed rhodostoxin. It is reported to be responsible for local necrosis through activation of tissue tumor necrosis factor- $\alpha$ .<sup>8</sup> However, the role of hemorrhagic metalloproteinases in systemic complications remains unclear. The MPV venom contains proteins that affect platelet aggregation. Aggretin is a heterodimeric C-type lectin that activates platelets by binding to the platelet glycoproteins GPIb and  $\alpha_2\beta_1$ .<sup>9</sup> It may induce thrombocytopenia in envenomed patients.<sup>10</sup> A protein with similar structure, called rhodocetin, was shown to inhibit collagen-induced platelet aggregation.<sup>11</sup> The venom also contains coagulation factor II and factor X activators and thrombin-like enzymes (ancrod), which are responsible for consumption coagulopathy and organ failure. Although consumption coagulopathy and thrombocytopenia are considered to be the major causes of systemic bleeding, hemorrhagic metalloproteinases could damage vascular endothelial cells, destroying vascular integrity, provoking platelet aggregation, and activating the coagulation cascade. This may terminate in disseminated intravascular coagulation. Almost all of the systemic effects of MPV venom can be successfully treated with supportive measures and species-specific antivenin. However, local toxicity from venoms, such as edema and myonecrosis, are not prevented or reversed by antivenin therapy. The MPV venom first causes pain, swelling, and ecchymoses, which appear within minutes after the bite (Fig. 2). The MPV, unlike cobras, has large fangs that can inject venom deep into tissues. Necrosis may therefore involve the skin, subcutaneous tissues, and muscle, which can progress slowly to end in dry gangrene. These effects prolong hospitalization and increase morbidity, which may result in loss of a digit or limb.<sup>12</sup> Factors contributing to clinical outcomes are related to the snake's size and the amount of venom injected. The geographic origin of the snake and the quality of medical care rendered may also influence outcomes. It is important, at the time of presentation, to make a rapid estimate of the degree of envenomation and to determine whether there is progression of symptoms. A recently published clinical tool is the snakebite severity score (SSS), shown in Table I. The SSS helps workers assess vital signs serially, search for evidence of progression of local tenderness, edema, and induration, and prepare a flow chart of systemic symptoms. It also allows scoring of the clinical severity and improves record keeping.<sup>13</sup> This study evaluated various factors that had effects on the severity of tissue necrosis and the systemic envenomation syndromes of patients bitten by a MPV.

### Methods

One hundred forty-five patients were collected prospectively, between April 2002 and June 2003, from 10 provincial hospitals throughout the country and from districts known to have many snakebite victims. Eighty hospital charts for snakebite patients

\*The College of Public Health, Chulalongkorn University, Bangkok, Thailand.

†Queen Saovabha Memorial Institute, Thai Red Cross Society, Bangkok, Thailand.

‡Department of Medicine, Faculty of Medicine, Chulalongkorn University Hospital, Bangkok, Thailand.

§Department of Microbiology, Faculty of Science, Mahidol University, and Laboratory of Immunology, Chulabhorn Research Institute, Bangkok, Thailand.

This manuscript was received for review in January 2004. The revised manuscript was accepted for publication in April 2004.

Reprint & Copyright © by Association of Military Surgeons of U.S., 2005.




Fig. 1. MPV (*C. rhodostoma*).



Fig. 2. Severe necrotic reaction for a Thai child bitten by a MPV. The necrosis involves deep and superficial tissues.

in Prachuap Khiri Khan province from the year 2001 were also reviewed retrospectively. Eligibility criteria for patient selection included a bite from a MPV for which the species of the snake was confirmed. Patients were included if the snake was not captured but the patient presented with a good description and clinical and laboratory features consistent with MPV envenomation. These features included a venous clotting time (VCT) of >30 minutes and multiple hemorrhagic blisters at the bite site that evolved rapidly.<sup>14,15</sup> Informed consent to participate in the prospective study was obtained from patients or their relatives.

Data from hospital records in the prospective and retrospective series were summarized by two nurses and reviewed by the principal investigator. The SSS was modified from the report by Dart et al.<sup>13</sup> as indicated in Table I. The following information was tabulated and analyzed: clinical signs and symptoms, pulse, temperature and blood pressure, urinalysis results, com-

plete blood counts, and VCT. Electrolyte levels and renal function parameters were evaluated where indicated. Data were collected at admission, 2 and 12 hours later, and daily thereafter. This information was compiled with the SSS level.<sup>13</sup> Statistical analyses used the  $\chi^2$  test or Student's *t* test.

## Results

The prospective study collected a total of 145 MPV bite victims from six provinces (Fig. 3). The victims came from the southern region, which included Trang (33.79%), Nakhon Si Thammarat (22.07%), Prachuap Khiri Khan (17.93%), Surat Thani (15.06%), and Songkhla (7.50%) provinces. Surprisingly, we found that Lampang, a northern Thai province, provided 2.76% of the cases (4 of 145 cases; Fig. 4). We found no MPV bite victims in Nakhon Ratchasima (northeast), Lop Buri (central), Ratchaburi (western), and Nakhon Sawan (northern) provinces. The peak snakebite season was in May, early in the monsoon season (19.31%; Fig. 5). All snakebite victims were Thai residents. There were no foreigners or tourists in this group.

Snakebites occurred throughout the day (from 8:00 a.m. to 4:00 p.m.), representing the time that victims worked in fields or rubber plantations, but some bites occurred at night (21.25%) and then mostly near the victims' homes. The size of snakes, as reflected by the distance between fang marks, was recorded only in the prospective study. Sizes ranged between 1.6 and 1.70 cm. We searched the literature in an effort to determine the relationship of the space between fang marks and snake size but found no published study. Local Thai herpetologists assured the authors that more space between fangs is related to larger snake size (L. Chanhom, personal communication). The number of snakes brought to the hospital for identification was higher in the prospective study (Table II).

More male patients experienced snakebites (age range, 27–50 years). However, there were five children (3.5%) < 15 years of age in the prospective group and eight (10%) in the retrospective group. Most bites occurred in rural areas, outdoors, and in dark or dusky places. Bites on the lower limbs, especially legs, represented 40 to 60% of cases. Bites on fingers or toes represented 30% (Table II).

The mean time between snakebite and arrival at a hospital was 175 minutes in the prospective study, and times did not differ significantly between patients who did or did not have severe tissue necrosis. The time was significantly longer (110–690 minutes) in the retrospective study. Most patients (60% in the prospective study and 100% in the retrospective study) had not applied tourniquets. The volume of antivenin administered was four to nine vials (i.e., 40–90 mL). Approximately 95% of the MPV antivenin used came from the Queen Saovabha Memorial Institute of the Thai Red Cross Society (Bangkok, Thailand), which is now the only antivenin manufacturer in Thailand.

The incidence of tissue necrosis at the MPV bite site was 95% in the prospective study and 94% in the retrospective study. The most common SSS levels were minimal (score of 1 for 78.6% and 86.25% of the prospective and retrospective groups, respectively) (Fig. 6). These patients had local pain and mildly inflamed wounds. They required only wound care to prevent or control infection. Patients with severe local necrosis (scores of 2–3) represented 23 (15.9%) of 145 prospectively studied patients. Six (26%) of 23 patients had moderate to severe local tissue

TABLE I  
MODIFIED SSS

Criteria	Level
Pulmonary system	
No symptom/sign	0
Minimal dyspnea, chest tightness, mild or vague systemic discomfort, or respirations of 20–25 breaths/minute	1
Moderate dyspnea (tachypnea, 26–40 breaths/minute; use of accessory muscle)	2
Cyanosis, air hunger, extreme tachypnea, or respiratory insufficiency/failure	3
Cardiovascular system	
No symptom/sign	0
Tachycardia (100–125 beats/minute), palpitations, generalized weakness, benign dysrhythmia, or hypertension	1
Tachycardia (126–175 beats/minute) or hypotension, with systolic blood pressure of <100 mm Hg	2
Extreme tachycardia (>175 beats/minute) or hypotension, with systolic blood pressure of <100 mm Hg, malignant dysrhythmia, or cardiac arrest	3
Local wound	
No symptom/sign	0
Pain, swelling, or ecchymosis within 5–7.5 cm of bite site	1
Pain, swelling, or ecchymosis involving less than one-half of the extremity (7.5 cm from bite site)	2
Pain, swelling, or ecchymosis extending beyond affected extremity (>100 cm from bite site)	3
Gastrointestinal system	
No symptom/sign	0
Abdominal pain, tenesmus, or nausea	1
Vomiting or diarrhea	2
Repeated vomiting, diarrhea, hematemesis, or hematochezia	3
Hematologic system	
No symptom/sign	0
Coagulation parameters slightly abnormal: PT, <20 seconds; PTT, <50 seconds; platelets, 100,000–150,000/mL; fibrinogen, 100–150 µg/mL; VCT, 10–20 minutes	1
Coagulation parameters abnormal: PT, <20 to 50 seconds; PTT, <50 to 75 seconds; platelets, 50,000–100,000/mL; fibrinogen, 50–100 µg/mL; VCT, 21–30 minutes	2
Coagulation parameters abnormal: PT, <50 to 100 seconds; PTT, <75 to 100 seconds; platelets, 20,000–50,000/mL; fibrinogen, <50 µg/mL; VCT, >30 minutes	3
Coagulation parameters markedly abnormal, with serious bleeding or the threat of spontaneous bleeding; unmeasurable PT or PTT; platelets, <20,000/mL; undetectable fibrinogen, and severe abnormalities of other laboratory values, including VCT, also fall into this category	4
Central nervous system	
No symptom/sign	0
Minimal apprehension, headache, weakness, dizziness, chills, or paresthesias	1
Moderate apprehension, headache, weakness, dizziness, chills, paresthesia, confusion, or fasciculation in area of bite site	2
Severe confusion, lethargy, seizures, coma, psychosis, or generalized fasciculation	3

PT, prothrombin time; PTT, partial thromboplastin time.

necrosis (scores of 2–3) that required surgical debridement. There was no need for amputation in either group. The severity of tissue necrosis was significantly influenced by the quality and timing of treatment rendered ( $p < 0.05$ ). However, two victims required surgical decompression because of swelling and circulatory impairment; bite sites were a finger or toe, locations without abundant soft tissue space, causing the compartment syndrome (scores of 2–3).

Systemic manifestations involved the central nervous, cardiovascular, pulmonary, gastrointestinal, and hematologic systems. An overall analysis of SSS values revealed 0.80 to 1.00 involving the central nervous system, 0.40 to 0.90 the pulmonary system, 0.03 to 0.17 the gastrointestinal system, 0.45 to 0.83 the cardiovascular system, 1.69 to 1.84 the hematologic system, and 1.01 to 1.17 bite site reactions. The highest scores appeared 12 hours after hospitalization (Fig. 7). No patients developed septicemia but two subjects had the disseminated intravascular coagulation syndrome and died as a result of intracranial hemorrhage. The first patient, a 60-year-old man,

came to the hospital comatose and with severe coagulopathy (VCT of >30 minutes and hematuria). He had been bitten by a MPV 3 days previously and had been treated by a traditional healer with herbal remedies and local potions. He was moribund and had moderately severe tissue necrosis (score of 2) at the bite site. The patient was intubated and given 30 mL of antivenin, and his VCT returned to normal after 6 hours. However, he had developed intracranial hemorrhage and never regained consciousness. The second fatality was a 72-year-old man. He was admitted to the hospital 1 hour after having been bitten by a MPV. On the first day of admission, he had pain at the bite site and was very apprehensive but had no abnormal systemic symptoms and no coagulopathy (VCT of 10 minutes). On the morning of his second hospital day, the patient developed bleeding from his gums and had a VCT of >30 minutes. There was no antivenin available at that time. In the afternoon, the patient went into shock, lost consciousness, and developed hematuria, hematemesis, and thrombocytopenia. He remained deeply unconscious and required vasopressors. During the afternoon of



Fig. 3. Map of Thailand, showing the 10 provincial locations.




Fig. 4. Distribution of snakebite victims among the 10 provincial hospitals, in the prospective study.

the second day, after it was decided that his case was hopeless, the patient was taken home to die.

Among victims with coagulopathy, 52.48% and 35.44% in the prospective and retrospective groups, respectively, had severe abnormal coagulopathy (VCT of >30 minutes). The mean VCT was >21 minutes on the first day and gradually decreased to normal by day 5 after treatment with antivenin. Most patients who had VCTs of >30 minutes and who received antivenin (three to five vials) showed improvement in the VCT 6 to 12 hours later, with return to normal 2 days later. In the prospective study, 25 of 145 victims did receive antivenin but their VCTs




Fig. 5. Seasonal frequency of snakebites by *C. rhodostoma* during April 2002 to June 2003.

were still >30 minutes 12 hours later. The patients were given an additional three to five vials, which rapidly normalized the VCT in all cases.

## Discussion

Our prospective and retrospective studies revealed some degree of bite site tissue injury for almost all victims. None of the patients required amputations, which indicates that even severe tissue necrosis of level 3 (by SSS) usually resolves in 2 to 3 weeks. Resolution can occasionally take 2 to 3 months, and for some victims the limb may remain permanently swollen because of vascular and/or lymphatic damage.<sup>16</sup>

Species-specific antivenin neutralizes circulating venom and reverses systemic symptoms. Local bite site necrosis, however, responds poorly (if at all) to antivenin administration.<sup>17</sup> This is partly attributable to the fact that tissue injury occurs rapidly and may be well established by the time the patient arrives at a hospital. Severe tissue necrosis was found in 2 (1.4%) of 145 cases in our prospective study and required surgical debridement. Both subjects appeared with a delay of 4 to 5 hours. They did receive 15 vials (150 mL) of antivenin after arrival at the hospital, but there was still progression of local necrosis. An additional four patients with tissue necrosis, who did not receive antivenin, had first been treated by traditional healers. The patients required surgical wound debridement. There were no cases of severe tissue necrosis in the retrospectively studied group. This finding is surprising and needs an explanation. Prachuap Khiri Khan provincial hospital, the site of the retrospective study, is located in a relatively prosperous region with good road infrastructure, a higher level of education, and good access to medical centers. It is likely that patients in this region reported more rapidly for medical care, rather than visiting traditional healers. This may be one explanation for less tissue injury and better outcomes. Only 61 (25%) patients in this group received antivenin.

Traditionally, patients bitten by venomous snakes in Thailand applied tourniquets to the bite site in an attempt to delay absorption of venom into the circulation. This is no longer recommended, because it was found to be dangerous.<sup>12</sup> A previous study of patients bitten by MPVs in Thailand showed that tourniquets applied by patients failed to inhibit the spread of venom into the general circulation.<sup>12,18,19</sup> Another study reported gangrene after tourniquet application.<sup>14</sup> Data concerning bite vic-

TABLE II  
VARIOUS FACTORS AFFECTING TISSUE NECROSIS

Variables	Prospective Study		Retrospective Study	
	No Tissue Necrosis	Tissue Necrosis	No Tissue Necrosis	Tissue Necrosis
Gender (no.)				
Male	64	9	44	6
Female	58	14	30	0
Location (no.)				
Urban	22	3	25	1
Rural	100	20	49	5
Place of bite (no.)				
Indoor	12	2	1	0
Outdoor	110	21	71	6
Predisposing factors (no.)				
Barefoot	92	19	23	1
Dusk	30	4	48	5
Site of bite (no.)				
Lower	85	16	61	3
Upper	37	7	13	3
First aid treatment (no.)				
Tourniquet	47	10	0	0
No tourniquet	75	13	68	5
Treatment (no.)				
Dressing	120	17	74	5
Debridement	2	6	0	1
First seen by traditional healers (no.)	7	6		
Age (years) <sup>a</sup>	48.61 ± 1.71	46.60 ± 4.26	39.95 ± 2.21	26.85 ± 8.05
Time of bite <sup>a</sup>	00:50 p.m. ± 30 minutes	2:18 p.m. ± 59 minutes	1:54 p.m. ± 45 minutes	12:50 p.m. ± 60 minutes
Duration of hospitalization (days) <sup>a</sup>	3.02 ± 1.73	4.35 ± 0.65 <sup>b</sup>	2.15 ± 0.13	4.33 ± 1.11
Time between bite and seeking medical advice (minutes) <sup>a</sup>	175.20 ± 44.16	187.17 ± 67.04	111.35 ± 26.20	690.00 ± 580.29
VCT (minutes) <sup>a</sup>				
First day	>21.58 ± 0.84	>25.91 ± 1.97 <sup>b</sup>	>19.97 ± 0.90	>24.00 ± 3.82
Third day	>11.26 ± 0.97	>14.71 ± 0.87 <sup>b</sup>	>18.00 ± 4.24	>20.93 ± 1.51

<sup>a</sup>Mean ± SE.

<sup>b</sup>Statistically significant,  $p < 0.05$ .




Fig. 6. Incidence of tissue necrosis after *C. rhodostoma* bites in the prospective and retrospective studies.

tims, circumstances, and sites of bites, as well as anatomic locations, were similar to those reported by Mitrakul,<sup>20</sup> Ismail and Memish,<sup>21</sup> and Da Silva et al.<sup>20-22</sup> The distance between fang marks is thought to be related to the size of the snake and possibly the amount of venom injected. Where this was recorded, bites had a mean space of 1.6 to 1.7 cm between fang marks. This may indicate that they came from female adult or young male snakes (L. Chanhome, personal communication).

When the SSS values involving the central nervous, cardiovascular, pulmonary, gastrointestinal, and hematologic systems were analyzed, it appeared that the score usually was severe only on the first day and there was rapid subsequent improvement. One study<sup>14</sup> noted nonclotting blood 1.5 to 72 hours (mean, 27.5 hours) after admission. Coagulopathies appeared within the first 24 hours in our study as well. Abnormal VCTs were significantly more common among patients who also had severe bite site tissue necrosis by the first and third days of hospitalization. Recurrences of coagulopathy after an initial response to antivenin were not uncommon and occurred more frequently among subjects with initial severe clotting abnormalities. Previous studies suggested that such recurrences of coagulopathy are related to decreases in circulating antivenin levels and/or reversible binding of antivenin to venom protein. There may also be a depot of non-neutralized venom at the bite site that is released slowly.<sup>12,23,24</sup>

Most patients presenting with MPV bites remained in stable condition after admission and received no antivenin. They were discharged after a period of observation in the emergency ward (30-40%). The mean duration of hospitalization for all patients was 3 to 5 days. All patients in this study who had a VCT of >30 minutes at admission received MPV antivenin. The average dose




Fig. 7. Severity of envenomation, as evaluated with the modified SSS.

was five vials, which is in accordance with the manufacturer's recommendations. We identified only two patients who received antivenin and developed mild to moderate serum sickness (1.8%). There were no cases of anaphylaxis among a total of 111 subjects who received purified, pepsin-digested antivenin of equine origin, demonstrating again the relative safety of purified, pepsin-digested, snake antivenin.

The severity of tissue necrosis in our two groups was decreased compared with previous studies reported more than a decade ago. Those studies found significant necrosis requiring surgical debridement and even amputation in 10.9% of cases (5 of 46 victims).<sup>14</sup> We identified only 8 cases (5.5%) among 145 patients in the prospective group and one case (1.25%) among 80 patients in

the retrospective study. There were no amputations among our patients, compared with 2.2% reported by Warrell et al.<sup>14</sup> However, 27 (18.6%) of 145 subjects had permanently swollen limbs because of vascular and/or lymphatic damage. The improved outcomes in our two groups are most likely attributable to better public education and road infrastructure and expanded health care facilities in rural areas. There was also considerable improvement in the quality of the antivenin used. Warrell et al.<sup>14</sup> demonstrated that at least one batch of *C. rhodostoma* antivenin, manufactured in Thailand before 1986, was of poor efficacy. Traditional healers, who may cause delays in treatment and increase the risk of wound infection, appear to play a lesser role in caring for snakebite victims today.

The MPV (*C. rhodostoma*) is prevalent through most of south-east Asia. It is a dangerous snake, because its coloration makes it difficult to spot in its natural environment. It often bites without warning and is an occupational hazard for rubber plantation workers and the military operating in the field. This snake is both tissue toxic and hematotoxic, and deaths resulting from coagulopathy and disseminated intravascular consumption have been reported. Purified, pepsin-digested, species-specific antivenin is manufactured in Thailand. It is active against coagulopathy and systemic symptoms but appears to do little to counteract local tissue necrosis, which is usually well established by the time victims present for treatment.

### Acknowledgments

This study would not have been possible without the help of the directors and staffs of the 10 provincial hospitals. Photographs and critical review were provided by Dr. Lawan Chanhom from the Thai Red Cross Snake Farm. This study was supported by grants from the Thailand Research Fund and is being submitted to the School of Graduate Studies of Chulalongkorn University, as part of the requirements for a doctoral degree, by the senior author. None of the authors has a financial interest in the production of antivenin.

### References

1. Reid HA, Thean PC, Chan KE, Baharom AR: Clinical effects of bites by Malayan viper [*Ancistrodon rhodostoma*]. *Lancet* 1963; 1: 617-21.
2. Looareesuwan S, Viravan C, Warrell DA: Factors contributing to fatal snake bite in the rural tropics: analysis of 46 cases in Thailand. *Trans R Soc Trop Med Hyg* 1988; 82: 930-4.
3. Chanhom L, Cox MJ, Wilde H, Jintakoon P, Chaiyabutr N, Sitprija V: Venomous snakebite in Thailand: part I: medically important snakes. *Milit Med* 1998; 163: 310-7.
4. Ponchanugool C, Wilde H, Bhangmanada K, et al: Venomous snakebite in Thailand: part II: clinical experience. *Milit Med* 1998; 163: 318-23.
5. Office of the Permanent Secretary for Public Health: Division of Epidemiology. Annual Summaries (1993-1997). Bangkok, Thailand.
6. Baramova EB, Shannon JD, Bjarnason JB, Fox FW: Degradation of extracellular matrix proteins by hemorrhagic metalloproteinases. *Arch Biochem Biophys* 1989; 275: 63-71.
7. Maruyama M, Sugiki M, Yoshida E, Shimaya K, Mihara H: Broad substrate specificity of snake venom fibrinolytic enzyme: possible role in haemorrhage. *Toxicon* 1992; 30: 1387-97.
8. Moura da-Silva AM, Laing GD, Paine MJ, et al: Processing of pro-tumor necrosis factor- $\alpha$  by venom metalloproteinases: a hypothesis explaining local tissue damage following snake bite. *Eur J Immunol* 1996; 26: 2000-5.
9. Navdaev A, Clemetson JM, Polgar J, et al: Aggretin, a heterodimeric C-type lectin from *Calloselasma rhodostoma* (Malayan pit viper), stimulates platelets by binding to  $\alpha_v\beta_3$  integrin and glycoprotein Ib, activating Syk and phospholipase  $C\gamma_2$ , but does not involve the glycoprotein VI/Fc receptor  $\gamma$  chain collagen receptor. *J Biol Chem* 2001; 276: 20882-9.
10. Sanders WE, Read MS, Reddick RL, Garriss JB, Brinkhous KM: Thrombotic thrombocytopenia with von Willebrand factor deficiency induced by botrocetin: an animal model. *Lab Invest* 1988; 59: 443-52.
11. Wang R, Kini RM, Chung MCM: Rhodocetin, a novel platelet aggregation inhibitor from the venom of *Calloselasma rhodostoma* (Malayan pit viper): synergistic and noncovalent interaction between its subunits. *Biochemistry* 1999; 38: 7584-93.
12. Ho M, Warrell DA, Looareesuwan S, et al: Clinical significance of venom antigen levels in patients envenomed by the Malayan pit viper (*Calloselasma rhodostoma*). *Am J Trop Med Hyg* 1986; 35: 579-87.
13. Dart RC, Hurlbut KM, Garcia R, Boren JB: Validation of a severity score for the assessment of crotalid snakebite. *Ann Emerg Med* 1996; 27: 321-6.
14. Warrell DA, Looareesuwan S, Theakston RD: Randomized comparative trial of three monospecific antivenoms for bites by the Malayan pit viper (*Calloselasma rhodostoma*) in southern Thailand: clinical and laboratory correlations. *Am J Trop Med Hyg* 1986; 35: 1235-47.
15. Brown L, Brown AE: The problem of snakebite diagnosis: a case report and two-year hospital review. *J Med Assoc Thai* 1987; 71: 456-60.
16. Reid HA, Theakston RDG: The management of snake bite. *Bull World Health Organ* 1983; 61: 885-95.
17. Gutiérrez JM, Chaves F, Bolanos R, et al: Neutrilización de los efectos locales del veneno de *Bothrops asper* por un antiveneno polivalente. *Toxicon* 1981; 19: 493-500.
18. Tun P, Tin N-S, Myint L, Warrell DA, Than W: The efficacy of tourniquets as a first-aid measure for Russell's viper bites in Burma. *Trans R Soc Trop Med Hyg* 1987; 81: 403-5.
19. Khin OL, Aye AM, Tun P, Thainge N, Min N: Russell's viper venom levels in serum of snake bite victims in Burma. *Trans R Soc Trop Med Hyg* 1984; 78: 165-8.
20. Mitrakul C: Clinical features of viper bites in 72 Thai children. *Southeast Asian J Trop Med Public Health* 1982; 13: 628-36.
21. Ismail M, Memish ZA: Venomous snakes of Saudi Arabia and the Middle East: a keynote for travellers. *Int J Antimicrob Agents* 2003; 21: 164-9.
22. Da Silva CJ, Jorge MT, Ribeiro LA: Epidemiology of snakebite in a central region of Brazil. *Toxicon* 2003; 41: 251-3.
23. Smith TW, Lloyd BL, Spicer N, Haber E: Immunogenicity and kinetics of distribution and elimination of sheep digoxin-specific IgG and Fab fragments in the rabbit and baboon. *Clin Exp Immunol* 1979; 36: 384-96.
24. Meyer WP, Habib AG, Onyade AA, et al: First clinical experiences with a new ovine Fab *Echis ocellatus* snake bite antivenom in Nigeria: randomized comparative trial with Institute Pasteur Serum (Ipsen) Africa antivenom. *Am J Trop Med Hyg* 1997; 56: 291-300.



Dr. Perlin presents the John D. Chase Award for Physician Executive Excellence to Dr. Michael J. Kussman-VA (BG, MC, USA (Ret.))

## A Study of Thai Cobra (*Naja kaouthia*) Bites in Thailand

**Guarantor:** Nualnong Wongtongkam, RN MSc

**Contributors:** Nualnong Wongtongkam, RN MSc\*; Henry Wilde, MD FACP†‡; Chitr Sitthi-Amorn, MD PhD‡; Kavi Ratanabanangkoon, PhD§

This study evaluated factors affecting the severity of bite site necrosis and systemic symptoms resulting from envenomation among patients bitten by Thai cobras (*Naja kaouthia*) in Thailand. We studied 45 victims prospectively. An additional 40 medical records were obtained for a retrospective study. Collected data included gender of the victims, anatomic locations of bites, where attacks took place, and predisposing factors and how they might have affected the clinical course. Most patients were asymptomatic or mildly symptomatic. Neurotoxic symptoms and respiratory failure developed in 31.11% and 12.50% in the prospective and retrospective groups, respectively. Only one patient died, from the effects of prolonged respiratory failure. There was some degree of tissue necrosis at the bite site for almost all victims. One victim required amputation of a digit in the retrospective study, and 33.60% of the prospective group and 20% of the retrospective group required minor surgical debridement. Snakebites in Thailand are still a public health problem, although rapid urbanization has decreased the number of victims because of degradation of the snake's habitat.

### Introduction

The Thai cobra (*Naja kaouthia*) (Fig. 1) is the most well-known venomous snake in Thailand. It is widely distributed throughout the country and is also found in northeastern India, the Andaman Islands, Nepal, Bangladesh, Myanmar, Malaysia, Laos, Cambodia, southern Vietnam, and southern China.<sup>1</sup> It is not the only *Naja* species found in Thailand. *Naja sumatrana*, also known as the golden spitting cobra, is found in southern Thailand and *Naja siamensis* is the most common spitting cobra seen in the western and central regions, especially in the provinces of Ang Thong, Suphan Buri, Kanchanaburi, and Tak. The king cobra (*Ophiophagus hannah*) is found in more remote regions of Thailand. It belongs to a different genus than *Naja*, but there are great similarities in the action of the venom. The lethal toxins of the genera *Naja* and *Bungarus* show a high degree of amino acid sequence homology.<sup>2</sup> The bite of *N. kaouthia* is dangerous, with a mortality rate that appears to be higher than that for other venomous snakes.<sup>3</sup> The main lethal component of the venom is a postsynaptic neurotoxin that binds to the nicotinic cholinergic receptor sites at the neuromuscular junction. It produces an effect similar to that seen with curare poisoning.<sup>4</sup> The neurotoxicity results in ptosis, ophthalmoplegia with blurred vision or diplopia, dysphagia with increased salivation, dysar-

thria, flaccid paralysis, loss of deep tendon reflexes, coma, and respiratory failure.<sup>5</sup> In addition to the neurotoxicity, local tissue swelling and inflammation, followed by tissue necrosis, are common manifestations<sup>6-8</sup> (Fig. 2). Cardiotoxin (also known as cytotoxin), direct lytic factors, and myotoxin are present in cobra venom and are almost certainly responsible for myonecrosis. However, tissue necrosis usually involves only skin and subcutaneous tissue (unlike with bites from most vipers, which have larger fangs) (Fig. 2). Once tissue damage has occurred, antivenin of high potency and in large quantity fails to prevent further tissue necrosis. Bite site injury prolongs hospitalization and may increase morbidity, particularly if complicated by secondary infection.<sup>7,9</sup> The severity of local tissue necrosis and respiratory failure are related to the timing, dose, and potency of venom injected and the quality of supportive care rendered.

It is important, at the time of presentation, to make a rapid estimate of the degree of envenomation and to determine whether there is progression of symptoms (i.e., worsening of clinical signs and symptoms over time). A recently published, useful, clinical tool is the snakebite severity score (SSS), shown in Table 1.<sup>10</sup> It helps workers assess vital signs serially, search for evidence of progression of local tenderness, edema, and induration, and prepare a flow chart of systemic symptoms. The SSS thus provides a method for scoring clinical severity and improves record keeping.

Respiratory failure and local tissue necrosis have been well recognized as major causes of death and morbidity. Therefore, this study evaluated various factors that had effects on the severity of tissue necrosis and on the systemic envenomation syndrome of patients bitten by Thai cobras.

### Methods

Forty-five patients were collected prospectively, between April 2000 and June 2003, from 10 provincial hospitals throughout the country that were known to treat a significant number of snakebite victims. Forty hospital charts of snakebite patients in Nakhon Sawan, a province north of Bangkok, from the years 1997-2000 were also reviewed retrospectively.

Eligibility criteria for patient selection included a bite from *Naja kaouthia* for which the species of the snake was confirmed. Patients were also included if the snake was not captured but the patient presented with a good description and clinical features of *N. kaouthia* envenomation, such as ptosis or dysphagia with increased salivation.<sup>5</sup> Informed consent to participate in the study was obtained from patients or their relatives.

Data from hospital records in the prospective and retrospective series were summarized by two nurses and reviewed by the principle investigator. The SSS, as modified from the report by Dart et al.,<sup>10</sup> was used to compile data (Table I). The following information was tabulated and analyzed: clinical signs and symptoms, pulse rate, and temperature. Data were collected at

\*The College of Public Health, Chulalongkorn University, Bangkok, Thailand.

†Queen Saovabha Memorial Institute, Thai Red Cross Society, Bangkok, Thailand.

‡Department of Medicine, Faculty of Medicine, Chulalongkorn University Hospital, Bangkok, Thailand.

§Department of Microbiology, Faculty of Science, Mahidol University, and Laboratory of Immunology, Chulabhorn Research Institute, Bangkok, Thailand.

This manuscript was received for review in December 2003. The revised manuscript was accepted for publication in April 2004.

Reprint & Copyright © by Association of Military Surgeons of U.S., 2005.



Fig. 1. Defensive position of *N. kaouthia*.

admission, 2 and 12 hours later, and daily thereafter. Statistical analyses used the  $\chi^2$  test or Student's *t* test.

### Results

The prospective study collected a total of 45 *N. kaouthia* snakebite victims from 10 provinces (Fig. 3). Most victims came from Nakhon Si Thammarat (southern) and Nakhon Sawan



Fig. 2. Cobra bite 1 week later. The tissue necrosis is superficial and does not involve muscle.

(northern) provinces, i.e., 13 (28.89%) and 11 (24.44%), respectively (Fig. 4). The peak prevalence was in May (13 of 45 cases; 28.89%), which is in the early part of the rainy season (Fig. 5). Snakebites by *N. kaouthia* were distributed throughout the

TABLE I  
MODIFIED SSS

Criteria	Level
Pulmonary system	
No symptom/sign	0
Dyspnea, minimal chest tightness, mild or vague discomfort, or respirations of 20-25 breaths/minute	1
Moderate respiratory distress (tachypnea, 26-40 breaths/minute; accessory muscle use)	2
Cyanosis, air hunger, extreme tachypnea, or respiratory insufficiency/failure	3
Cardiovascular system	
No symptom/sign	0
Tachycardia (100-125 beats/minute), palpitations, generalized weakness, benign dysrhythmia, or hypertension	1
Tachycardia (126-175 beats/minute) or hypotension, with systolic blood pressure of <100 mm Hg	2
Extreme tachycardia (>175 beats/minute) or hypotension, with systolic blood pressure of <100 mm Hg, malignant dysrhythmia, or cardiac arrest	3
Local wound	
No symptom/sign	0
Pain, swelling, or ecchymosis within 5-7.5 cm of bite site	1
Pain, swelling, or ecchymosis involving less than one-half the extremity (7.5 cm from bite site)	2
Pain, swelling, or ecchymosis extending beyond affected extremity (>100 cm from bite site)	3
Gastrointestinal system	
No symptom/sign	0
Pain, tenesmus, or nausea	1
Vomiting or diarrhea	2
Repeated vomiting, diarrhea, hematemesis, or hematochezia	3
Hematologic system	
No symptom/sign	0
Coagulation parameters slightly abnormal: PT, <20 seconds; PTT, <50 seconds; platelets, 100,000-150,000/mL; fibrinogen, 100-150 $\mu$ g/mL	1
Coagulation parameters abnormal: PT, <20 to 50 seconds; PTT, <50 to 75 seconds; platelets, 50,000-100,000/mL; fibrinogen, 50-100 $\mu$ g/mL	2
Coagulation parameters abnormal: PT, <50 to 100 seconds; PTT, <75 to 100 seconds; platelets, 20,000-50,000/mL; fibrinogen, <50 $\mu$ g/mL	3
Coagulation parameters markedly abnormal, with serious bleeding or the threat of spontaneous bleeding; unmeasurable PT or PTT; platelets, <20,000/mL; undetectable fibrinogen and severe abnormalities of other laboratory values, including venous clotting time, also fall into this category	4
Central nervous system	
No symptom/sign	0
Minimal apprehension, headache, weakness, dizziness, chills, or paresthesia	1
Moderate apprehension, headache, weakness, dizziness, chills, paresthesia, confusion, or fasciculation in area of bite site, ptosis, and dysphagia	2
Severe confusion, lethargy, seizure, coma, psychosis, or generalized fasciculation	3

PT, prothrombin time; PTT, partial thromboplastin time.



Fig. 3. Map of Thailand, showing the 10 provincial locations.

day and were most common between 11:00 a.m. and 4:00 p.m. (48.9% and 37.5% in the prospective and retrospective studies, respectively). Some victims were bitten at night (15%). The size of the snakes, as reflected by the distance between fang marks, was recorded only in the prospective study. Distances ranged between 1.45 and 1.50 cm. The number of snakes presented for identification was larger (27 of 45 cases, 60%) in the prospective study (Table II).

More male patients experienced snakebites, with a mean age range of 37 to 50 years. However, there were two children (4.5%) <15 years of age in the prospective study and five (12.5%) in the retrospective study. Most bites were on toes and fingers (40%). Bites of the lower limbs, especially feet, represented 25 to 35% (Table II). Most bites occurred in rural areas, outdoors, and in dusky or dark places. Times between the bite and arrival at a hospital or health center ranged between 65 and 120 minutes (1–2 hours) and did not differ significantly between patients who did or did not have severe tissue necrosis, in both groups. Twenty-seven (60%) of 45 victims applied tourniquets in the prospective study, and 7 (17.50%) of 40 victims did so in the retrospective study. Only two patients in the prospective group and none in the retrospective group were first treated by traditional healers. Antivenin therapy ranged from 1 to 20 vials (mean of 2–6 vials, 20–60 mL, in both groups). The antivenin used for all groups was from the Queen Saovabha Memorial Institute of the Thai Red Cross Society (Bangkok, Thailand). The




Fig. 4. Distribution of snakebite victims among the 10 provincial hospitals, in the prospective study.




Fig. 5. Seasonal frequency of snakebites by *N. kaouthia* during April 2002 to June 2003. The monsoon season is May to October.

range of hospitalization was 2 to 6 days, with a maximum of 29 and 20 days in the prospective and retrospective studies, respectively (Table II). The severity of tissue necrosis was related to the length of hospitalization only in the retrospective study ( $p < 0.05$ ).

Tissue necrosis at the bite site was seen for 91.11% (41 of 45 cases) and 65% (26 of 40 cases) of cases in the prospective and retrospective studies, respectively. The most common SSS level was minimal tissue necrosis (score of 1 for 84.5% and 60% in the prospective and retrospective groups, respectively). There was no case of severe tissue necrosis in either group (Fig. 6). Twenty-nine of 45 victims (64.5%) in the prospective group and 77.5% in the retrospective group required some surgical wound care to prevent or control infection, but the wounds were of moderate local severity (score of 2 for 33.6% and 20% in the prospective and retrospective groups, respectively). Only one patient required amputation of the thumb, in the retrospective group. Lack of abundant soft tissue to allow expansion with swelling (the compartment syndrome) was the cause.<sup>11</sup>

TABLE II  
VARIOUS FACTORS AFFECTING TISSUE NECROSIS

Variables	Prospective Study		Retrospective Study	
	No Tissue Necrosis	Tissue Necrosis	No Tissue Necrosis	Tissue Necrosis
Gender (no.)				
Male	4	21	11	15
Female	0	20	3	11
Location (no.)				
Urban	1	4	1	14
Rural	3	37	13	22
Place of bite (no.)				
Indoor	1	12	3	8
Outdoor	3	29	11	18
Predisposing factors (no.)				
Barefoot	3	32	3	6
Darkness	1	9	11	20
Site of bite (no.)				
Lower	2	25	6	18
Upper	2	16	8	8
First aid (no.)				
Tourniquet	3	24	1	6
No tourniquet	1	17	13	20
Treatment (no.)				
Dressing	3	26	13	18
Debridement	1	15	1	7
Amputation			0	1
Age (years) <sup>a</sup>	46.25 ± 7.77	41.92 ± 2.50	43.26 ± 4.12	37.30 ± 4.06
Time of bite <sup>a</sup>	11:25 a.m. ± 51 minutes	00:30 p.m. ± 49 minutes	4:01 p.m. ± 49 minutes	2:50 p.m. ± 42 minutes
Time between bite and seeking medical advice (minutes) <sup>a</sup>	68.75 ± 38.42	83.65 ± 11.73	120.00 ± 18.50	95.19 ± 19.09
Distance between fang marks (cm) <sup>a</sup>	1.50 ± 0.28	1.45 ± 0.10		
No. of antivenin vials used <sup>a</sup>	6.25 ± 3.42	4.37 ± 0.84	4.07 ± 2.19	2.15 ± 1.01
Duration of hospitalization (days) <sup>a</sup>	5.50 ± 3.17	5.65 ± 0.88	2.87 ± 0.51	5.34 ± 0.89 <sup>b</sup>

<sup>a</sup>Mean ± SE.

<sup>b</sup>Statistically significant,  $p < 0.05$ .




Fig. 6. Tissue necrosis after *N. kaouthia* bites in the prospective and retrospective studies.

Clinical manifestations involved the central nervous, cardiovascular, pulmonary, gastrointestinal, and hematologic systems and injury at the bite site (Fig. 7). An overall analysis of SSS values revealed 1.05 to 1.62 involving the central nervous

system, 1.07 to 1.42 the pulmonary system, 0.13 to 0.27 the gastrointestinal system, 0.55 to 0.89 the cardiovascular system, 0 the hematologic system, and 0.70 to 0.98 bite site injury. The highest scores were observed 12 hours after hospitalization; most patients presented with no or only mild symptoms, which gradually decreased to level 0 (no symptoms) after 5 days of hospitalization, in both the prospective and retrospective studies. On the first day of hospitalization, 14 (31.11%) of 45 victims in the prospective group and 5 (12.50%) of 40 victims in the retrospective group had dysphagia, flaccid paralysis, and respiratory failure. The patients were intubated. Only one patient, in the prospective study, died as a result of respiratory failure. He was 66 years of age and came to the hospital deeply comatose, in advanced respiratory failure. He was intubated, given 50 mL of antivenin for the first dose, and given the same antivenin dose 2 hours later. He never regained consciousness in >4 days of hospitalization and was taken home to die at the request of his family.

## Discussion

Our two studies revealed some degree of tissue necrosis and progression of systemic symptoms for almost all victims. The




Fig. 7. Severity of envenomation, as evaluated with the modified SSS.

incidence of tissue necrosis was 91.11% (41 of 45 cases) in the prospective group and 65% (26 of 40 cases) in the retrospective group. Most patients had minimal tissue necrosis and no patients had severe tissue necrosis. However, one victim required amputation of the thumb and almost one-third required minor surgical debridement. Patients presented to medical facilities quickly, with times between bite and admission ranging from 60 to 120 minutes. It is noteworthy that evidence of tissue necrosis

was already present at arrival. There was no difference in the severity of tissue necrosis among patients who arrived more rapidly after being bitten and those who arrived with significant delay. A previous study showed that antivenin administration within 30 minutes after the bite did not prevent local tissue damage.<sup>12</sup> Histopathologic findings suggested that local reactions at the bite sites result from vasculitis and thrombosis of superficial and deep dermal vessels and may extend to subcu-

taneous tissues.<sup>9</sup> Neurotoxic symptoms and respiratory failure developed in 31.11% and 12.5% of cases in the prospective and retrospective groups, respectively. They required intubation and assisted respiration, and one patient died as a result of respiratory failure, because of a delay in treatment and irreversible central nervous system damage.

Species-specific antivenin neutralizes circulating venom and reverses systemic symptoms. It does not immediately release venom bound to binding sites at neuromuscular junctions. Our data indicated that 9 of 10 victims who required intubation regained spontaneous respiratory function, with reversal of dysphagia, dysarthria, and flaccid paralysis, within 2 to 3 days after receiving *Naja* antivenin (mean of 11 vials, 110 mL). A previous study showed that 100 mL is adequate for recovery of respiratory paralysis and has the same beneficial effect as 200 mL.<sup>7</sup> Antivenin administered early and in adequate doses shortens the duration of paralysis and the time of respiratory assistance.<sup>13,14</sup> Pochanugool et al.<sup>7</sup> showed that, even if species-specific antivenin is administered in a timely manner, severely envenomed patients still require intubation and respiratory assistance.

Traditionally, patients bitten by venomous snakes in Thailand applied tourniquets to the bite site, in an attempt to delay absorption of venom into the systemic circulation. In our study, 27 (60%) of 45 victims used a tourniquet in the prospective study and 7 (25.92%) of 27 patients required intubation and assisted respiration. Among patients who did not apply tourniquets, 38.89% (7 of 18 patients) developed respiratory failure. Previous reports showed that tourniquets applied by patients failed to inhibit the spread of venom into the general circulation.<sup>15-17</sup> There was no relationship between the use of restriction bandages and the kinetics of serum venom levels among patients after *N. kaouthia* bites.<sup>18</sup> Our data concerning bite victims, circumstances, and sites of bites, as well as anatomic locations, were similar to those reported by others.<sup>9,18,19</sup>

Most patients who presented with *N. kaouthia* bites remained asymptomatic or mildly symptomatic. They received no antivenin (45-75%) and were discharged after periods of observation ranging from 3 to 4 days. Cases with moderate to severe symptoms (SSS of 5), with involvement of the central nervous, cardiovascular, pulmonary, gastrointestinal, and/or hematologic systems, showed that these symptoms were severe only during the first day and improved rapidly thereafter. The mean amounts of antivenin used were two to six vials (20-60 mL). The beneficial effects on neurotoxic and systemic symptoms of the administration of specific antivenin were shown by Pochanugool et al.<sup>7</sup> We identified only one patient who had a positive skin test result with equine cobra antivenin; this patient was not given antivenin and received only prolonged supportive treatment. No patients developed serum sickness (0 of 45 cases).

Hospitalization was prolonged significantly for patients with tissue necrosis. The range of hospitalization times was 20 to 29 days. Most patients were rural farm workers with a high risk of secondary infection if discharged home.

In 1988, Pongprasit et al.<sup>9</sup> showed that 51% of cobra bite victims presented with severe tissue necrosis and 27.66% required full-thickness skin grafts. More recently, Chulalongkorn University Hospital found that 60.3% of patients bitten by cobras presented with tissue necrosis.<sup>18</sup> Many (if not most) of

those patients represented a selected population referred to Chulalongkorn University Hospital from other medical facilities. Rural medical care has been significantly expanded and improved in the past decades, and fewer patients are referred to secondary and tertiary care facilities. Also, the number of snakebites throughout the country has decreased, because of reduction of the snake population as a result of snakes being hunted for their hides and flesh<sup>7</sup> and degradation of their habitat.

## Acknowledgments

This study would not have been possible without the help of the directors and staffs of the 10 provincial hospitals. Photographs and critical review were provided by Dr. Lawan Chanhome from the Thai Red Cross and Dr. Trinh Xuan Kiem from the Center of Venom Research (Ho Chi Minh City, Vietnam). This study was supported by grants from the Thailand Research Fund and is being submitted to the School of Graduate Studies of Chulalongkorn University, as part of the requirements for a doctoral degree, by the senior author. None of the authors has a financial interest in the production of antivenin.

## References

1. Cox MJ: The Snakes of Thailand and Their Husbandry. Malabar, FL, Krieger Publishing, 1991.
2. Tan NH, Saifuddin MN: Enzymatic and toxic properties of *Ophiophagus hannah* (king cobra) venom and venom fractions. *Toxicon* 1989; 27: 689-95.
3. Looareesuwan S, Viravan C, Warrell DA: Factors contributing to fatal snake bite in the rural tropics: analysis of 46 cases in Thailand. *Trans R Soc Trop Med Hyg* 1988; 82: 930-4.
4. Le Goas R, Laplante SR, Mikou A, et al:  $\alpha$ -Cobratoxin: proton NMR assignments and solution structure. *Biochemistry* 1992; 31: 4867-75.
5. Minton SA: Neurotoxic snake envenoming. *Semin Neurol* 1990; 10: 52-61.
6. Kuo TP, Wu CS: Clinicopathological studies on snakebites in Taiwan. *Snake* 1972; 4: 1-22.
7. Pochanugool C, Linthongkul S, Wilde H: Management of Thai cobra bites with a single bolus of antivenin. *Wilderness Environ Med* 1997; 8: 20-3.
8. Liao WB, Lee CW, Tsai YS, Liu BM, Chung KJ: Influential factors affecting prognosis of snakebite patients management: Kaohsiung Chang Gung Memorial hospital experience. *Chang Gung Med J* 2000; 23: 577-83.
9. Pongprasit P, Mitrakul C, Noppakun N: Histopathology and microbiological study of cobra bite wounds. *J Med Assoc Thai* 1988; 71: 475-80.
10. Dart RC, Hurlbut KM, Garcia R, Boren J: Validation of a severity score for the assessment of crotalid snakebite. *Ann Emerg Med* 1996; 27: 321-6.
11. Huang TT, Blackwell SL, Lewis SR: Hand deformities in patients with snakebite. *Plast Reconstr Surg* 1978; 62: 32-6.
12. Homma M, Tu AT: Antivenin for the treatment of local tissue damage due to envenomation by Southeast Asian snakes: ineffectiveness in the prevention of local tissue damage in mice after envenomation. *Am J Trop Med Hyg* 1970; 19: 880-4.
13. Alandikar KK: Cobra poisoning. *Indian Med Gaz* 1920; 55: 60-8.
14. Croley VSE: Notes on a case of recovery from the bite of a cobra. *Trans R Soc Trop Med Hyg* 1922; 16: 57-60.
15. Ho M, Warrell MJ, Warrell DA, Bidwell D, Voller A: A critical reappraisal of the use of enzyme-linked immunosorbent assays in the study of snake bite. *Toxicon* 1986; 35: 253-60.
16. Tin P, Tin N-S, Mynt L, Warrell DA, Than W: The efficacy of tourniquets as a first aid measure for Russell's viper bites in Burma. *Trans R Soc Trop Med Hyg* 1987; 81: 403-5.
17. Khin OL, Aye AM, Tin P, Thainge N, Min N: Russell's viper venom levels in serum of snake bite victims in Burma. *Trans R Soc Trop Med Hyg* 1984; 78: 165-8.
18. Hung DZ, Liao MY, Lin Shiao SY: The clinical significance of venom detection in patients of cobra snakebite. *Toxicon* 2003; 41: 409-15.
19. Watt G, Padre L, Tuazon ML, Theakston RDG, Laughlin LW: Tourniquet application after cobra bite: delay in the systemic absorption of neurotoxins and dangers of sudden release. *Trans R Soc Trop Med Hyg* 1988; 38: 618-22.

# SEXUAL BEHAVIORS AND OPINIONS ON SEXUALITY OF ADOLESCENTS IN A SLUM COMMUNITY IN BANGKOK

Ratana Somrongthong<sup>1</sup>, Panee Panuwatsuk<sup>2</sup>,  
David Amarathithada<sup>3</sup>, Orapin Chaipayom<sup>1</sup> and Chitr Sitthi-amorn<sup>1</sup>

<sup>1</sup>College of Public Health, Chulalongkorn University, <sup>2</sup>Department of Nursing,  
Faculty of Medicine, Ramathibodi Hospital, Mahidol University, Bangkok, Thailand;

<sup>3</sup>School of Public Health, University of Illinois at Chicago, USA

**Abstract.** This study was a survey research aiming to investigate sexual behaviors and opinions on sexuality of adolescents in a slum community. The study group comprised of 377 adolescents aged 12-22 years in a slum community in Bangkok randomly selected, and data were collected using self-administered questionnaires. Results indicated that 18.8% of the adolescents were sexually experienced with the average age of 15 years old at first intercourse. 63.1% of the adolescents had unprotected sexual intercourse with lovers or friends. Almost one-third of the population believed that premarital sexual activity was acceptable. One-sixth of the adolescents agreed that having sexual intercourse with a lover is safe, assuming that they had trustworthy partners and that having sexual intercourse was the best way to prevent their lover from having sexual activities with other partners. In addition, gender and age range were found to be the factors that significantly related to the adolescents' opinions that premarital sexual activity was acceptable and having sexual intercourse with a lover was safe ( $p < 0.05$ ), whereas the relationship between the opinions and education level was statistically insignificant. It is recommended that familial, academic, community and public health support are necessary in educating the adolescents on reproductive health and family planning in order to reduce high risk behaviors associated with acquiring HIV and other STDs.

## INTRODUCTION

Presently, adolescents are categorized as a risk group for acquiring sexually transmitted diseases (STDs) (Anonymous, 1994) due to the curious and experimental nature and as they begin to develop sexual desires (Steinberge, 1993). Sexual activity among adolescents was found to increase in developing countries (Hedberg *et al*, 1998). This was endorsed by the World Health Organization report, which found that 50% of new HIV infected patients globally were in the 10-24 year old age range (WHO, 1989). These adolescents included those who were outside education systems, those with inferior social opportunity, and youth laborers, all of whom were at greater risk for acquiring STDs, especially HIV/AIDS, than subjects in other groups (Cates and Pheeters, 1997, unpublished paper).

This study was, therefore, a survey research to investigate opinions about sexuality and sexual behaviors of adolescents with inferior social opportunities. Klong Toey community was selected for this survey because it is a slum community with the majority of its population being low-income and facing problems of poverty, employment insecurity and a lack of land ownership because the Port Authority of Thailand has rights on the land (Office of the Prime Minister, 1994). This study aimed to survey risky sexual behaviors and opinions on sexuality of adolescents in Klong Toey to reduce high-risk behaviors among disadvantaged youths.

## MATERIALS AND METHODS

The study group was randomly recruited from 14,800 teenagers in Bangkok's Klong Toey community. A study group of 375 was determined by using Taro Yamane Table. The researcher, however, conducted additional surveys giving a sample size of 377. The error of the

---

Correspondence: Ratana Somrongthong, College of Public Health, Chulalongkorn University, Bangkok 10330, Thailand.

study group was determined to be 0.05.

### Sampling method

A cluster stage random sampling technique was utilized. Randomly selected subjects from 25 communities in Klong Toey district obtaining a total of 23 communities. In sampling of community populations, systematic random sampling was adopted to determine the ratio of the populations in the communities. Sampling of household subjects was done by determining the ratio and sample size of the subjects by systematic random sampling obtaining the ratio of 1:9.

### Instruments

Questionnaires on demographical data and on sexual behaviors and opinions about sexuality of adolescents were employed in this study.

### Data analysis

The data were analyzed using the SPSS PC program by dividing the analysis as follow: frequency distribution, percentage, mean, and standard deviation were obtained in analysis of demographical data and sexual behaviors of the study groups. Chi-square test was employed in analysis of opinions on sexuality.

## RESULTS

### General characteristics of the study group

As shown in Table 1, the ratio of female to male subjects is 1.5: 1. The average age was 16 years old with 19.1 % of the subjects in the 14-17 years old range and 37.7 % of the subjects in the 18-22 years old range. The ratio of adolescents inside education systems to those outside education systems was 2.5:1.

### Sexual experiences of adolescents

The result indicates that 81.2% of the study group never had sexual experiences, while 18.8% of that had sexual experiences with the average age of 15 years old at first intercourse.

### Sexual partners and uses of condom during sexual activities

Sexual partners of the sexually active adolescents mainly comprised of their lovers (70.5%). When having sexual intercourse with their lovers, 63.1% of the subjects did not use condoms, while only 12.3% always used condoms. When having sexual intercourse with friends, 42.8% of the subjects did not wear condoms, 28.6% always used condoms. However, when having sexual intercourse with pros-

Table 1  
Demographical data.

Variables	Number	%
<b>Gender</b>		
Male	166	44.0
Female	211	56.0
<b>Age</b>		
12-13 years old (Early teenage)	72	19.1
14-17 years old (Mid teenage)	163	43.2
18-22 years old (Late teenage)	142	37.7
Mean ( $\bar{x}$ ) = 16.0	377	100.0
Standard deviation = 3.02		
<b>Education background</b>		
Studying both inside and outside conventional education systems	272	72.1
Not studying	105	27.9

Table 2  
Percentage of partners whom the adolescents had sexual activities which classified by percentages of frequency of condom uses.

Sexual partners of the adolescents (No. of subjects)	Frequency of condom uses during sexual activities		
	Always used (%)	Occasionally used (%)	Never used (%)
Lovers (50)	12.3	24.6	63.1
Friends (14)	28.6	28.6	42.8
Female/male prostitutes (7)	71.4	14.3	14.3

titutes, the majority of the subjects always used condoms, while only 14.3% of those never used condoms during sexual intercourse as shown in Table 2.

#### Opinions on sexuality of adolescents

70.6 % of the study population believed that having premarital sex was unacceptable. 84.9% of the adolescents disagreed that having sexual intercourse with a lover is safe, reasoning that they did not trust their lovers or that they did not know if their partners had STDs. Only 15.1% of the adolescents agreed with the opinion, reasoning that their partners were trustworthy or that it was the best prevention from their partners having multiple sexual partners.

#### Opinions on the risk for acquiring sexually transmitted diseases

74.8 % of the population believed that they were not at risk for STDs because they never had sexual experiences. 7.4 % believed that they were at risk for STDs because they had unprotected sexual intercourse with their lovers or friends until present time and their lovers or friends also had unprotected sexual intercourse with other partners. However, 17.8% of the subjects were unsure of such risk even though they had unprotected sexual experiences.

#### Factors relating to the opinions on sexuality of the adolescents

The factors that significantly related to the opinions of the adolescents that 'having sexual intercourse with a lover is safe' included gender

( $p=0.000$ ) and age range ( $p=0.0002$ ) (Table 3).

The factors that significantly related to the opinions of the adolescents that 'having premarital sexual activities are acceptable' included gender ( $p=0.0002$ ) and age range ( $p=0.0002$ ) (Table 4).

## DISCUSSION

The study results found that one-fourth of teenagers were sexually experienced with the average age of 15 years at first sexual intercourse. This corresponds to a study, which found that the majority of adolescents in slum communities became sexually experienced between the ages of 14-16 years of age (Kanungsukaseam, personal communication). Interestingly, adolescents who had sexual intercourse with lovers or friends did not wear condoms, believing that their partners were trustworthy and that it was the way to prevent their partners from having sexual activities with other people. Consequently, the adolescents in this slum community were at a high risk for acquiring and spreading HIV and other STDs, which corresponds with the WHO report that found that teenagers under the age of 20 years old in developing countries are a risk group for acquiring STDs, especially HIV/AIDS (Reid, 1994).

Almost one-third of the adolescents believed that premarital sexual activity is acceptable. These opinions reflect the tendency of the adolescents in the slum community to have liberal sexual attitudes. Furthermore, teenagers typi-

Table 3  
Factors relating to the opinion that having sexual intercourse with lovers is safe.

Variables	Opinion that having sexual intercourse with lovers is safe		p-value of $\chi^2$
	Agree (%)	Disagree (%)	
<b>Gender</b>			
Male	11.4	32.6	0.000
Female	3.7	52.3	
<b>Age</b>			
12-13 years old (Early teenage)	4.0	14.9	0.0002
14-17 years old (Mid teenage)	8.7	34.3	
18-22 years old (Late teenage)	2.4	35.7	

Table 4  
Factors relating to the opinion that having premarital sexual activities are acceptable.

Variables	Opinion that having premarital sexual activities are acceptable		p-value of $\chi^2$
	Agree (%)	Disagree (%)	
<b>Gender</b>			
Male	17.5	12.0	0.0002
Female	11.9	58.6	
<b>Age</b>			
12-13 years old (Early teenage)	4.0	14.8	0.0002
14-17 years old (Mid teenage)	8.8	34.5	
18-22 years old (Late teenage)	16.6	21.3	

cally imitate the behaviors of their peer group, which can increase risky behavior (Brown, 1990). Such sexual attitudes might be used to predict incidence of illness among adolescents with high-risk sexual practices. These facts emphasize that teenagers in this community are potentially at high risk for acquiring AIDS and other STDs. Furthermore in Thai societies, discussion about sexuality is taboo, especially among young women. Because of the culturally sensitive nature of sexual health, Thai adolescents face limited information and knowledge of reproductive health and health service provisions.

For these reasons, the pattern of health care service provision should be promptly improved by increasing services to support education on reproductive health, reproductive system diseases, proper sexual practices, prevention of STDs, and family planning. Health facilities and community organizations need to be coordinated in preparing and integrating such courses into school teaching to suit the actual problems of the adolescents in the slum community. Health care providers need to be trained for the specific needs of adolescents in terms of health service provisions and information on safe sex practices and family planning. Moreover, health professional should provide familial support and communication should be promoted among the parents of these adolescents to reduce the sex taboo associated with Thai culture.

#### ACKNOWLEDGEMENTS

We thank the Thailand Research Fund (TRF) and WHO Thailand for funding the study. We specially wish to acknowledge Dr EJ Love for their kindness and guidance for his project.

#### REFERENCES

- Anonymous. The HIV/AIDS pandemic: 1994. World Health Organization, 1994: 1-13.
- Brown B. Peer groups. In: Feldman S, Elliot G, eds. At the threshold: the developing adolescent. Cambridge: Harvard University Press, 1990.
- Hedberg VA, Klein JD, Andersen E. Health counseling in adolescent preventive visits: effectiveness, current practices, and quality measurement. *J Adolesc Health* 1998; 23: 344-53.
- Office of the Prime Minister. National Statistical Office. The 1994 demographic survey of population in congested community of Bangkok Metropolis. Bangkok : National Statistical Office, 1994.
- Reid E. Young women: silence, susceptibility and HIV epidemic, HIV and development program. New York: United Nation Development Program, 1994.
- Steinberge L. Adolescent. New York: McGraw-Hill, 1993.
- World Health Organization. The health of youth, fact for action: youth and AIDS. Geneva: WHO, 1989.

## **Existing health needs and related health services for adolescents in a slum community in Thailand**

Ratana Somrongthong, MA and Chitr Sitthi-amorn, MD, PhD

*College of Public Health, Chulalongkorn University,  
Bangkok, Thailand*

### **ABSTRACT**

The health problems of youth have dramatically shifted in the last 30 years from biological to social causes of morbidity and mortality. In this population, major health problems that are becoming increasingly important for preventive intervention include substance abuse, teenage pregnancy and AIDS. Presently, there is limited research on the health status and health care needs of adolescents, particularly those who are difficult to track and quantify. The objective is to carry out a situation analysis of the health needs of adolescents, including the availability of existing services for them. The study setting is Klong Toey, the biggest slum community in Bangkok, where 38 adolescents were selected for focus group discussion, including in-depth interviews of youth leaders, government and non-government officers and schoolteachers, who work with adolescents. The result showed that adolescents in Thailand are older than traditionally defined. The culture and life-style of Thai society influence adolescents to stay with their families longer. Most adolescents do not have adequate and appropriate knowledge about sex and sexual relationships. The most important health problems are drug dependency and unwanted pregnancy among schoolgirls, which have commonly led to illegal abortions, and subsequent termination of schooling due to regulations and social norms. For general illness, such as pain associated with menstruation, headache and common cold, most have resorted to

---

Correspondence: R. Somrongthong and S. Sitthi-amorn, The College of Public Health, Chulalongkorn University, 10<sup>th</sup> Floor, Institute Building 3 Phayathai Rd., Bangkok 10330, Thailand. Fax: (662)-255-6046; Email: sratana3@chula.ac.th

self-medication with drugs obtained from the local drugstore. Males tend to acquire injuries from violence, accidents and drug abuse. The need for female adolescents is a clinic with female staff who will be able to provide both consultative and curative care, while the need for males is qualified staff for consultation, especially regarding drug abuse. Adolescents under-utilize existing government health services due to their unavailability and adolescents' dissatisfaction with their services. This study was conducted in only one community, however, the findings are important for the planning and delivery of appropriate and effective health services for young people living in urban slums.

## BACKGROUND

Although adolescents are often considered to possess good physical health, it is a period of physical, social and psychological change. The socio-cultural, political, religious environments and the economic conditions in which adolescents develop, have a strong influence on their development and health in later life. Due to development and rapid social change, the health problems of youth have dramatically shifted in the last 30 years from biological to social causes of morbidity and mortality. In fact, the literature and experience have indicated that adolescents' health problems have become serious, particularly in the developing countries, and require prompt action. This has resulted in an increasing volume of literature relating to the development of adolescent health services that meet the needs of their target group. However, a lack of understanding and a lack of emphasis on adolescent needs in policy and programming continues, especially in Asia. The outcome is that many still have limited access to health care, information concerning their growth and development, or counseling services. Even when health services do exist, they may be culturally inappropriate, insensitive to adolescent health needs, or delivered by staff with insufficient training in adolescent health. Thus, adolescents are unlikely to use them. This is alarming in a time of increased sexual activity among unmarried adolescents, a younger age at first intercourse, and evidence of an increasing incidence of STDs and HIV/AIDS in this group. Risk behaviors are also heavily concentrated in youth, leading to more unprotected sex, teenage pregnancies, substance abuse, and their consequent negative health effects.

In Thailand, many studies have shown that adolescents experience similar risks to the ones outlined above. For example, of the 22,064 drug addicts treated at rehabilitation clinics in Thailand, 24% of them were adolescents. This study also states that only 35.7% of adolescent males' first intercourse is protected, while only 25.9% of females' is protected. In addition, 41.2% of all AIDS patients in Thailand are 15-29 years old and 50.4% of all HIV+ people fall into this category. There is resistance to sex education by many Thai adults, as they still consider that it will probably lead to increased sexual activity among adolescents. These studies, among others, point to the need for appropriate adolescent health programs, especially in a time of economic crisis where less money is being spent in the public sector. It is often adolescents and young people who suffer from budget and service cuts. Such programs should be able to learn and reflect the needs of adolescents, acquire their help in planning and delivery, have the support of local leaders and parents and be trusted by the adolescents themselves. They must be culturally appropriate and institutionalized to ensure support and cost-effectiveness. Most studies focus on youth and adolescents in general and very few studies have targeted disadvantaged adolescents, particularly those living in slum communities. As so little is known about the specific problems and strengths unique to this group, this project was initiated. It is hoped that it will result in a better understanding of adolescents' health needs and behavior so that appropriate and accessible health services can be provided to this often overlooked group.

Our study was conducted in Klong Toey slum, the biggest slum community in Bangkok, the capital of Thailand. The area is about 3 square km., encompassing 25 different slum communities with a population of more than 70,000. There are around 17,000 inhabitants who are aged between 12-22 years. The community was first established in 1952 as a dwelling place for port workers and other laborers. Although many residents have lived there all their lives, the majority of them have no rights to the land they occupy, which creates a feeling of instability among the populace. Most people are laborers or unskilled workers. The biggest problems are poverty, lack of job security, limited education, and debt. In terms of health, the community suffers from overcrowding, unhygienic conditions, widespread drug and alcohol abuse, lack of clean drinking water, and broken families. There is a private doctor's clinic within the slum and a health center run by the Bangkok Metropolitan Authority.

## OBJECTIVES

To carry out a situation analysis of the health needs of adolescents, including the availability of existing services for adolescents in an under-privileged community of Bangkok. The information is needed to develop actions and further research to increase access to health care for adolescents, formulate the role of local and the central government in adolescent health, as well as to propose ways to improve adolescents' social environment, to promote healthy adolescent lifestyles.

## METHODOLOGY

A group of adolescents was assembled for focus group discussions, to understand their needs (health and non-health), their knowledge of, and access to, existing services, as well as their perception of the adequacy and quality of those services. Thirty-eight adolescents, aged between 12-23 years, were gathered for focus group discussions. Two groups were male and two groups were female. Each group consisted of 8-10 participants. They were selected by purposive sampling, i.e., with the help of the youth leaders in each of the 18 sections within the slum community. Two adolescents represented each section. The discussion followed a pre-prepared interview guideline plus observation of the behaviors of the adolescents, including their interaction and physical characteristics, during the discussion sessions. The product of the discussion was to be the variables that were to be used to construct an interview guideline.

In-depth interviews with 10 youth leaders, 4 health providers of the health center in the community and 4 non-government organization staff who were working with adolescents, were conducted in the community. The objective of the in-depth interviews was to understand the perception of each individual group, related to adolescents' health problems, actual utilization of the health services by adolescents, and the needs for health services.

Organizing the meeting among stakeholders and relevant agencies; a group of 21 individuals from various agencies in Thailand was gathered at the Health System Research Institute to discuss potential research activities pertaining to adolescent health. These included the Health System Research Institute, Mahidol University, the Department of Mental Health, the Division

of Family Health and Population, the Division of Health Promotion, the Institute of Health Research, the Bangkok Metropolitan Administration, the Department of Obstetrics and Gynecology at Chulalongkorn Hospital, the Division of School and Youth Health, private sector organizations dealing with youth, adolescents and AIDS, and the College of Public Health, Chulalongkorn University. The experiences and the activities of these agencies related to the adolescents' health were summarized. It was agreed that the College of Public Health, Chulalongkorn University, would serve as an information center for adolescent health. It was also agreed that periodic meetings should be organized to exchange resource persons, research results and other information about adolescent health.

## RESEARCH FINDINGS

### Focus Group Discussions among the Adolescents

The main objectives of focus group discussions were: 1) to understand the perceptions of the health problems of adolescents in the slum community, and 2) to understand their perceived needs for health services. The discussions followed a pre-prepared interview guideline plus observation of the behaviors of adolescents, including their interactions and physical characteristics during the discussion sessions.

#### *Findings:*

Most of the female adolescents were well-behaved, well-dressed, clean, pleasant and appeared to be happy. One member of the female group was a little reserved and silent (It was later reported by an officer from a private organization that she might be an addict and her household had been involved in illegal drug trafficking.) All females in the groups were unmarried. Three members came from broken families (the parents were separated).

Most of the male adolescents in the discussions were happy and appeared to be healthy. They were not as well-dressed as the females and used the language of their peers. One of the members was a heroin addict who said that he had already given up the habit. He appeared to be sick and unhealthy. Most of the males had had some sexual experiences. One of the members had

already had a family and was divorced.

*The major health problems of the female adolescents were:*

1. Pain associated with menstruation. Most had taken care of this problem themselves by using a rubber hot-water bottle and common analgesics.
2. General illness, such as headache and common cold. Most had resorted to self-medication with drugs obtained from local drugstores.
3. More than half of the female adolescents had used cleaning agents to clean their sex organs. Information about these products had been obtained from advertisements in the mass media and from various direct sales. Friends and close relatives had been important sources of information as well. Various cleaning agents had been used, particularly during menstruation.
4. Most had taken care of their external physical appearance, especially weight control. However, it had been difficult to control their eating habits.
5. Most had taken particular care of their skin complexion and the avoidance and treatment of acne.
6. Some had had boyfriends and all denied having ever had pre-marital sex.
7. Friends had been the main source of consultation about sex and sex organs. Some had had bad experiences consulting mothers who had tended to suspect an association between the questions about sex and sex organs with hidden sexual relationships with men.
8. Some of the adolescents had become metamphetamines addicts because their households had been involved in drug pushing (methyl amphetamines). Some had been selling the drugs to friends in their schools.
9. About one-third of the females in the focus groups knew about the use of "Postinal" for the prevention of pregnancy. The information had been obtained from friends with experiences and from married neighbors. Most had not had any sexual experiences.
10. Some had used "Dian" to increase the size of their breasts. The information about the drug came from friends.
11. Some had taken "Frimelude" to delay their menstruation for special occasions, such as if they planned to go for a picnic up-country.

12. Most had not dared to experiment with illegal drugs. Only two smoked occasionally.
13. Most of the stressful problems included problems with boyfriends, girlfriends and performance at school. Most had spent time thinking about the problems during their walks at shopping centers.
14. Everyone knew about AIDS but thought that it was not a problem since most did not indulge in drugs and extra-marital sexual relationships.
15. Mothers had been the most frequent consultants for general illnesses, friends or elder sisters for sexual organs and sexual relationships, and classmates for problems with performance in school.
16. Most would wait for minor illnesses to disappear and would buy their own drugs if the symptoms did not disappear.

***The major services used by the female adolescents were:***

1. Private clinic services, which were preferred due to their efficiency.
2. BMA Health Stations, which had not provided satisfactory services and could consume a lot of time.

***The major perceived health service needs for female adolescents were:***

1. A clinic with female service providers who could give consultation about various female health problems.
2. A consultation services for family and other problems, in addition to health problems.
3. A fitness club for physical exercise and weight control.

***The major health problems for male adolescents were:***

1. Most minor ailments were not problems since males should be perceived as being strong and should not worry about minor illnesses. If the illnesses persisted, most had resorted to self-medication.
2. Most of the problems had been due to accidents and injuries at the workplace.
3. Most male adolescents had had some involvement in community activities, such as being volunteers for the prevention of fire and arson.
4. Most had not had any opportunity for formal education. Most had enrolled in schools for adults and thus had had ample time during the

day. Most had lived with their parents. Only one lived on his own. Most had sought occasional work at a private organization in the community and received daily wages.

5. Questions about sex and sex organs had been discussed among friends, peers and close relatives, more than with parents or other household members.
6. Most had had some experiences with cigarettes, alcohol, and some illicit drugs. Some had been addicted to heroin but reported having quit the habit already. The opportunity to work in a private non-government organization was cited as the main reason for quitting drugs. Their work had kept them from the drug environment in the community.
7. Masturbation was cited as a way to reduce sexual desire and most considered masturbation as a natural activity, which did not incur any feeling of guilt.
8. Some had had accidents and injuries, including motorcycle accidents.

***The major uses of health services among the male adolescents were***

1. Most had not used BMA Health Services because of unfriendly attitudes towards the officers of the health station. The health officers had been perceived as being stern and not willing to spend adequate time to care for their problems. Most had preferred self-treatment.
2. Health services in schools had had limited capacity for only minor ailments. Most had not used school health services.
3. Most male adolescents had had experiences with females, such as their sweethearts, girlfriends, prostitutes and others. Most had not used condoms, especially with sweethearts and friends. As for prostitutes, they had chosen only those who "looked clean" and often had requested their partners to clean up before intercourse.

***The major perceived health service needs for male adolescents included:***

1. Community health services capable of giving advice about drug addiction, including treatment. The services should be appropriately distributed at various sites.
2. Periodic focus group discussions about the problems and experiences of male adolescents.

3. Adequate time for health providers to give adequate consultation.
4. A comprehensive set of services including consultation, referral and treatment.

### **In-Depth Interviews**

#### ***In-depth interview of staff of the private organization working with adolescents:***

**Findings:** It was perceived that most adolescents did not have adequate appropriate knowledge about sex and sexual relationships. They did not quite understand the physiological changes associated with adolescence. They had little idea about appropriate health promotion and care for their sex organs, due mainly to natural socialization among their peers. Most of the adolescents' problems related to their family, individual problems, problems with friends, problems with boyfriends and girlfriends. Some had problems with their studies. Many experienced mental and emotional stresses.

It was perceived that the health problems of the adolescents were not significant. This might be due to the nature of the services given by the private organization in question, which dealt mainly with giving consultations, rather than health services in general. Many of the adolescents preferred to seek consultations from someone they did not know, rather than their parents.

The executive of the organization thought that care for adolescents was important and that the efforts in Thailand had been hampered by discontinuity, due to a lack of a steady flow of funds, and competent human resources who had adequate experience working with adolescents. It is possible that the lack of concern for adolescent health may stop the flow of future funds.

#### ***In-depth interview of officers in an organization in the slum community:***

The **objectives** of the interviews were: 1) to understand the social and environmental situations and other problems in the community; 2) to get access to the organizations within the community who had worked with adolescents; and 3) to document the perceptions of these officers about the needs for health services for adolescents in the community.

**Findings:** It was found that of all the 18 communities in Klong Toey, drug dependence was perceived as the most important health problem. Some communities had been active in the promotion of illicit drugs. Many adolescents in the slum had not had an opportunity to continue their formal education due to poverty. Many had been striving to study in adult informal education and thus had had ample time to be idle and become involved with drugs.

Besides drug dependency, pregnancy among schoolgirls had been common, leading to illegal abortions and subsequent termination of schooling. These girls had confided with their mothers about the unwanted pregnancies. Most of the unwanted pregnancies had ended up in separation of the girls and their sexual partners.

AIDS and HIV had been common due mainly to needle sharing. However, asymptomatic HIV subjects had had frequent sex with changing partners. Most of the adolescents who had sex with HIV subjects had thought that they had little likelihood of being infected with the virus. This erroneous perception among adolescents is a cause for concern, and must be changed.

#### ***In-depth interviews of youth leaders in the community:***

The **objectives** of the interviews of community youth leaders were: 1) to understand the perceptions of the leaders about the health problems of adolescents; 2) to understand their needs for health services; and 3) to understand the actual utilization of health services by the adolescents.

**Findings:** The nature and extent of health problems among the adolescents in the slum community depend on the sex of the adolescents. Males tended to acquire injuries from violence and accidents, while females had been affected by general illnesses. Drug dependence had been a major problem among the males. Many of the addicts had come from families with drug dealers in the household. Some had had too much idle time, because they had been too poor to go to school. For women, teenage pregnancies had been common, leading to illegal abortion, premature marriage and eventual termination of formal education.

Health services are needed in the community. Consultation and advice about adolescents' health problems are most needed. Health services should be available 24 hours a day, since the problems associated with violence and injuries among males occur on a constant basis.

Most ad  
Metropolitan  
different pers  
staff and inac  
of the adolesc

#### ***In-depth inte (BMA) Healt***

The **objectiv**  
at the health  
utilization of  
problems and  
**Findings:** Th  
general publi  
rehabilitation.  
and child hea  
and care for  
provided to th  
10 beds for a  
there were no  
to adolescent  
illnesses. The  
had been insu

#### **CONCLUSIO**

This study w  
clients' perspe  
and these fac  
community, :  
adolescents. F  
general adoles  
community in  
study were ol  
cultural and li

Most adolescents do not use the services provided by the Bangkok Metropolitan Administration Health Center. The main reasons include different personalities between the providers and the adolescents, inadequate staff and inadequate time given to satisfactorily address the health problems of the adolescents.

***In-depth interview of staff of the Bangkok Metropolitan Administration (BMA) Health Station:***

The **objectives** of the interviews were: 1) to understand the services available at the health center for the general public and the adolescents; 2) the utilization of health services by the adolescents, and 3) the perceived problems and barriers in providing services for the adolescents.

**Findings:** The BMA Health Station had given comprehensive care to the general public such as disease prevention, health promotion, treatment and rehabilitation. The most common types of care sought by people were mother and child health services, treatment for general illnesses, care for the elderly and care for chronic diseases such as hypertension and diabetes. Care provided to the adolescents had been minimal. The BMA Health Station had 10 beds for admission. This was to be expanded in the future. At that time, there were no consultation services for the adolescents. Most of the care given to adolescents had been in the area of accidents, violence and general illnesses. The main barrier to the BMA Health Station in performing its tasks had been insufficient human resources.

## CONCLUSION

This study was conducted through situation analysis of the providers' and clients' perspectives of adolescents' health care needs and their accessibility, and these factors were ascertained through surveys of facilities, including community, school-based and hospital services currently available to adolescents. From qualitative data, conclusions have been drawn about the general adolescent health situation in the Klong Toey slum, the biggest slum community in Bangkok. It was perceived that the Thai adolescents in this study were older than the traditional definition, as a result of unique Thai cultural and lifestyle factors, which increase the period of adolescence. It was

felt that most of the adolescents did not have adequate or appropriate knowledge about their sex and sexual relationships. The most important health problems, as suggested by both health workers and adolescents, were drug dependency, unwanted pregnancy among schoolgirls (leading to subsequent termination of schooling), general illnesses such as pain associated with menstruation, headache, and the common cold. Most individuals tended to self-medicate at the local drugstore. Males reported their main health concerns were related to injuries acquired from violence, accidents and drug abuse. Needs for female adolescents were stated to be for a clinic with female staff who could provide both consultative and curative services, while males felt a need for qualified staff for consultations, especially regarding drug abuse. Health and fitness equipment was requested. The adolescents stated the reason for not frequenting existing government health services was that they felt the services provided were either inappropriate or irrelevant and they often felt dissatisfied with the treatment they received.

This study was undertaken in only one community, and these adolescents' problems may not be common to all in Thailand. However, the findings are important inputs to the planning and delivery of appropriate and effective health services to young people living in urban slums. Action research in the service sector is needed to develop and test prototypes for adolescent health services.

## REFERENCES

1. Kanungsukaseam A, et al. Sexual behaviour of young males in slum community (Thai version). Mahidol, Thailand: Institute of Population Study, Mahidol University, 1994.
2. Bennett DL. Worldwide problem in the delivery of adolescent health care. *Publ Health* 1989;96:334-40.
3. Curic CE, Todd J, Platt S. Indicator of socioeconomic status for adolescents: The WHO health behaviour in school-aged children. *Health Ed Res* 1997;12(3):385-97.
4. Diekstra RFW. Suicide behaviour in adolescents and young adults: The international picture *Crises* 1989;10 (1):16-35.
5. King A, Wold B, Smith TC, Yossi H. *The Health of Youth : A Cross –*

- National Survey. Copenhagen: WHO Regional Publ, European Series, 1996, 69.
6. Mench BS, Bruce J, Greene ME. The Uncharted Passage: Girls' Adolescence in the Developing World. Population Council, Inc. 1998.
  7. Verasakdi J, ed. The Proceeding of the Fourth National Meeting on Health Behaviours: Socio-Cultural Health (Thai version). Songkla, Thailand: Songkla University, 1997.

## **BITS 'N PIECES**

### **Association between postnatal catch-up growth and obesity in childhood: Prospective cohort study**

The objective of this study was to identify predictors of postnatal catch-up growth from birth to two years and its relation to size and obesity at five years. 848 full term singtons from a 10% random sample of the Avon longitudinal study of pregnancy and childhood were studied.

Overall, 30.7% (260 o 848) of infants showed a gain in SD score for weight greater than 0.67 SD scores between zero and two years, indicating clinically significant catch-up growth. These children had lower weight, length and ponderal index at birth than other children and were more often from primoparous pregnancies. They also had taller fathers than other children and their mothers had lower birth weights and were more likely to smoke during pregnancy. Children who showed catch-up growth between zero and two years were heavier, taller and fatter at five years than other children.

Mechanisms that signal and regulate early catxh-up growth in the postnatal period may influence associations between small size at birth and risks for disease in adulthood.

**Ong KKL, Ahmed ML, Emmett PM, Preece MA, Dunger DB. BMJ 2000;320:967-71.**

# FACTORS AFFECTING THE EXPERIENCES OF DRUG USE BY ADOLESCENTS IN A BANGKOK SLUM

Patrapan Laoniramai<sup>1</sup>, Orapin C Laosee<sup>2</sup>, Ratana Somrongthong<sup>3</sup>, Sunanta Wongchalee<sup>3</sup>  
and Chitr Sitthi-amorn<sup>2,3</sup>

<sup>1</sup>Faculty of Social Sciences, Kasetsart University; <sup>2</sup>Institute of Health Research, Chulalongkorn University; <sup>3</sup>College of Public Health, Chulalongkorn University, Bangkok, Thailand

**Abstract.** The purpose of this research was to study the demography, financial status, social status, knowledge of amphetamines, perceived harmfulness of amphetamines, and life skills in the prevention of drug abuse in adolescents. The factors leading to drug use among young people were also studied. The study group was composed of 354 subjects aged 12 to 22 years, living in 2 slums in Bangkok. The research showed that about 7% of the sample group had used drugs before. Four percent had never used drugs, but someone had tried to talk them into using them. Almost 20% had friends who had used drugs, and 11% had friends who were still using drugs. About 13% of the adolescents in the study group had family members who used drugs and another 9% had family members who were still using drugs. In our study, we found that the most common drug group was amphetamines. On average, the participants had a low level of understanding about drug abuse, especially of the symptoms, side effects, and legal penalties. Most of the adolescents realized how harmful amphetamines and other drugs were and had a high degree life skills. Factors influencing adolescent drug use were (1) personal factors, such as monthly income/allowance and life skills; (2) family environment, such as drug abuse history in the family; and (3) social environment, such as a drug abuse history among friends. When studying the life skill factors of the adolescents, which is an independent factor capable of influencing the experience with drugs, the researchers found that the time spent with other members of the family and the family members' drug experiences were the only factors leading to life skills in the prevention of drug abuse in adolescents. In addition to letting children learn on their own, training them to acquire life skills is beneficial when faced with problematic situations. Creating relationships between adolescents and other members of the family, friends, and society can increase their life skills, diminishing the risk of drug abuse.

## INTRODUCTION

Many researchers have sought the cause of drug abuse/addiction, so that effective treatments and preventive methods can be implemented. Research has found that the problem arises from many aspects: family sources, such as mother-father relationships, the way parents raise their children, the financial situation of the family, and the community, such as illicit gathering places, wrong or bad values, and neglect (Pisuttiwongse, 1996). The people adolescents spend time with also affect his/her habits. Adolescents who have friends using drugs are more

prone to become addicts themselves. Friends tend to introduce others to drug use (Adger, 1992). Another factor affecting drug abuse is the lack of knowledge of the drug's effects.

Because of the complexity of the problem, collaboration between several groups must be implemented to increase adolescents' understanding amphetamines and themselves, to enable them to develop self-worth and to prepare them to face drug temptations, learning to say "No" to drugs. Society should support these adolescents in preparing them for such situations. Friends, family, teachers and other members of the community should give support to adolescents so that they are emotionally and mentally ready to fight against drugs.

This research is part of a project called "Creative communication between friends to support drug (amphetamine) prevention habits among

Correspondence: Orapin C Laosee, Institute of Health Research, Chulalongkorn University, 4<sup>th</sup> Floor, Institute Building 2, Chulalongkorn Soi 62, Phya Thai Road, Pathum Wan, Bangkok 10330, Thailand.  
Tel: 66 (0) 2218-8208; Fax: 66 (0) 2253-2395  
E-mail: Orapin.c@chula.ac.th

adolescents: a case study of a Bangkok slum". The research investigated basic information about adolescents in Bangkok's slums as a basis for implementing activities supporting the prevention of drug abuse among young people.

## MATERIALS AND METHODS

This research was based on a survey. The sample group was composed of adolescents ranging in age from 12-22 years, living in Bangkok slums. A purposive sampling method was used to select 2 slums with different characteristics. Simple random sampling was used to select the 354 participants from the 2 communities. Data were collected by questionnaires filled out by the adolescents. A staff member was present to give assistance to the participants, such as explaining the meaning of the questions or how to fill out the questionnaire, and answer any concerns raised by the participants. Descriptive statistics were used to analyze the data collected. Frequencies, averages and standard deviations were calculated. Logistic regression analysis was used to study the factors influencing drug use habits.

## RESULTS

### Demographic, economic, and social characteristics

The sample group was comprised of more women than men, aged 15-19. Thirty percent of the participants' parents were separated. The parents were mostly employees or salespersons in street. They were mostly primary-school educated. Most of the subjects had finished middle school. The average family income was about 10,000 baht per month, providing for 5 family members. The adolescents received, on average, a 2,400-baht monthly allowance from their parents. The main problem the participants faced was financial. When any problems occurred, the participants turned to their parents first, then to their friends. The adolescents spent most of their time listening to music, watching television, and hanging out with friends in order to relax.

### The drug habits of adolescents and their close ones

The research shows that about 7% of the

sample group had used drugs before. Four percent had never used drugs but someone had tried to talk them into using them. Almost 20% had friends who had used drugs, and 11% had friends who were still using drugs. About 13% of the adolescents in the study group had family members who had used drugs and another 9% had family members who were still using drugs. In this study, the most common drug group used was amphetamines.

### Knowledge, perceived harmfulness, and promoting of life skills to prevent amphetamine addictive behavior

The adolescents had a low level of understanding regarding amphetamines, especially regarding their symptoms and side effects. Their main sources of knowledge were educators or schools. Nevertheless, these adolescents had high levels of perceived harmfulness of amphetamines, and had high score levels for life skills (Tables 1-3).

### Factors influencing adolescent drug use behavior

The personal factors influencing the drug habits of adolescents were the amount of money the adolescent received per month, and the life skills promoting amphetamine-prevention attitudes. Adolescents with no financial problems were less likely to use drugs. Adolescents with skills to prevent drug abuse were at lower risk for drug use. The family environmental factor that influenced adolescent drug habits was the experience of drug use in a family member. An adolescent who had a family member who used drugs was 1.4 times more likely to have used them than an adolescent who had no family member using drugs. Previous drug use by friends, being a social factor, also increased the risk of adolescent drug habits. In other words, an adolescent who had a friend using drugs was about 1.5 times more likely to use drugs than someone who did not have a friend using drugs (Table 4).

### Factors influencing adolescent life skills

On additional analysis, an independent factor, namely life skills, also influenced the drug experience of adolescents. The research found that the family environment, which included

Table 1  
Percentage of adolescents classified by score level of knowledge of amphetamines, and their experience using drugs.

Score level for knowledge of amphetamines	Experience using drugs		Total
	Never	Experienced	
Low level of knowledge	43.0	33.3	42.4
Moderate level of knowledge	45.5	62.5	46.6
High level of knowledge	11.5	4.2	11.0
Total	100.0	100.0	100.0
(Number)	(330)	(24)	(354)
Average score	3.99	3.96	4.02

Table 2  
Percentage of adolescents classified by score level for perceived harmfulness of amphetamines, and their experience using drugs.

Score level for perceived harmfulness of amphetamines	Experience using drugs		Total
	Never	Experienced	
High level	69.7	45.8	68.4
Rather high level	22.7	41.7	23.8
Moderate level	5.8	8.3	5.8
Rather low level	1.8	4.2	2.0
Total	100.0	100.0	100.0
(Number)	(330)	(24)	(354)
Average score	4.21	4.01	4.20

Table 3  
Percentage of adolescents classified by score level for life skills promoting amphetamine-prevention attitudes, and their experience using drugs.

Score level for life skill	Experience using drugs		Total
	Never	Experienced	
Low level of life skills	2.8	8.7	3.2
Moderate level of life skills	37.4	60.9	39.0
High level of life skills	59.7	39.0	57.9
Total	100.0	100.0	100.0
(Number)	(318)	(23)	(341)
Average score	25.53	23.65	25.19

spending time with family members and the drug experiences of family members, was the only factor influencing adolescent life skills: the more time spent with their family, the higher the level of life skills. Someone who had a family member with a drug history would have a higher level of life skills than someone who did not (Table 5).

## DISCUSSION

Even though most of the participants had never used drugs before, some of them were still at great risk for drug use due to a family drug history. The most dangerous was when a friend in the group was using drugs, because adoles-

Table 4  
Logistic regression analysis results of factors influencing adolescent drug use experiences.

Factors	B	SE	p-value	Exp(B)
<b>Personal factors</b>				
Gender (female <sup>a</sup> )	1.205	0.693	0.082	3.335
Age	0.124	0.143	0.389	1.131
Income	-0.001	0.000	0.019 <sup>b</sup>	0.999
Knowledge	-0.013	0.185	0.944	0.987
Perception	0.033	0.031	0.291	1.034
Skill	-0.228	0.097	0.018 <sup>b</sup>	0.796
<b>Family environment factors</b>				
Number of family members	-0.113	0.136	0.408	0.894
Marital status of parents (separated <sup>a</sup> )	1.319	0.785	0.093	3.740
Warmth	0.098	0.471	0.674	1.219
Restriction	-0.209	0.345	0.544	0.811
Quarreling	0.287	0.357	0.421	1.332
Conflicts	-0.003	0.425	0.994	0.997
Concern	-0.726	0.412	0.078	0.484
Spending time	0.140	0.325	0.666	1.151
Drug use experience in a family member (No family member experienced drug use <sup>a</sup> )	-1.486	0.662	0.025 <sup>b</sup>	0.226
<b>Socio-environmental factors</b>				
Community (participation <sup>a</sup> )	-1.086	0.851	0.202	0.338
Education (Vacation <sup>a</sup> )	0.495	0.939	0.598	1.641
Participation in drug activity in the community (never <sup>a</sup> )	0.912	0.793	0.250	2.489
Drug use experience among friends (never <sup>a</sup> )	-1.534	0.754	0.042 <sup>b</sup>	0.216

<sup>a</sup>Control group; <sup>b</sup>p<0.05

B = Unstandardized regression coefficient; Exp(B) = Exponential of B; SE = Standard error of B

cents bestow a great deal of importance upon friends, putting the subject at increased risk for adolescent drug abuse. Suphap (1997) stated that adolescent behavior was greatly influenced by the behavior of members of the group or the group leader. Analysis of the factors influencing adolescent drug habits (mostly amphetamine addiction) showed that family or friends' drug habits do influence adolescent risk of drug use. This is consistent with the results of the research by Sussman *et al* (2000), which found that the drug use habits of friends were a predictive factor for adolescent drugs habits. Adolescents were more likely to be influenced if a member of the family, especially a parent, had a history of drug use. Adger (1992) found that the family factor influences the probability of alcohol and drug abuse by an individual. For instance, if a father

drank alcohol, his son was more likely to drink alcohol. An adolescent who had a drug-addicted friend was more likely to use drugs than someone with no friends who use drugs.

Adolescent culture revolves around friends. They are easily influenced by friends because of the need for peer recognition. Being shunned and isolated from the group is considered humiliating and the worst punishment. Curiosity increases the risk of drug abuse. To prevent the occurrence of such problems, a new referral group needs to be established and friends need to draw adolescents away from drugs.

Having a family member or friend who has used drugs does not mean that all adolescents will follow in his/her footsteps. Adolescent life skills also play an important role here. The research indicates that the higher the life skill level,

Table 5  
Multiple regression analysis results of factors influencing adolescent life skills promoting drug-prevention behavior.

Factors	B	SE	Beta	t	p-value
<b>Constant</b>	22.794	2.356		9.675	0.000 <sup>b</sup>
<b>Personal factors</b>					
Gender (female <sup>a</sup> )	-0.056	0.379	-0.010	-0.150	0.881
Age	-0.089	0.069	-0.100	-1.312	0.191
Income	-0.000	0.000	-0.003	-0.043	0.966
Knowledge	0.029	0.090	0.022	0.332	0.740
Perception	0.016	0.016	0.069	1.060	0.290
<b>Family environment factors</b>					
Number of family members	-0.011	0.063	-0.011	-0.173	0.863
Marital status of parents (separated <sup>a</sup> )	-0.464	0.392	-0.008	-0.118	0.906
Warmth	0.176	0.257	0.056	0.685	0.494
Restriction	0.079	0.211	0.025	0.376	0.707
Quarreling	0.093	0.194	0.034	0.477	0.634
Conflicts	0.037	0.218	0.012	0.169	0.866
Concern	0.117	0.248	0.039	0.471	0.638
Spending time	0.412	0.178	0.159	2.311	0.022 <sup>b</sup>
Drug use experience in a family member (No family member experienced drug use <sup>a</sup> )	0.918	0.421	0.140	2.179	0.030 <sup>b</sup>
<b>Socio-environmental factors</b>					
Community (participation <sup>a</sup> )	-0.057	0.379	-0.010	-0.150	0.881
Education (Vacation <sup>a</sup> )	-0.793	0.467	-0.111	-1.697	0.091
Participation in drug activity in the community (never <sup>a</sup> )	0.379	0.358	0.068	1.057	0.292
Drug use experience among friends (never <sup>a</sup> )	-0.496	0.399	-0.081	-1.243	0.215

$R^2 = 0.112$ ;  $F = 1.800^b$

<sup>a</sup>control group; <sup>b</sup> $p < 0.05$

B = Unstandardized regression coefficient; SE = Standard error of B

Beta = Standardized regression coefficient; t = t value for B

R = Correlation coefficient;  $R^2$  = Coefficient of determination; F = F value for regression model

the lower the risk of drug abuse. These skills include life adaptations, problem solving, communication, negotiation, analysis, reasoning, and coping with stress, especially when faced with peer pressure. Adolescents acquiring the latter skills can prevent drug abuse (Lapsirianankul, 2000).

To acquire such skills, adolescents need to learn from life experiences, and from their interactions with other members of society. Our study shows that monthly allowance also influences adolescent drug habits. Participants with no financial worries were less likely to use drugs. This does not mean that parents or guardians must

increase their children's allowances. Teaching financial values to adolescents can minimize financial problems.

Forty-two percent of participants had a low level of understanding regarding amphetamines, while 46.6% had a moderate level. Schools were the primary source of information regarding amphetamines and other drugs. The results of the study reflect those by Varalakshna (1995), which indicated that educators were the ones who really understood the threat of drugs and who communicated these risks to the students. One problem is that the age range of students influenced by drugs now starts in primary school.

Thus, drug awareness programs to prevent drug abuse need to be implemented in primary school age children. Even though adolescents had a low level of understanding of amphetamines and their side effects, the majority of them (68.4%) did comprehend the power of drugs over people. This can be considered a first step in preventing most young people from using drugs.

The study also shows that an important factor influencing life skill levels is the family environment. This means that the family is the source of life skills acquisition among adolescents, from spending time with members of the family. The more time spent with the family, the greater the learning process from parental experience and human interaction. These activities shape the values and behavior of adolescents. However, this can also be harmful to adolescents, since a family drug history can negatively influence children. This is consistent with a study by Low *et al* (1996), which found that being beyond the concern of the family can lead to drug use in adolescents. Nevertheless, family drug experience increases the life skills of those who have experienced them, compared with those who have not. This may be because they have faced drug problems before, and learned to say "No" to drugs, developing the necessary life skills.

This does not mean that people with no family drug experience have fewer life skills, since these skills can be acquired by one's own experience, and through interaction with other members of society in many different situations. Drug awareness activities can also be a method of teaching adolescents to understand the harm of using drugs and how to prevent such situations. Although a one-day, two-day, or one-week drug awareness activity cannot develop immediate skills for drug prevention, it can be combined with other life experiences to prevent the influence of drugs. Ko (1997) found that youth camps can be used to inform and educate adolescents about drugs and how to prevent drug use. Thus, a shield against amphetamines and other drugs can be achieved with close collaboration between the adolescent, parents, friends, educa-

tors, the community, and society.

## ACKNOWLEDGEMENTS

We thank the Thailand Research Fund (TRF) for funding this study. We wish to acknowledge Duang Prateep Foundation's staff for their kind assistance through this project.

## REFERENCES

- Adger Jr H. Alcohol and other drug use and abuse in adolescents. Adolescents at risk: medical and social perspective. Seventh Conference on Health Policy. Cornell University, Medical College: Westview Press, 1992.
- Ko P. The effectiveness of health education program on amphetamine preventive behaviors of Mathayom Suksa 1 students in Phra Nakhon Si Ayutthaya Province. Bangkok: Srinakharinwirot University, 1997. MS Thesis. 239 pp.
- Lapsirianankul J. The development of life skills for prevention of amphetamine use among primary school students through student club in Ban Pong district, Ratchaburi. Nakhon Pathom: Mahidol University, 2000. MS Thesis. 137 pp.
- Low WY, Zulkifli SN, Yusof K, Batumalail S, Aye KW. The drug abuse problem in Peninsular Malaysia: parent and child differences in knowledge, attitudes and perceptions. *Drug Alcohol Depend* 1996; 42: 105-15.
- Pisuthiwongse S. The participation of the community leaders in the prevention of amphetamine problem in Nakhon Pathom Province. Nakhon Pathom: Mahidol University, 1996. MA Thesis. 112 pp.
- Sussman S, Dent CW, Leu L. The one-year prospective prediction of substance abuse and dependence among high risk adolescent. *Subst Abuse* 2000; 12: 378-86.
- Suphap S. Sociology, 16<sup>th</sup> ed. Bangkok: Thai Watana Panich, 1997.
- Varalaksna V. Mass and interpersonal communication exposure on narcotics and the prevention among students of the lower secondary education in Muang District, Chiang Mai Province. Bangkok: Chulalongkorn University, 1995. MA Thesis. 170 pp.

# PATTERNS, APPROPRIATENESS, AND PREDICTORS OF ANTIMICROBIAL PRESCRIBING FOR ADULTS WITH UPPER RESPIRATORY INFECTIONS IN URBAN SLUM COMMUNITIES OF BANGKOK

Siritree Suttajit<sup>1</sup>, Anita K Wagner<sup>2</sup>, Ruangthip Tantipidoke<sup>3</sup>, Dennis Ross-Degnan<sup>2</sup>  
and Chitr Sitthi-amorn<sup>4</sup>

<sup>1</sup>College of Public Health, Chulalongkorn University, Bangkok; <sup>2</sup>Department of Ambulatory Care and Prevention, Harvard Medical School and Harvard Pilgrim Health Care, Boston, MA, USA; <sup>3</sup>Faculty of Pharmaceutical Sciences, Chulalongkorn University, Bangkok; <sup>4</sup>Institute of Health Research, Chulalongkorn University, Bangkok, Thailand

**Abstract.** Upper respiratory tract infections (URIs) are the most common infections worldwide. Their frequent inappropriate treatment with antibiotics is likely to increase antibiotic resistance, contribute to morbidity and mortality, and waste scarce resources. Using data from registration books and prescriptions, we measured patterns and assessed appropriateness and predictors of antibiotic prescribing for viral and bacterial URIs treated in health centers located in two slum communities in Bangkok, Thailand. Based on recorded diagnoses and symptoms, 91% of the patients probably had viral URIs; 60% of viral and 89% of bacterial URI patients were prescribed an antibiotic. Compliance with the national treatment guideline was 36.4% for treatment of viral URIs and only 1.7% for treatment of bacterial URIs. Amoxicillin was the most frequently prescribed antibiotic regardless of diagnosis. Among viral URI patients, those who were young, male, and self-paying were more likely to receive antibiotics; part-time physicians were more likely to prescribe antibiotics for these patients. Among patients with bacterial URIs, those who paid for drugs by themselves were more likely to receive antibiotics compared to patients covered by the national health insurance plan. We used these formative results as input to the design of health center and community interventions to encourage more appropriate prescribing for URI among adults.

## INTRODUCTION

Upper respiratory tract infections (URIs) are the most common infections worldwide (Bamberger and Jackson, 1995). In Thailand, acute URIs are an important health problem. Nearly half of respondents to the 2000 National Health and Welfare Survey (NSO, 2000) reported having had a respiratory disease during the two preceding weeks, the most frequently reported acute health problem.

Although viruses cause most URIs (Therapeutic Guidelines, 1998), antimicrobial agents are still among the most frequently prescribed

drugs for URI. Unnecessary antimicrobial use for URI, both at health facilities and in the community, is one of the most common drug use problems (Grand et al, 1999). Adverse consequences of antimicrobial use for viral URIs are particularly unfortunate in developing countries (Kunin, 1993). Since most patients pay out-of-pocket for their medicines, they waste scarce household resources on unnecessary care. Moreover, inappropriate antimicrobial use increases drug resistance in the community, requiring use of more expensive and potentially more toxic alternative drugs when antimicrobial treatment is needed. These more expensive antimicrobials are less likely to be affordable in resource-poor settings. In Thailand, data from the Drug Control Division (2003) showed that antimicrobials are the most frequently used drugs and their use increases every year.

---

Correspondence: Siritree Suttajit, College of Public Health, Chulalongkorn University, Bangkok 10330, Thailand.  
Tel: +66 (0) 2218 8198; Fax: +66 (0) 2255 6046  
E-mail: siritree@gmail.com

According to the WHO Strategy for the Containment of Antimicrobial Resistance (WHO, 2001), developing countries play an important role in the emergence of antimicrobial resistance. In developing countries, bacterial resistance to antimicrobial agents is common in isolates from healthy persons (Okeke et al, 1999). In urban slum communities with poor sanitation, high incidences of HIV/AIDS, multi-drug resistant tuberculosis (MDRTB), and drug addiction, resistance is likely to spread easily.

Studies in developed countries suggest that both clinical and socio-cultural factors contribute to inappropriate antimicrobial prescribing (Avorn and Solomon, 2000; Murray et al, 2000; Pechere, 2001). A recent review by Radyowijati and Haak (2003) highlighted the lack of data to characterize the importance of different determinants of antibiotic use in developing countries. There are few studies of antimicrobial use for URIs in these settings and most report on treatment of URIs in children. Patterns of antimicrobial use in low-income countries differ from those in industrialized countries, and complex relationships exist between antibiotic use and local culture, socioeconomic characteristics, and poor health care infrastructure (Radyowijati and Haak, 2003).

To provide formative data to guide the design of interventions to improve treatment of URI in adults, we assessed patterns of antimicrobial and other drug treatment for adults with URIs treated in health centers in two slum communities in Thailand.

## MATERIALS AND METHODS

### Setting

We conducted the present study in two of the 61 health centers run by the Bangkok Metropolitan Administration (BMA), which provide care in under-served slum communities in Bangkok, Thailand. More than 125,000 individuals live in 22,063 households in a two square kilometer slum area covered by the two health centers. People in these communities face problems of substandard housing, poor sanitation, unemployment, low income, high incidence of HIV/AIDS, MDRTB, sexually transmitted diseases, and drug addiction.

### Data sources

We retrospectively identified in registration books of the two health centers all patients over 18 years of age who were treated for URI in 2001. We included cases with diagnoses of URI, cough and cold symptoms, pharyngitis, tonsillitis, otitis media, or sinusitis. We abstracted the prescriptions these patients had received, which were written by a total of 19 physicians.

Information about patients collected from registration books included age, gender, and diagnoses. Type, amount, and cost of prescribed drugs, type of payment for drugs, and physician name were collected from copies of the prescriptions maintained in the clinic pharmacies. Additional information about physician characteristics (age, gender, full-time or part-time employment status, specialty, and number of years practicing) was gathered directly from the physicians and from health center administrators.

The study was approved by the Human Research Ethical Committee at the Faculty of Medicine of Chulalongkorn University; the Ethical Review Committee for Research in Human Subjects, Ministry of Public Health, Thailand; and the Boston University Medical Center Institutional Review Board.

### Definition of 'likely viral' and 'likely bacterial' URIs

We classified patients diagnosed with URI or common cold, unspecified URI, or symptoms like cough, rhinitis or headache as having 'likely viral' URIs. Patients diagnosed with pharyngitis, sore throat, tonsillitis, acute otitis media, or sinusitis were classified as having 'likely bacterial' URIs. Patients with more than one URI diagnosis were classified according to the most severe diagnosis.

### Definition of appropriate treatment

We classified treatments according to the National Standard Treatment Guideline for Respiratory Tract Infections (MoPH, 1996). For viral URIs, such as nonspecific URI, acute bronchitis, rhinitis, or influenza, the Guideline emphasizes that antibiotics should not be given unless patients have high fever, severe cough, or abscesses. For bacterial URIs, the Guideline recommends penicillin V, penicillin G, or erythromycin for 7-10 days for pharyngitis and tonsillitis;